



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>


LANE MEDICAL LIBRARY STANFORD  
J57 G368 1905 STOR  
Post-mortem pathology : a manual of post



24503312833



# PLATE I



Drawing showing the normal situation of the parts of the human trunk, introduced for the purpose of quickly and accurately preparing outline figures to illustrate the situation and extent of lesions found at any postmortem. 1, esophagus; 2, left common carotid; 3, right subclavian artery; 4, first rib; 5, aorta; 6, sternum, with heart beneath; 7, adrenal; 8, pylorus; 9, gall-bladder; 10, ascending vena cava; 11, left common iliac; 12, ureter; 13, appendix; 14, vertebral column; 15, right common carotid; 16, clavicle; 17, left subclavian artery; 18, pulm-; 19, tricuspid valve; 20, thoracic duct; 21, tricuspid valve; 22, thoracic duct; 23, adrenal; 24, left sol-; 25, umbilicus; 26, small intestine; 27, right common; 28, right common; 29, ureter; 30, ant-; 31, lumbar vertebra; 32, ovaries; 33, oviducts.

*Dr. B. Wilbur*



Gift of Dr. Wilbur



# POST-MORTEM PATHOLOGY

A MANUAL OF POST-MORTEM EXAMINATIONS  
AND THE INTERPRETATIONS TO BE  
DRAWN THEREFROM

*A PRACTICAL TREATISE FOR STUDENTS AND PRACTITIONERS*

LANE BY GRARY

HENRY W. CATTELL, A.M., M.D.

AMERICAN EDITOR AND TRANSLATOR OF ZIEGLER'S SPECIAL PATHOLOGY; PATHOLOGIST TO THE WEST  
PHILADELPHIA HOSPITAL FOR WOMEN; AND SOMETIME DIRECTOR OF THE JOSEPHINE M. AYER  
CLINICAL LABORATORY OF THE PENNSYLVANIA HOSPITAL; SENIOR CORONER'S PHYSICIAN OF  
PHILADELPHIA; PATHOLOGIST TO THE PRESBYTERIAN AND PHILADELPHIA HOSPITALS;  
PROSECTOR OF THE AMERICAN ANTHROPOMETRIC SOCIETY; DEMONSTRATOR  
OF MORBID ANATOMY IN THE UNIVERSITY OF PENNSYLVANIA, ETC.

---

**SECOND, REVISED AND ENLARGED EDITION**

---

COPIOUSLY ILLUSTRATED WITH COLORED PLATES AND FIGURES

"Rotto dal mento insin dove si trulla.  
Tra le gambe pendevan le minugia;  
La corata pareva, e il tristo sacco  
Che merda fa di quel che si trangugia."

—DANTE

PHILADELPHIA AND LONDON  
J. B. LIPPINCOTT COMPANY

1905

d



VBA 981: 3BA

Copyright, 1903, by J. B. LIPPINCOTT COMPANY

Copyright, 1905, by J. B. LIPPINCOTT COMPANY

PRINTED BY J. B. LIPPINCOTT COMPANY, PHILADELPHIA, U.S.A.

J57  
C368  
1905



TO THE MEMORY OF  
MY FRIENDS

DR. THOMAS G. MORTON  
DR. THOMAS S. KIRKBRIDE, JR.

37520



## PREFACE TO THE SECOND EDITION



To this edition have been added a chapter on the pathology of the bones and joints, a summary of the literature on legal medicine and the technic of post-mortem examinations, and twenty-seven new illustrations, including six in colors. The chapters on bacteriologic investigations, medicolegal suggestions, and the examination of the exterior of the body have been thoroughly revised, and much other matter has been rearranged and wholly or in part rewritten.

I am indebted to Dr. ADELAIDE W. PECKHAM, Professor of Bacteriology in the Woman's Medical College of Pennsylvania, for suggestions regarding Chapter XXIII, Bacteriologic Investigations, and to Dr. ELLEN P. CORSON-WHITE for her assistance upon Chapter V, Examination of the Exterior of the Body, and upon the references contained in Chapter XXIX. Those who helped me in the preparation of the first edition of this work have again performed a similar service, for which I am grateful.

HENRY W. CATTELL.

3709 SPRUCE STREET, PHILADELPHIA, January 18, 1905.



## PREFACE TO THE FIRST EDITION



This book has been written for those who ought to make autopsies but do not and for those of whom such investigations are required, as medical students, hospital interns, and coroner's physicians. While it would seem to be quite needless to urge upon a practitioner the importance of performing post-mortem examinations, it is a fact that extremely few are made outside of hospitals, and even there necropsies are usually conducted by the untrained resident or the substitute of the pathologist. It cannot be questioned, however, that the physician who improves his opportunities for pathological study on the cadaver will be a better diagnostician and safer therapist, will have a more enduring reputation, and will receive a greater pecuniary return than he who neglects such means of investigating morbid processes.

While the author has mainly relied upon his personal experiences in the preparation of the subject-matter of this manual, he has freely used classifications and material derived from ORTH'S *Pathologisch-Anatomische Diagnostik*, OSLER'S *Practice of Medicine*, NAUWERCK'S *Sections-Technik*, and other publications mentioned in the foot-notes and in the text. He is, therefore, much indebted to these authorities, as well as to Dr. GEORGE ROBINSON and Mr. LOUIS SCHMIDT for most of the drawings, all of which were prepared under the writer's direction, to his friends and former students Drs. WILLIAM S. WADSWORTH, MARY E. LAPHAM, E. D. BURKHARD, and EDWARD LODHOLZ for suggestions in the preparation of the book, and to that excellent proof-reader Mr. T. GROW TAYLOR for seeing the work through the press.

HENRY W. CATTELL.

3709 SPRUCE STREET, PHILADELPHIA. March 31, 1903.

# CONTENTS



CHAPTER	PAGE
I. GENERAL CONSIDERATIONS .....	1
II. ORDER OF EXAMINATION AND POST-MORTEM RECORDS .....	13
III. POST-MORTEM INSTRUMENTS AND HOW TO USE THEM .....	28
IV. THE CARE OF THE HANDS AND THE TREATMENT OF POST-MORTEM WOUNDS .....	38
V. EXAMINATION OF THE EXTERIOR OF THE BODY .....	46
VI. TECHNIC OF OPENING THE ABDOMINAL CAVITY AND THE TOPOGRAPHIC EXAMINATION OF THE PARTS CONTAINED THEREIN .....	79
VII. TECHNIC OF EXPOSING THE THORACIC CAVITY AND THE CRITICAL EXAMINATION OF THE PARTS CONTAINED THEREIN .....	92
VIII. DISEASES OF THE HEART, BLOOD, BLOOD-VESSELS, AND LYMPH-VESSELS .....	113
IX. DISEASES OF THE RESPIRATORY TRACT AND ACCESSORY PARTS .....	141
X. CRITICAL EXAMINATION OF THE ORGANS OF THE ABDOMINAL CAVITY ..	159
XI. DISEASES OF THE GENITO-URINARY TRACT .....	199
XII. DISEASES OF THE LIVER AND PANCREAS AND THEIR DUCTS .....	213
XIII. EXAMINATION OF THE SKULL AND BRAIN .....	225
XIV. EXAMINATION OF THE SPINAL CANAL AND CORD .....	242
XV. DISEASES OF THE BRAIN AND CORD .....	246
XVI. EXAMINATION OF THE NASOPHARYNX, EYES, AND EARS .....	256
XVII. EXAMINATION AND DISEASES OF THE BONES AND JOINTS .....	261
XVIII. POST-MORTEM EXAMINATION OF THE NEW-BORN .....	275
XIX. RESTRICTED POST-MORTEM EXAMINATIONS .....	280
XX. RESTORATION AND PRESERVATION OF THE BODY .....	283
XXI. DISEASES DUE TO MICRO-ORGANISMS, PARASITES, AND HÆMATOZOA ...	290
XXII. THE PRESERVATION OF TISSUES FOR MACROSCOPIC AND MICROSCOPIC PURPOSES .....	326
XXIII. BACTERIOLOGIC INVESTIGATIONS .....	346
XXIV. WEIGHTS AND MEASURES .....	357
XXV. COMPARATIVE POSTMORTEMS .....	373
XXVI. MEDICOLEGAL SUGGESTIONS .....	397
XXVII. THE PRUSSIAN REGULATIONS FOR THE PERFORMANCE OF AUTOPSIES IN MEDICOLEGAL CASES .....	436
XXVIII. USUAL CAUSES OF DEATH; THEIR NOMENCLATURE, COMPLICATIONS, AND SYNONYMS .....	448
XXIX. REFERENCES .....	465
XXX. GLOSSARY INDEX .....	489



# LIST OF ILLUSTRATIONS



PLATE	PAGE
I. Outline Chart of Human Body .....	<i>Second page of cover</i>
II. Normal Size of Familiar Objects .....	18
III. Post-Mortem Scale for estimating Hæmoglobin. Moisture Scale.	
	<i>Third page of cover</i>
IV. Chart of Tumors .....	134
V. Bacteriologic Chart .....	350
VI. Powder Markings .....	409
FIG.	
1. Portable post-mortem table .....	8
2. Body in coffin prepared for post-mortem examination .....	8-9
3. Post-mortem table for babe, constructed of chairs and drawing-board ..	8-9
4. Post-mortem room of Ayer Clinical Laboratory, Pennsylvania Hospital	8-9
5. Refrigerator box for preservation of bodies .....	9
6-8. Working plans for preparing refrigerator box for storage of bodies ....	10
9, 10. Plans for post-mortem table .....	10-11
11, 12. Working plan and drawing of combination electric, gas, and water fixtures for post-mortem table .....	11
13. Section- or cartilage-knife .....	28
14. Cartilage-knife with projection on back .....	28
15. Post-mortem knife with a faulty point and improper belly .....	28
16. Coplin's brain-knife .....	28
17. Bread-knife for incising large organs .....	28
18. Valentine's knife .....	28
19. Pick's myelotome .....	28
20. Forceps; saw; needles; hammer; steel tape measure; hone and strop; and rounded hard-wood handle for scalpel .....	28-29
21. Saw for post-mortem work .....	28-29
22. Butcher's saw for post-mortem work .....	28-29
23. Hey's saw .....	28-29
24. Metacarpal saw .....	28-29
25. Luer's double rhachiotome .....	28-29
26. Cryer's electric surgical engine .....	30
27. Strong scissors with short blades .....	29
28. Scissors with one dull point and with bent handles .....	29
29. Separate-bladed scissors .....	29
30. Proper form of enterotome .....	29
31. Improper form of enterotome .....	29



FIG.	PAGE
32. Proper form of costotome .....	30
33. Improper form of costotome .....	30
34. Steel hammer .....	30
35. Steel side chisel .....	30-31
36. Curved chisel .....	30-31
37. Brunetti's curved spinal chisel .....	30-31
38, 39. Various forms of forceps .....	30-31
40. Straight grooved director .....	30-31
41. Satterthwaite's calvarium clamp .....	30-31
42. Iron clamp for removing brain .....	31
43. Bigelow clamp .....	31
44. Folding head-rest .....	31
45. Cornell folding clamp .....	32
46. Measuring stick .....	32
47. Raspatory .....	33
48. Pocket-case of instruments .....	32
49. Cones for measuring orifices .....	32-33
50. Glass balls for measuring orifices .....	32-33
51. Linen twine wrapped of proper length ready for use .....	32-33
52. Formad's pocket-case for post-mortem instruments .....	32-33
53. Method of holding cartilage-knife .....	33
54-56. Lines showing various methods for opening thoracic and abdominal cavities .....	79
57, 58. Method of making initial incision over sternum .....	78
59, 60. Method of removing skin over sternum and ribs .....	78-79
61. Method of opening abdominal cavity .....	78-79
62. Method of incising ribs .....	78-79
63, 64. Method of separating sternoclavicular attachment .....	79
65. Method of incising first rib and sternoclavicular articulation with costo- tome .....	92
66. Severance of diaphragm .....	92
67, 68. Removal of bony parts covering thoracic cavity .....	92-93
69. Breastplate after its removal from body .....	92-93
70. Methods of enlarging abdominal cavity and of protecting operator's hands from injury .....	92-93
71. Method of opening pericardium .....	93
72-74. Lines for opening heart .....	102
75. Method of removing heart from body .....	93
76. Situation of pulmonary veins .....	104
77. Method of opening right auricle .....	104-105
78. Method of opening left ventricle .....	104-105
79. Method of opening pulmonary artery .....	104-105
80. Interior of left auricle and ventricle fully exposed .....	105

# LIST OF ILLUSTRATIONS

xi

FIG.	PAGE
81. Reconstruction of heart after its examination .....	105
82. Methods of opening lung .....	108
83. Lung laid open for minute inspection .....	108-109
84. Method of opening pulmonary vein and its branches .....	108-109
85. Method of opening bronchi and their ramifications .....	109
86. Method of removing tongue, tonsils, œsophagus, trachea, etc., in a single piece .....	110
87. Examination of organs of neck .....	110-111
88. Method of opening trachea posteriorly .....	110-111
89. Examination of trachea and vocal cords .....	110-111
90. Finger method of tying intestine .....	165
91. Method of tying intestine preparatory to its removal .....	111
92. Bucket method of opening and cleansing intestines .....	111
93. Method of removing small intestines .....	166
94. Opening of small intestines after their removal from the body .....	166
95. Method of incising kidney with its ureter still attached .....	166
96. Method of opening kidney .....	177
97. Method of finishing opening of kidney .....	178
98. Method of removing capsule of kidney (without gloves) .....	166-167
99. Method of removing capsule of kidney (with gloves) .....	166-167
100. Relations of pancreas, kidney, ureter, adrenal, and solar plexus .....	182
101. Position for body in examination of rectovaginal region .....	166-167
102-109. Post-mortem extirpation of bladder, uterus, and adnexa through vagina, and subsequent restoration of parts .....	167
110. Method of opening uterus .....	183
111. Uterus and adnexa after being opened .....	184
112-114. Author's method of examining testicles, epididymis, spermatic cord, etc., without disfigurement .....	186
115. Method of opening seminal vesicles .....	186-187
116. Relation of gall-ducts and duodenum .....	186-187
117. Examination of bile-ducts .....	186-187
118. Method of examining stomach .....	187
119. Removal of liver from body .....	187
120. Method of incising liver .....	194
121. Starting-point of incision in removal of brain .....	226
122. Parting of hair so as not to injure it in removal of brain .....	226
123. Method of sawing skullcap .....	226-227
124. Angular method of removing brain .....	226-227
125. Method of breaking up inner table with an old knife after sawing ..	226-227
126. Method of drawing off skullcap with a retractor .....	226-227
127. French method of opening skull .....	228
128. French method of opening dura .....	229
129. Appearance of dura mater after removal of calvarium .....	227

FIG.	I
130. Appearance of brain after removal of dura .....	
131. Method of removing brain from skull .....	230-
132-135. Virchow's method of dissecting brain .....	230-
136-138. Virchow's method of dissecting brain (continued) .....	
139. Nauwerck's method of dissecting brain .....	236-
140. Basal ganglia, with cerebellum, pons Varolii, and medulla oblongata attached, in Meynert's method of dissecting brain .....	236-
141. Flechsig's, Brissaud's, and Dejerine's transverse sectioning of brain .....	
142. Dejerine's incisions for brain previous to hardening .....	
143. Lines for removing spinal cord and brain, the latter through a small triangular occipital incision .....	
144-149. Removal of spinal cord .....	244-
150. Method of examining nasopharynx, eyes, and ears .....	
151. Harke's method of examining nasopharynx .....	
152. Restoration of parts in Harke's method of examining nasopharynx .....	
153. Use of chain saw in examining auditory apparatus .....	
154. Examination of umbilical vessels .....	
155. Examination of ductus arteriosus .....	
156. Removal of spinal cord in child .....	276-
157, 158. Method of examining the femur for syphilitic osteochondritis .....	276-
159, 160. Method of examining nasal cavities, antrum of Highmore, etc. ....	
161. Slee's method of fixing skullcap .....	
162. Author's method of holding skullcap in place .....	
163, 164. Sewing up body .....	284-
165. Method of withdrawing blood from a body previous to injection of embalming fluid .....	284-
166. Injection of body with embalming fluid .....	284-
167. Refrigerating room .....	
168. Preparation of bodies with preservative fluids after removal from re- frigerating room .....	
169. Method of determining rectal temperature in a guinea-pig .....	
170. Animal holder .....	
171. Method of performing peritoneal injection in a rabbit .....	354-
172. Ear method of inoculating rabbit .....	354-
173. Post-mortem examination of guinea-pig .....	354-
174. Post-mortem examination of rabbit .....	
175-177. Post-mortem examination of horse .....	377, 378, and
178. Lines for opening cephalic cavities of horse .....	
179. Lines for opening cranial cavity of horse .....	
180. Lines for exposing cranial and nasal cavities in ruminants .....	
181. Appearance of cranial cavity in a cow .....	
182. Post-mortem examination of dog .....	
183. Exposure of oral and pharyngeal cavities in a dog .....	

# POST-MORTEM EXAMINATIONS



## CHAPTER I

### GENERAL CONSIDERATIONS

**HISTORICAL.**—The Code of Hammurabi, the Old Testament, and the ancient classics abound in references to violent death and the shedding of blood, but are silent up to the time of Herophilus (320–250 B.C.) as to any opening of the body for legal or pathologic purposes. It is known, however, that the Jewish priests examined the carcasses of animals killed for food to detect impurities, and the knowledge of anatomy displayed at an early date shows that human dissection must have been practised. In the Middle Ages postmortems were performed in cases of poisoning, Charles V. in 1530 empowering the judge to call in physicians as experts. In 1562 Paré made a judicial post-mortem, and thus established a medicolegal status which has continued until the present time. The office of coroner is an old and important one. It was created during the reign of King Athelstan, 925 A.D., and its duties were clearly defined soon after the Norman conquest. From England the institution was brought to America by the colonists, where the first post-mortem examination appears to have been made in 1639 in a case of fracture of the skull, and the second in 1643, death being caused by a bullet wound.<sup>1</sup> The authority of the coroner to hold an inquest is not confined to the body of a person who may have died within his jurisdiction, but extends to all cadavers brought within his territory, no matter where death may have taken place. (Becker.) Massachusetts in 1877 abolished the office of coroner, substituting therefor medical examiners, and the New York legislature in 1904 passed a bill for a similar purpose, which was, however, returned to the Governor with a veto by the Mayor of New York City.

---

<sup>1</sup> HOADLEY, *Proceedings Conn. Med. Soc.*, 1892, pp. 207–17, quoted by STEINER, *Johns Hopkins Bull.*, Aug., 1903. See also PACKARD, *Phila. Med. Jr.*, Feb. 17, 1902; EDITORIAL, *Jr. Amer. Med. Assoc.*, Oct. 28, 1893, p. 661.



**DEFINITION.**—Postmortem, autopsy, and necropsy are synonymous terms applied to the systematic exposure and critical examination of the cadaver with the object of determining the cause of death or of studying morbid anatomy in any of its various aspects. Other synonyms employed are necroscopy, mortopsy, section, *sectio cadaveris*, *sectio anatomica*, and *post* (colloquial). A medicolegal post-mortem differs from an ordinary postmortem only in the application of the information obtained to the furtherance of the ends of justice. The German word *Obduction* is correctly applied only to a medicolegal postmortem. The use of the word "autopsy," in the sense now generally accepted, was first made by v. Rühl, Crighton, and Bluhm, in an account by them of the examination of the body of the Empress Maria Feodorowna, of Russia.<sup>1</sup>

**PURPOSE.**—As the object of a post-mortem examination is the acquisition of exact data, the method of procedure should be scientific and systematic. This is especially important in medicolegal cases, which frequently involve not only the reputation and liberty, but even the life of a human being. If the examination be conducted in a perfunctory or desultory way, some detail of the greatest importance may be overlooked or the information obtained may be so ill arranged as to be practically valueless for statistical or demonstrative purposes. In no other department of medical science are the faculties of observation and discrimination more vigorously called into play, and in none other are sound knowledge and accurate work so indispensable.

Opportunities for the study of normal structures offered by post-mortems upon presumably healthy individuals killed by accident should not be neglected, as thorough familiarity with the appearance of the various organs and tissues in their normal condition is necessary in order that morbid changes or slight variations from health may be recognized. Such subjects also often afford favorable opportunities for the study of the earliest manifestations of disease, particularly in case of tumors and the infective granulomata. New anomalies also may be found, and these, as in the case of polydactylism, may be studied in order to support or disprove Mendel's and Galton's laws of inheritance. As the science of medicine advances, new discoveries

---

<sup>1</sup> *Salzb. med.-chir. Ztg.*, 1829, vol. i, p. 107; FOSTER'S *Encyclopædic Med. Dict.*, p. 516, quoting from KRAUS'S *Kritisch-etymologisches medicinisches Lexikon*.

necessitate a constant revision of the statistics of even the most common diseases.

Autopsies present exceptional opportunities for reviewing the study of anatomy and also for acquiring dexterity in the practice of surgery. To this end, it is permissible in suitable cases to perform surgical operations that entail no visible disfigurement of the body. Some of the more recently devised surgical procedures, such as the decapsulization of the kidney, the mechanical irritation of the hepatic peritoneum, the transplantation of ovarian tissue, the Lorenz operation for congenital dislocation of the hip, Gersung's injection of paraffin for the correction of deformities, the formation of an anterior and posterior cusp in the cervical os to prevent conception without interference with the outflow of the menstrual fluid, etc., will at once suggest themselves as being worthy of practice upon the cadaver as opportunity occurs.

**PERMISSION.**—When a postmortem is desired, the first step in every instance is to secure the legal right to make it. When not performed by order of a regularly appointed officer of the law, consent (preferably in writing) should be obtained from the next of kin to the deceased, or, in the absence of relatives, from the person in charge of the funeral. The feelings of friends and relatives must be fully respected: scientific zeal is no excuse for wounding them. In a suit for damages brought a few years ago against a Philadelphia hospital on account of a postmortem that had been made without the consent of the nearest relative, the judge severely deprecated the procedure, but held that no damages could be recovered in this instance, as the hospital was a charitable institution.

The method to be pursued in gaining permission will depend largely on the nature of the case, but the exercise of tact will nearly always overcome sentimental objections and secure consent except where religious scruples stand in the way. Thus, one resident in a hospital will obtain the opportunity of making an autopsy upon almost every patient dying in the wards during his term of service, while another interne of the same institution will, for one reason or another, meet refusal in the great majority of his cases. The curiosity of relatives and friends may be aroused, or the humane plea of doing no harm to the dead but possibly much good to the living will often appeal to the better judgment of those from whom consent is to be obtained. The author recalls a case in which those interested expressed

**DEFINITION.**—Postmortem, autopsy, and necropsy are synonymous terms applied to the systematic exposure and critical examination of the cadaver with the object of determining the cause of death or of studying morbid anatomy in any of its various aspects. Other synonyms employed are necroscopy, mortopsy, section, *sectio cadaveris*, *sectio anatomica*, and *post* (colloquial). A medicolegal post-mortem differs from an ordinary postmortem only in the application of the information obtained to the furtherance of the ends of justice. The German word *Obduction* is correctly applied only to a medicolegal postmortem. The use of the word "autopsy," in the sense now generally accepted, was first made by v. Rühl, Crichton, and Bluhm, in an account by them of the examination of the body of the Empress Maria Feodorowna, of Russia.<sup>1</sup>

**PURPOSE.**—As the object of a post-mortem examination is the acquisition of exact data, the method of procedure should be scientific and systematic. This is especially important in medicolegal cases, which frequently involve not only the reputation and liberty, but even the life of a human being. If the examination be conducted in a perfunctory or desultory way, some detail of the greatest importance may be overlooked or the information obtained may be so ill arranged as to be practically valueless for statistical or demonstrative purposes. In no other department of medical science are the faculties of observation and discrimination more vigorously called into play, and in none other are sound knowledge and accurate work so indispensable.

Opportunities for the study of normal structures offered by post-mortems upon presumably healthy individuals killed by accident should not be neglected, as thorough familiarity with the appearance of the various organs and tissues in their normal condition is necessary in order that morbid changes or slight variations from health may be recognized. Such subjects also often afford favorable opportunities for the study of the earliest manifestations of disease, particularly in case of tumors and the infective granulomata. New anomalies also may be found, and these, as in the case of polydactylism, may be studied in order to support or disprove Mendel's and Galton's laws of inheritance. As the science of medicine advances, new discoveries

---

<sup>1</sup> *Salzb. med.-chir. Ztg.*, 1829, vol. i, p. 107; FOSTER's *Encyclopædic Med. Dict.*, p. 516, quoting from KRAUS's *Kritisch-etymologisches medicinisches Lexikon*.

**DEFINITION.**—Postmortem, autopsy, and necropsy are synonymous terms applied to the systematic exposure and critical examination of the cadaver with the object of determining the cause of death or of studying morbid anatomy in any of its various aspects. Other synonyms employed are necroscopy, mortopsy, section, *sectio cadaveris*, *sectio anatomica*, and *post* (colloquial). A medicolegal post-mortem differs from an ordinary postmortem only in the application of the information obtained to the furtherance of the ends of justice. The German word *Obduction* is correctly applied only to a medicolegal postmortem. The use of the word "autopsy," in the sense now generally accepted, was first made by v. Rühl, Crichton, and Blum, in an account by them of the examination of the body of the Empress Maria Feodorowna, of Russia.<sup>1</sup>

**PURPOSE.**—As the object of a post-mortem examination is the acquisition of exact data, the method of procedure should be scientific and systematic. This is especially important in medicolegal cases, which frequently involve not only the reputation and liberty, but even the life of a human being. If the examination be conducted in a perfunctory or desultory way, some detail of the greatest importance may be overlooked or the information obtained may be so ill arranged as to be practically valueless for statistical or demonstrative purposes. In no other department of medical science are the faculties of observation and discrimination more vigorously called into play, and in none other are sound knowledge and accurate work so indispensable.

Opportunities for the study of normal structures offered by post-mortems upon presumably healthy individuals killed by accident should not be neglected, as thorough familiarity with the appearance of the various organs and tissues in their normal condition is necessary in order that morbid changes or slight variations from health may be recognized. Such subjects also often afford favorable opportunities for the study of the earliest manifestations of disease, particularly in case of tumors and the infective granulomata. New anomalies also may be found, and these, as in the case of polydactylism, may be studied in order to support or disprove Mendel's and Galton's laws of inheritance. As the science of medicine advances, new discoveries

---

<sup>1</sup> *Salzb. med.-chir. Ztg.*, 1829, vol. i, p. 107; FOSTER'S *Encyclopædic Med. Dict.*, p. 516, quoting from KRAUS'S *Kritisch-etymologisches medicinisches Lexikon*.



necessitate a constant revision of the statistics of even the most common diseases.

Autopsies present exceptional opportunities for reviewing the study of anatomy and also for acquiring dexterity in the practice of surgery. To this end, it is permissible in suitable cases to perform surgical operations that entail no visible disfigurement of the body. Some of the more recently devised surgical procedures, such as the decapsulization of the kidney, the mechanical irritation of the hepatic peritoneum, the transplantation of ovarian tissue, the Lorenz operation for congenital dislocation of the hip, Gersung's injection of paraffin for the correction of deformities, the formation of an anterior and posterior cusp in the cervical os to prevent conception without interference with the outflow of the menstrual fluid, etc., will at once suggest themselves as being worthy of practice upon the cadaver as opportunity occurs.

**PERMISSION.**—When a postmortem is desired, the first step in every instance is to secure the legal right to make it. When not performed by order of a regularly appointed officer of the law, consent (preferably in writing) should be obtained from the next of kin to the deceased, or, in the absence of relatives, from the person in charge of the funeral. The feelings of friends and relatives must be fully respected: scientific zeal is no excuse for wounding them. In a suit for damages brought a few years ago against a Philadelphia hospital on account of a postmortem that had been made without the consent of the nearest relative, the judge severely deprecated the procedure, but held that no damages could be recovered in this instance, as the hospital was a charitable institution.

The method to be pursued in gaining permission will depend largely on the nature of the case, but the exercise of tact will nearly always overcome sentimental objections and secure consent except where religious scruples stand in the way. Thus, one resident in a hospital will obtain the opportunity of making an autopsy upon almost every patient dying in the wards during his term of service, while another interne of the same institution will, for one reason or another, meet refusal in the great majority of his cases. The curiosity of relatives and friends may be aroused, or the humane plea of doing no harm to the dead but possibly much good to the living will often appeal to the better judgment of those from whom consent is to be obtained. The author recalls a case in which those interested expressed

great satisfaction on learning that death was not due, as had been diagnosed during life, to consumption. An invitation to a member of the family to be present at the postmortem or a promise to make a death-mask (see page 289) will often secure the desired permission. The laity should be encouraged to ask for an autopsy. A carefully performed postmortem often secures ready consent to, or even a voluntary request for, others in the vicinity in which the physician resides. The blank forms which accompany insurance papers often contain the query, "Was an autopsy made?" and an affirmative answer greatly strengthens the holder's claim. Indeed, insurance companies should encourage the making of autopsies, as it is to their own pecuniary advantage so to do. Our Boards of Health, with their enormous power for good or evil, in some States have the legal right to compel the performance of postmortems, a prerogative that has already been advocated more warmly by the lay press than by the profession at large. The offer of a small sum of money will often secure permission to make a necropsy among the indigent foreigners who are so numerous in our large cities, but a threat to refer the case to the coroner unless permission is voluntarily granted should never be employed. Undertakers who oppose the making of autopsies should not be recommended.

The pecuniary value that dead bodies may have sometimes gives rise to legal contests. The Supreme Court of California has decided that one cannot dispose of his own corpse by will. A man bequeathed his body to the managers of a medical college, in the hospital of which he had been treated, to be used for scientific purposes. The man's relatives claimed the cadaver, and applied to the courts for an injunction restraining the medical college from using it. The kinsfolk won, the court holding that the custody of the corpse and the right of burial belong to the next of kin.<sup>1</sup> There are in America and in France several societies the members of which sign cards granting permission for the performance of postmortems on their bodies: it would, however, on account of the decision of the court just referred to, seem best to have the card endorsed by the legal heirs.

Yet even when permission has been given circumstances may prevent the performance of the autopsy. Thus, in the case of Phillips Brooks, who was a member of the American Anthropometric Society,

---

<sup>1</sup> *American Medicine*, April 6, 1901.

the prosector of the society, on reaching Boston, could not perform the postmortem, as, a public funeral being universally desired, the body had been placed in an hermetically sealed coffin, death being due to diphtheria.

In the case of *Loesch vs. the Union Casualty and Surety Co.*,<sup>1</sup> the Supreme Court of Missouri held that the autopsy made without notice to the company was no bar to recovery. The physicians making the examination and the mother of the deceased, who tacitly assented to its performance, were in ignorance of the fact that there was a clause in the policy stating that if a postmortem was held without notice to the company in time to have its medical adviser present all claims under the policy should be forfeited. As soon as the error was discovered, which was in time for a re-examination, the company was notified.

There should be a law permitting post-mortem examinations of the bodies of all persons dying in charitable institutions. Such a rule exists in the hospitals in Germany, and this precedent for some time prevailed in the Philadelphia Hospital with practically no opposition, until a lawsuit, now pending, caused it to be abolished. In cadavers allotted to the anatomical board care should be taken not to destroy the arteries commonly used for injection. If in the course of an autopsy conditions are found which indicate foul play, as injuries or the presence of poison, the examination should be immediately suspended, and steps at once taken to have the coroner or other legal officer take charge of the case. If properly authorized by the coroner or the person who is legally acting in his stead, the examination may proceed in the manner prescribed for conducting medicolegal post-mortems.

When portions of the body are desired for preservation or for future study, permission to remove them should be obtained from some one connected with the household, though not necessarily from the nearest relative, as in gaining consent for the performance of the autopsy; it is, of course, unnecessary to tell how much is to be taken away! Should, however, the person authorizing the autopsy forbid the removal of any portion of the body from the house, no specimens should be taken. Consent can nearly always be obtained for the removal of small pieces of tissue for microscopic purposes, even

---

<sup>1</sup> *Jr. Amer. Med. Assoc.*, September 26, 1903.



DEFINITION.—Postmortem, autopsy, and necropsy are synonymous terms applied to the systematic exposure and critical examination of the cadaver with the object of determining the cause of death or of studying morbid anatomy in any of its various aspects. Other synonyms employed are necroscopy, mortopsy, section, *sectio cadaveris*, *sectio anatomica*, and *post* (colloquial). A medicolegal post-mortem differs from an ordinary postmortem only in the application of the information obtained to the furtherance of the ends of justice. The German word *Obduction* is correctly applied only to a medicolegal postmortem. The use of the word "autopsy," in the sense now generally accepted, was first made by v. Rühl, Crichton, and Bluhm, in an account by them of the examination of the body of the Empress Maria Feodorowna, of Russia.<sup>1</sup>

PURPOSE.—As the object of a post-mortem examination is the acquisition of exact data, the method of procedure should be scientific and systematic. This is especially important in medicolegal cases, which frequently involve not only the reputation and liberty, but even the life of a human being. If the examination be conducted in a perfunctory or desultory way, some detail of the greatest importance may be overlooked or the information obtained may be so ill arranged as to be practically valueless for statistical or demonstrative purposes. In no other department of medical science are the faculties of observation and discrimination more vigorously called into play, and in none other are sound knowledge and accurate work so indispensable.

Opportunities for the study of normal structures offered by post-mortems upon presumably healthy individuals killed by accident should not be neglected, as thorough familiarity with the appearance of the various organs and tissues in their normal condition is necessary in order that morbid changes or slight variations from health may be recognized. Such subjects also often afford favorable opportunities for the study of the earliest manifestations of disease, particularly in case of tumors and the infective granulomata. New anomalies also may be found, and these, as in the case of polydactylism, may be studied in order to support or disprove Mendel's and Galton's laws of inheritance. As the science of medicine advances, new discoveries

---

<sup>1</sup> *Salzb. med.-chir. Ztg.*, 1829, vol. i, p. 107; FOSTER'S *Encyclopædic Med. Dict.*, p. 516, quoting from KRAUS'S *Kritisch-etymologisches medicinisches Lexikon*.



necessitate a constant revision of the statistics of even the most common diseases.

Autopsies present exceptional opportunities for reviewing the study of anatomy and also for acquiring dexterity in the practice of surgery. To this end, it is permissible in suitable cases to perform surgical operations that entail no visible disfigurement of the body. Some of the more recently devised surgical procedures, such as the decapsulization of the kidney, the mechanical irritation of the hepatic peritoneum, the transplantation of ovarian tissue, the Lorenz operation for congenital dislocation of the hip, Gersung's injection of paraffin for the correction of deformities, the formation of an anterior and posterior cusp in the cervical os to prevent conception without interference with the outflow of the menstrual fluid, etc., will at once suggest themselves as being worthy of practice upon the cadaver as opportunity occurs.

**PERMISSION.**—When a postmortem is desired, the first step in every instance is to secure the legal right to make it. When not performed by order of a regularly appointed officer of the law, consent (preferably in writing) should be obtained from the next of kin to the deceased, or, in the absence of relatives, from the person in charge of the funeral. The feelings of friends and relatives must be fully respected: scientific zeal is no excuse for wounding them. In a suit for damages brought a few years ago against a Philadelphia hospital on account of a postmortem that had been made without the consent of the nearest relative, the judge severely deprecated the procedure, but held that no damages could be recovered in this instance, as the hospital was a charitable institution.

The method to be pursued in gaining permission will depend largely on the nature of the case, but the exercise of tact will nearly always overcome sentimental objections and secure consent except where religious scruples stand in the way. Thus, one resident in a hospital will obtain the opportunity of making an autopsy upon almost every patient dying in the wards during his term of service, while another interne of the same institution will, for one reason or another, meet refusal in the great majority of his cases. The curiosity of relatives and friends may be aroused, or the humane plea of doing no harm to the dead but possibly much good to the living will often appeal to the better judgment of those from whom consent is to be obtained. The author recalls a case in which those interested expressed



FIG. 2.—Body in a coffin prepared for a post-mortem examination. In this case the board supporting the body is elevated by means of a soap-box.



FIG. 3.—Post-mortem table for a babe, constructed of chairs and a drawing-board; the marble slab from a washstand or bureau may be used for the same purpose.





FIG. 3.—Post-mortem room of the Ayer Clinical Laboratory of the Pennsylvania Hospital.



FIG. 5.—Refrigerator box for the preservation of nine bodies, with scales so arranged as to weigh the corpse while it is in the refrigerator, if placed in the lower compartment next to the scales

tion. The fact that the appearance of the exposed parts is improved by the loss of blood and by its gravitation into the larger cavities of the body as a result of the post-mortem section may be mentioned to him as an argument in favor of the procedure. Fortunately, the formalin injecting fluid now generally employed for embalming purposes does not interfere with the microscopic study of tissues as did the arsenical preparation formerly used. Indeed, one of the special methods for hardening the brain is based on its previous injection with formalin by means of a cannula introduced through the orbit or nasal cavities.

The amount of preparation necessary for an autopsy will depend somewhat on whether the examination is to be made (I.) in a private house or at an undertaker's establishment, or (II.) in a hospital or morgue.

I. In the former case a table on which to lay the cadaver is rarely available, and a substitute must be provided. There are in the market several portable operating tables<sup>1</sup> which may be used for this purpose, as the one shown in Fig. 1. The postmortem may be performed while the body lies in the coffin, on the coffin-lid, or, still better, on the bottom of the inverted coffin, on the wooden slab usually found in the box, or on a door taken from its hinges and placed upon two kitchen chairs. The undertaker may have prepared the corpse for the autopsy, as seen in Fig. 2. For the body of a child the marble slab from the top of a bureau or wash-stand placed on the backs of chairs may be used. (Fig. 3.)

To facilitate the necessary manipulations, the cadaver must lie at a proper height. If placed too low, the stooping position required in making the autopsy is most fatiguing. A piece of oil-cloth, mackintosh, or old carpet should be placed under the table or its substitute, to prevent soiling the floor. In addition to the articles brought by the operator (see page 35), two buckets half filled with lukewarm water, an empty basin, and several newspapers should be provided.

Scrupulous cleanliness in the performance of an autopsy is of the greatest importance. The reasons for this are apparent. We owe it to our fellow-men to leave no malignant organisms in the place where the postmortem was performed. Besides, the pathologist can see bet-

---

<sup>1</sup> SHERMAN, *American Medicine*, October 26, 1901. Illustration from *International Clinics*, vol. I., Twelfth Series, 1902.



ter and his sense of touch is finer if the organs, fingers, and rubber gloves are not besmeared.

If the operator be careful not to soil his own person, the surrounding objects will be more likely to escape contamination. For this reason, he may accustom himself in private work to make necropsies on non-contagious cases with but little protection to his clothing. In France the usual dress consists of a hospital blouse, overalls of homespun, an apron reaching to the feet, and a pair of sabots or wooden shoes. In our hospitals the regulation duck trousers, shirt-sleeves, bare arms, rubber gloves, and an apron are most frequently seen. There should be a bountiful supply of water, a basin for the hands, and a board on which to arrange the instruments. The parts under examination should be cleansed as occasion requires by a stream of water squeezed from a sponge, the sponge itself not being permitted to touch the tissues. Mucous and serous surfaces should be carefully inspected before washing.

In private work the laity are likely to estimate the skill of the pathologist by the neatness displayed in sewing up the body and the appearance of the room after the autopsy is completed. The greatest care should be exercised that no blood-stains be left behind. Incense or cascarrilla may be burnt or ground coffee strewn on red-hot coals to remove the odor from the apartment, which should then be thoroughly aired.

II. In a hospital or morgue the facilities for making postmortems are much more complete. The room set apart for this purpose should be clean, well lighted, and secure against intrusion. The author remembers once having seen, much to his annoyance, a number of convalescents in the grounds of a hospital watching the performance of an autopsy through an open window. If practicable, the lead-house should communicate by an underground passage with all the wards of the hospital, and a covered or screened court for the undertaker's wagon should also be provided. A well-appointed mortuary room, like the one at the Ayer Clinical Laboratory of the Pennsylvania Hospital (Fig. 4), should have a refrigerator box with scales so arranged that bodies can be weighed within it. At the author's suggestion, the Fairbanks Scale Company fitted one of their scales to a Ridgeway refrigerator for this laboratory (Fig. 5), in such a way that the cadaver could be weighed while in the ice-chest and the result noted without opening the doors. For structural plans see Figs. 6, 7, and 8. The




FIG. 6.—Working plans for preparing refrigerator with eight compartments for the storage of bodies preparatory to their removal for burial. Front view.




FIG. 7.—Ground plan for a truck  
a, guide; b, track.




FIG. 8.—Cross-section.




FIG. 9.—Plans prepared for the writer by Mr. Addison Hutton for a post-mortem table at the Ayer Clinical Laboratory. (See also Fig. 10.)






FIG. 10.—Plan for a post-mortem table. (See also FIG. 9.)




FIG. 11.—Working plan for combination electric, gas, and water fixture above post-mortem table, prepared for the Ayer Clinical Laboratory.




FIG. 12.—Combination electric, gas, and water fixture to be placed above post-mortem table.

corpse should be weighed as soon as it is brought into the dead-house, as it will usually be found to lose weight after a time. Each box should have two doors, one opening into the post-mortem room, and the other into a waiting-room on the opposite side, through which the body may be viewed by friends and removed by the undertaker. This arrangement prevents the transmission of noises and odors. The waiting-room ought also to be such that religious services may be held in it, if desired.

The operating table should be strongly built, about seven feet long, two feet nine inches high, and three feet six inches wide. The top may be slate, soapstone, zinc, or copper; its surface should slope gently towards a central perforated depression connected with a drain and a ventilating shaft operated by an electric fan, and should be provided with sunken grooves converging towards the centre. (Figs. 9 and 10.) The drains of the post-mortem room should not connect with those of the hospital, but empty directly into the main sewer. An ideal though expensive plan would be to sterilize the waste water. A scale of feet, inches, and fractional parts of an inch (or of centimetres) should be laid off on the top of the table, or, if this be of slate, upon a metal rule sunk into it in such a way that no edges are exposed. For class instruction, a revolving table is required, upon which by an ingenious fulcrum and lever attachment the body can be weighed. An extra iron table like those used in the dissecting-room may be provided, in case it is desired to conduct two autopsies at once. The making of several necropsies simultaneously was discontinued in one Philadelphia hospital owing to the fact that three livers were found in a body subsequently exhumed on account of suspected poisoning. Additional tables, upon which to place instruments, scales, plates, and other requisites, should also be at hand.

Ample illumination should be provided, preferably by a northern skylight for day and by a combination gas-light and electric-light fixture directly over the table (Figs. 11 and 12) for night work. Plenty of water, hot and cold, should be supplied by means of an overhead spigot with rubber tubing attached, so that by the use of a mixer a steady stream of water at any required temperature may at once be had wherever desired.

To support the head there should be a solid block or a rest similar to those used by undertakers. This block should be about forty centimetres long, twenty centimetres high, ten centimetres broad, and

hollowed out on top to receive the nape of the neck. For children smaller sizes are to be employed. (See Figs. 66, 121, and 163.)

A board upon which organs may be placed after their removal, for convenience in making sections, etc., should also be at hand, as the slate slab becomes slippery from exuded fluid and the organs are held with difficulty while being incised. It is the custom abroad to set a stool upon which instruments are arranged within easy reach of the operator over the upper ends of the thighs. To avoid the spattering of dripping fluids when opening the cranium, it will be well to place a piece of previously moistened horse-blanket or a mop on the floor beneath the head. If the operator be subject to rheumatism, he should, while making the autopsy, stand on a piece of dry board rather than on the cement or tile floor usually found in mortuaries. The latticed wood flooring found on ships is well adapted to this purpose. The lavatories should preferably be the surgical kind operated by the feet. All linen, towels, etc., used in the dead-house ought to bear some distinguishing mark, and should be put at once into a proper disinfectant or sterilized apart from the other linen of the hospital.

A desk for the post-mortem book, a revolving chair, a slop-sink, a wash-stand, several cabinets, a work-table supplied with ordinary chemicals, a bacteriologic outfit, preservative fluids, and apparatus for preparing frozen sections complete the furniture of a well-equipped mortuary. The preparation of the latter adds greatly to the interest and value of an autopsy by enabling the operator to compare the microscopic and macroscopic appearances of a part while it is still in a fresh state. The use of ethyl chlorid as the freezing agent, where the more elaborate carbon dioxid or ether freezing apparatus is not at hand, may sometimes be advisable.<sup>1</sup> A library and a museum should be attached to the dead-house when possible.

---

<sup>1</sup> CATTELL, *International Medical Magazine*, December, 1896.



## CHAPTER II

### ORDER OF EXAMINATION AND POST-MORTEM RECORDS

PRECISION, with simplicity of technic, being the key-note for the proper performance of an autopsy, the following three rules will immediately suggest themselves as proper ones to be rigidly observed in the making of post-mortem examinations.

I. *Never disturb any part or organ until its position relative to adjacent tissues and organs has been accurately determined.*

II. *Never unnecessarily remove a part or organ if the proper inspection of remaining parts or organs will thereby be rendered difficult or impossible.*

III. *When an organ is to be opened in order to examine its cavities, walls, or component parts, the requisite incisions should be made in such a way as to permit, as far as possible, of the reconstruction of the organ in its original shape and condition.*

In the fulfilment of these conditions it is, therefore, best to begin by making a topographic examination of the contents of the cavity about to be explored. In the case of the trunk, the organs of the abdominal cavity are inspected first, those of the thorax next, and those of the pericardium last, whereas the removal of the organs and their minute description should be made in the reverse order. The abdomen should be examined before the thorax is opened, in order that the position of the diaphragm and the relative situations of the various abdominal organs can be determined before the entrance of air into the relaxed thoracic walls has altered the normal relationship, before the heart has been emptied of its blood by cutting the abdominal veins, and before the escape of blood and other liquids has obscured the appearances of the parts under consideration.

In order that nothing of importance shall be overlooked, the pathologist should have a definite plan of survey that he follows at every autopsy. The following order of examination is recommended:

1. Inspection of the exterior of the body.
2. Topographic exploration of the abdominal cavity.<sup>1</sup>
3. Topographic exploration of the thoracic cavities.<sup>1</sup>

---

<sup>1</sup> The organs are not yet incised nor are their relations markedly disturbed.

4. Pericardium.
5. Arch of the aorta.
6. Heart.<sup>1</sup>
7. Lungs. (a) Left. (b) Right.<sup>2</sup>
8. Larynx and trachea; external examination of the œsophagus.
9. Omentum, mesentery, and other portions of the peritoneum.
10. Spleen.
11. Intestines, except the duodenum.
12. (a) Left adrenal body and semilunar ganglion. (b) Left kidney. (c) Right adrenal body and semilunar ganglion. (d) Right kidney.
13. Ureters and bladder.
14. (a) In the male: Prostate gland, spermatic cord, urethra, testicles, etc. (b) In the female: Uterus, tubes, ovaries, broad ligaments, urethra, etc.
15. Duodenum and its ducts.
16. Stomach and œsophagus.
17. Liver and gall-bladder.
18. Pancreas and adjacent fat.
19. Retroperitoneal glands, the diaphragm, psoas muscle, thoracic duct, thoracic and abdominal aortæ, venæ cavæ, abdominal sympathetics, abdominal portion of the spermatic duct, etc.
20. Head. (a) Scalp and skull. (b) Meninges. (c) Encephalon. (d) Eye. (e) Ear. (f) Nasopharyngeal cavities. (g) Region of neck.
21. Spinal cord.
22. Bones, joints, peripheral nerves, arterial trunks of the extremities, muscles, tendons, etc.
23. Portions preserved, and the character of fluid employed.
24. Microscopic, chemic, bacteriologic, and physiologic examinations.

As a general rule, the order above suggested will be found convenient and practical. It must, of necessity, be subject to more or less variation, depending on the circumstances of the case. For example, in a medicolegal necropsy it is often advantageous to examine the seat of the suspected fatal lesion at once, and afterwards resume the order given above as nearly as possible. Thus, after death by poison the abdominal cavity is immediately inspected, while in a case of gunshot wound of the head the cephalic cavity is first investigated. The finding of anomalies, malformations, adhesions, etc., or the necessity of undertaking special lines of investigation may also cause a departure from the ordinary procedure. Thus, in autopsies on the remains of those who have died from nervous diseases it is often best to remove the brain and cord before opening the body.

<sup>1</sup> While the heart is being examined, time may be saved by having an assistant undertake the opening of the skull, as, theoretically, the heart should be exposed before the head is opened and the brain inspected before the heart is incised.

<sup>2</sup> The pleural cavities, already superficially examined, are to be most carefully inspected after the removal of each lung.

Letulle advises that the thoracic and abdominal organs be removed *en masse* from the body and first examined from their posterior aspect, as follows:<sup>1</sup>

(1) Large, or right, and small, or left lower, azygos veins. (2) Thoracic duct (dissection). (3) Suprarenal glands (dissection and removal). (4) Ureters (dissection). (5) Kidneys and their pelves (dissection and removal). (6) Thoracico-abdominal aorta (opened). (7) Inferior vena cava (opened). (8) Main portion of the portal vein and its branches of origin. (9) Common bile-duct and its two canals of origin. (10) Pancreas (dissection of posterior surface, tail, and borders). (11) Removal of the thoracico-abdominal aorta. (12) Dissection of the œsophagus to its point of entrance into the stomach. (13) Organs of mouth and pharynx: (a) incision of the pharynx; (b) dissection of the velum palati; (c) tonsils; (d) tongue; (e) sublingual glands. (14) Incision of the œsophagus at its point of origin. (15) Epiglottis and larynx (examination and opening). (16) Trachea and primitive bronchi. (17) Pulmonary roots (examination). (18) Lymphatic glands of the posterior region of the body (deep cervical, posterior mediastinal, diaphragmatic, prelumbar, retrorectal). (19) Cervicothoracic portion of the pneumogastric nerves.

After the posterior examination is completed, the parts are turned so that their anterior aspect comes into view. In doing this care is to be taken that the attachments are not twisted on their axes. The following order of examination from the anterior surface is then to be adopted:

(1) Thymus gland (examination and removal). (2) Thyroid gland (dissection and removal). (3) Opening of the superior vena cava and its branches of origin. (4) Study of the termination of the thoracic duct and the great lymphatic vein. (5) Pericardium (inspection and opening of). (6) Examination of the cardiac plexus. (7) Dissection of arch of the aorta and the thoracic aorta down to the seventh costal artery. (8) Pulmonary artery and its extrapulmonary branches (separation and opening of). (9) Pulmonary veins, extrapulmonary portion (separation and opening of). (10) Hilum of the lung (examination). (11) Examination of the exterior of the heart. (12) Removal of the heart. (13) Removal of the lungs. (14) Diaphragm (examination). (15) Liver and extrahepatic biliary ducts (examination and removal). (16) External examination and separation of spleen, stomach, pancreas, and duodenum. (17) Removal of œsophagus, stomach, pancreas, and duodenum. (18) Exterior examination, dissection, and removal of intestinal canal, with the exception of the rectum: (a) small intestine, (b) cæcum, (c) vermiform appendix, (d) colon, (e) rectum, (f) anus. (19) Examination of the peritoneum: (a) mesentery, (b) omentum, and (c) parietal peritoneum; (d) pelvic cavity. (20) Urinary apparatus (separation and examination of): (a) kidneys; (b) ureters; (c) bladder; (d) urethra. (21) Genital organs: (a) prostate, vesiculæ seminales, vasa deferentia, and testicles; (b) oviducts, broad ligaments, ovaries, vulva, vagina, and uterus.

---

<sup>1</sup> mode of procedure presents greater advantages in a child than in an adult.

A lesion found in one portion of the body may indicate the existence of pathologic conditions in another perhaps remote part. For example, multiple melanotic sarcomata of the liver are frequently secondary to a primary growth in the eye; embolism in the brain often arises from malignant endocarditis; hætomata of the ears will suggest chronic meningo-encephalitis, with thickening of the cranial meninges; and the presence of miliary tuberculosis should lead to an examination of the pulmonary arteries for tuberculous thrombi arising from caseous tuberculous glands. Again, particles of coal-dust embedded in the hands demand a careful inspection of the lungs for anthracosis, while bronzing of the skin will suggest scrutiny of the adrenals and of the sympathetic ganglia (Addison's disease).

If the ascertainment of the cause of death be the object in view, the line of inquiry should be based upon a hypothetical or tentative diagnosis suggested by the clinical history or special circumstances of the case. This may subsequently be corrected, modified, or abandoned as the autopsy proceeds; but the final diagnosis should, of course, not be made until the autopsy has been completed and any material requiring subsequent investigation reported upon by those undertaking this part of the work. There are cases in which it is impossible to state positively the cause of death, even on the completion of the autopsy, after a most thorough and painstaking examination. In such instances, as in all others, the accuracy of the conclusions drawn will depend upon the care exercised in the observation of details. Fortunately for those having to do with cases coming under the notice of the coroner, sudden death is nearly always attended by well-marked pathologic lesions. When no such cause of death is found, chemic or early microbic poisoning should be suspected. Any epidemic disease, such as smallpox, which is now (1904) so widely distributed throughout America, should always be thought of during the time of its prevalence, as death therefrom may occur before the characteristic rash or symptoms have made their appearance.

The following characteristics of each organ are to be noted, particular attention being given to those structures which are most vitally connected with the functional activity of the part.

1. Situation and relation to other parts.
2. Size and weight.
3. Shape, contour, borders, and coverings (capsule, serosa, mucosa, etc.).
4. Color.



5. Consistency.
6. Anomalies and malformations (congenital and acquired).
7. Fractures, dislocations, and lacerations.
8. Cut surfaces and liquid exuded.
9. Odor.
10. New growths.
11. Other pathologic conditions, taking into account the condition of the vessels to and from as well as in the part under consideration.

1. *Situation and Relation to other Parts.*—This takes into account any departure from the normal position or attachments of the organ. There are a number of regional landmarks frequently used; thus, in the case of the diaphragm we speak of its height in relation to the ribs or the intercostal spaces; of the stomach, as extending so many inches above or below the umbilicus; of the heart, in its relation to the nipples and the xiphoid cartilage; and of the cord, as to the vertebræ. While it may usually be easy to distinguish from which side an organ has been taken when there are no marked changes in its shape, the author has found that much time is saved and confusion avoided by marking each of the double organs as it is removed from the body, —one nick for the left and two nicks for the right-sided organs.

Plate I, which will be found upon the inside of the front cover, is based upon Cunningham's Anatomy, and will be helpful in the preparation of outline charts of the body for recording the situation and extent of lesions discovered at postmortems. The drawing shows the normal relations of all the important thoracic and abdominal viscera. Those who are unable to draw can place a sheet of thin paper over the figure and prepare an outline, which, after being filled in according to the exigencies of the case, may be pasted in the notes for future reference. One may purchase in England gummed outline charts in sheets of the brain and other parts of the body, upon which may be sketched in black ink the lesions which it is desired to record. Such drawings may then be sent at once to the engraver for reproduction in order to illustrate matter intended for publication.

2. *Size and Weight.*—For tables of weights and measures of the body, see Chapter XXIV. Whenever possible, it is advisable to give dimensions in centimetres and weight in grammes; if, however, measures and weights are to be used when giving testimony in a court of justice, it is well to convert them into inches and pounds, ounces, and grains avoirdupois before going on the witness stand. It should be remembered that a large organ is not necessarily a heavy one.

Atrophy and hypertrophy may be present in the same part, as seen in cases of hypertrophic cirrhosis of the liver in which acute yellow atrophy has supervened.

For determining bulk various means are employed besides actual measurement. A number of familiar objects at once suggest themselves, which may be used for comparison in describing the size of a lesion or part. Most persons have only vague and more or less erroneous notions concerning the dimensions of many common things, such as the head of a horse (usually underestimated) or the height from the ground of a stationary wash-stand (generally overestimated). A lesion may appear larger or smaller by being elevated or depressed and of a color like or unlike that of the surrounding parts. Virchow had, in the old Charité dead-house before its destruction by fire, a cabinet containing specimens of various familiar objects, such as beans, peas, lentils, barley, etc., with which pathologic lesions could be compared. (Plate II.) Later on, recognizing the relation of specific gravity to size and weight, he estimated the size by noting the quantity of water displaced by the organ when placed in a large flask of known capacity.


3. *Shape, Contour, Borders, and Coverings (Capsule, Serosa, Mucosa, etc.).*—All deviations from the normal in these particulars should be noted. It is often advisable to use the name of some well-known object in describing the configuration,—*e.g.*, cauliflower growth, hobnail liver, etc. As to the external appearance of a solid organ, its surface may be smooth, granular, nodular, shrivelled, or puckered. Here also we describe the capsules of organs, the serous coverings of the various parts, the mucous membrane of the stomach, vermiform appendix, etc. The borders of organs that have undergone infiltration are usually rounded and filled out; in degenerations they are generally flatter, thinner, and sharper than normal. Thus, in fatty infiltration the edges of the liver are rounded, while in cirrhosis its margins, often so largely composed of connective tissue as to contain practically no liver-cells, are sharply defined. The general contour of the blood-vessels may be markedly changed, as in aneurisms.

4. *Color.*—It is most difficult to describe colors or to reproduce them satisfactorily. Three-color printing does not give such good results as lithography. The best color values are obtained by the *kromskop*,<sup>1</sup> which has a wide field of usefulness not as yet recognized.

---

<sup>1</sup> CATTELL, *International Clinics*, vol. ii., Tenth Series, 1900.

Representation of certain familiar objects, with scales in millimetres and inches, useful for comparison in describing the size and form of lesions found at the postmortem.







Various shades of red are the most common tints found in the body; there is no such thing as pure white, even the conjunctivæ being a pearl-grayish pink. In pathology the word "pale" means relative deficiency of color. Note the color of the organ as soon as possible after exposure, as air, light, and water tend to alter it considerably, though naturally more or less change brought about by death has already occurred. Thus, the pericardium, which during life is transparent, is at autopsy only translucent. An organ should not be washed before its color is described, as water removes part of the coloring matter present, acts on the proteids, and modifies the original consistence of the organ; these changes may readily be demonstrated by placing the thymus gland with the surrounding areolar tissue in running water for five minutes. Air oxidizes the blood, so that a bluish stain may in a short time change to bright red. In the case of a congested lung it is well to note its appearance both before and after the blood has become oxidized. Certain abdominal organs are frequently discolored by a greenish slate tint supposed by some to be due to the deposition of the iron from the hæmoglobin by the hydrogen sulphid arising from decomposition. In a case of ammonium hydrate poisoning observed by the author, although the body was well preserved, the characteristic discoloration had penetrated the substance of the liver to a depth of three-quarters of an inch. Poisons often change the color of the blood markedly, and degenerations and infiltrations alter the appearance of the various parts affected. For a further description of the blood, the reader is referred to page 113.

5. *Consistency*.—This is learned only by experience, and is determined by pinching the organ between the thumb and the index-finger and by noticing its behavior when held in the hand. Palpation may often be practised with advantage. It should be remembered that consistency is affected by the season of the year, by the temperature of the cadaver and of its surroundings, by the length of the interval between dissolution and the making of the autopsy, by the manner of death, by the means used for the preservation of the body, and by various other influences.

6. *Anomalies and Malformations (Congenital and Acquired)*.—Each part or organ is subject to its own peculiar anomalies and malformations, and an entire chapter might readily be written upon the various altered conditions, congenital and acquired, revealed by autopsies. Thus, the writer has seen perforation of a typhoid ulcer in

a Meckel's diverticulum; free calcified bodies in the abdominal cavity; peculiar curvatures of the iliac arteries; the left kidney shaped like the spleen; the tip of the vermiform appendix resting near the pyloric end of the stomach; an artificial anus made by the rupture of a typhoid ulcer; the vermiform appendix in a *left* femoral hernia and the sigmoid flexure in a *right* inguinal hernia; a fish-bone in the omentum, etc.

7. *Fractures, Dislocations, and Lacerations.*—Every degree of injury may be represented. It should be remembered that from the external appearances alone it is not possible to state definitely the extent of the internal lesions. A heavy wagon may run over a child without rupturing the skin, though the internal organs may be lacerated and torn to a remarkable extent. The writer had an instructive case where a man struck a lamp-post with a push-cart, the handle of which entered his hepatic region; the external injury was not larger than a silver dollar, but hemorrhage from laceration of the liver finally caused death. Injuries are especially apt to be overlooked in those parts which are covered with hair.

8. *Cut Surfaces and Liquid exuded.*—When an organ is incised, describe first that which is most striking, as, for example, the presence of a hydatid cyst that is exposed on section of the spleen. Note the color of the exposed surface; whether it is smooth or granular; the amount, character, and chemic reaction of the fluid that is spontaneously exuded or is obtained by scraping with a knife; and the condition of the blood-vessels, especially as to atheroma and thrombosis. Numerous incisions may lead to the discovery of new lesions or afford an opportunity of studying the morbid process in its various stages.

Under the term "liquid exuded" are included not only blood, transudates, and exudates that follow incision of the part, but also any fluid that may be contained in the cavity of a hollow organ or in a cyst present, and the juice that appears on scraping or squeezing.

Œdema of an organ may be detected by squeezing it. In the lungs a frothy œdema shows the absence of a pneumonic infiltration. Surfaces should be scraped and the material thus obtained examined with the microscope.

In describing cavities pay especial attention to the lining membranes, noting their color, lustre, smoothness or roughness, and the presence of any adhesions; also the quantity, color, consistence, odor, and reaction of their contents and any sediments found therein.

9. *Odor*.—It is safe to predict that more and more attention will be given to the significance of odor. The organ of smell is imperfectly developed, and varies greatly in different individuals and in the same individual at different times. The peculiar odor that accompanies the growth of certain bacteria, such as the *Bacillus coli communis*, is well known. Smallpox, measles, cancerous ulcerations, and gangrene of the lung have their peculiar stenches. We may also mention the odor of acetone in diabetes, the pus-like odor in leucocythæmia, the butyric-acid-like or alcoholic odor from the brains of those who have drunk heavily before death, the uræmic odor, the odor in cases of carbolic or hydrocyanic acid poisoning, etc. The following poisons may also be recognized by their odor: opium, methylic alcohol, paraldehyde, chloroform, ether, formalin, creosote, iodine, iodoform, bromine, bromoform, chlorine, phosphorus, nitrobenzol, ethereal oils, amyl nitrite, ammonia, tellurium salts, etc. Fischer and Pentzold<sup>1</sup> state that the one five-millionth part of a gramme of chlorphenol or of mercaptan may be recognized by the sense of smell. For another illustration, see the action of the *Penicillium brevicaulis* on arsenical preparations, p. 331. Too often a case of apoplexy is taken to a police-station and the diagnosis is there recorded as one of alcoholism, simply because the odor of alcohol is found on the person arrested.

10. *New Growths*.—It is important to determine at once the presence of tumors, cysts, worms, etc., in a part, as subsequent manipulations may have to be markedly altered by their discovery.

11. *Pathologic Conditions*.—Every death may be attributed to one or more of the three following proximate causes: I. Interference with respiration, called asphyxia or apnoea; II. Interference with the heart's action, called syncope; and III. Interference with the nervous system, called coma or shock. The number of distinct diseases capable of producing death is limited, as will be seen by reference to the Bertillon classification of the causes of death given in Chapter XXVIII., the list not being nearly so large as one might expect without due consideration of this subject. There is also a distinct repetition of morbid processes in the different diseases and in the different parts. It is also well to remember that the histologic structure of an organ often at once gives information as to the lesions which will possibly be present in an affected part. In making a postmortem the diseases from which

---

<sup>1</sup> *Jr. Amer. Med. Assoc.*, April 23, 1904.



a person is liable to die and the lesions which may be found in any individual part or organ should always be carefully considered. Bearing this point in mind, considerable care has been taken in the preparation of the index in order to aid the reader in reaching a proper diagnosis by refreshing his memory as to the possible diseases or disturbances that may take place in the part or organ under study. By exclusion, the character of a lesion under observation may often at once be reduced to two or three possibilities, the final diagnosis being reached, in many cases, only by microscopic study. The index will also be found of use as suggestive in the preparation for an examination in pathology before a State Board or elsewhere.

NOTE TAKING.—Relying upon one's memory for records is a treacherous practice, and appearances which seem to be of no importance while the organ is before you are often of value to others who for various reasons may be called upon to read the protocol of the autopsy, but who have not had the opportunity of examining the parts in which they are interested. It is important, especially in medicolegal cases, to write "examined" or "normal"<sup>1</sup> after the number referring to the part under study, where no lesion exists, as this shows that an actual examination of this portion of the body has been made.

The notes should always be dictated in a distinct tone of voice and in easily understood language while the autopsy is in progress, and should consist exclusively of descriptions of the conditions then and there observed. Numbers and doubtful words should be at once repeated by the scribe, as an additional safeguard against error. Like the anæsthetist at an operation, the amanuensis should pay strict attention to the work assigned him. Names of diseases should be omitted in the notes themselves, but are to be inserted under the heading of "Pathologic Diagnosis" at the head of the report. The record of morbid changes present ought to be full, clear, and exact, so that from it alone the pathologic lesions can be inferred by another pa-

---

<sup>1</sup> Objection to the use of the word "normal" may properly be raised, for what one person may consider normal another would class as abnormal, while its use by an inexperienced person might lead to the omission of certain data which might be of importance in the future. It is, therefore, well to describe the part in detail. This will not only impress upon the obducent the normal appearances, but also lead him to make a more critical examination than he otherwise would be likely to do. The comparison of one organ with its fellow or of one part of the organ with another is often of value in this connection.

thologist as well as by the one who performed the necropsy. If the post-mortem record is rewritten, any descriptions given during the superficial examination may be combined with the detailed account of the parts removed from the cavity examined, thus permitting of the omission of any possible repetitions. One well-worded description of an autopsy dictated to a reliable amanuensis during the progress of the work is of much more value than scores written from memory after their completion. Drawings, photographs, skiagraphs, kromoskopie pictures, casts, microscopic slides, properly mounted museum specimens, and cultures of micro-organisms make valuable additions to a well-written account of a postmortem.

The liability to mistake, of which every day furnishes examples, is nowhere more forcibly exemplified than in the performance of post-mortems and the description of the appearances of the parts examined. What serious errors may result from poor writing or through misunderstanding, as in conversations over the telephone, is shown by the following illustrations. A pathologist communicated by telephone to the secretary of a surgeon the diagnosis of adenocarcinoma. The report received by the surgeon was to the effect that the patient "had no" carcinoma.<sup>1</sup> Often the word "atypical" is understood in the sense of "a typical." There is also an amusing side to this subject. In abstracting an article by Banti for the *International Medical Magazine* in 1895, the author wrote of the bacillus there described as being 4  $\mu$  long by 1  $\mu$  broad. The style of the office where the magazine was printed was to spell out numbers, and it appeared in the galley proof that "The bacilli are four feet long by one foot broad"!

Post-mortem records may be kept in a book specially prepared for that purpose, or on sheets to be filed away with the clinical history of the case under consideration. To every autopsy performed by myself I give a specific number, and lately have preserved my records on sheets kept in a flat-opening note cover-book, until they are ready to be filed away and indexed in properly-made manila covers. The interchangeable sheets in the note-book measure seven by eight and one-half inches. By means of an ingenious clasp opening in the centre, one end being fixed and the other movable, the leaves are held in place by passing the clasp through two small circular openings on the left-hand side of the page. When the clasps are closed, the leaves can be turned

---

<sup>1</sup> *Amer. Med.*, Dec. 5, 1903.

**PENNSYLVANIA HOSPITAL.**  
**POST-MORTEM RECORDS OF THE AYER CLINICAL LABORATORY.**

Autopsy number,										
Hospital number,										
Name,										
Occupation,	Ward,	Bed,	Sex,	Age,	Color,					
Died,	Nationality,	Weight,	Height,	feet,	inches.					
Weather,	Commenced autopsy	hours after death, and finished same at								
Permission given by	Clinical diagnosis of <sup>1</sup>									
	Performed by									
REMARKS.	PATHOLOGICAL DIAGNOSES.									

---

<sup>1</sup> This should always be entered before the beginning of the autopsy, and if there is any doubt as to the clinical diagnosis of the cause of death, such doubt should be stated, and the most likely cause written first.

like a book; when open, one or more sheets may readily be removed or others inserted. This method I find superior to the practice of keeping the records in special books or on the large index cards which are used by many physicians in preserving their private case records.<sup>1</sup>

In post-mortem books prepared for hospital records it is advantageous to have some data printed at the top of each page if the book be a large one or at the top of the left-hand page alone if the book be less than ten by fifteen inches, so as to afford ample room for notes. In my service at the Pennsylvania Hospital I used the form given on the opposite page.

The routine order of examination to be employed in the making of the autopsy, as given on p. 16, may then follow, or a card showing this order may be displayed in such a manner as readily to be seen by the one making the autopsy and the person to whom the notes are being dictated. Figures corresponding to the numbers of the divisions in the list may then be placed just before the notes describing the lesions to be sought for in the parts under examination.

Many writers advise the use of more or less elaborate printed descriptions of the various anatomic regions and organs, with blank spaces to be filled in at the time of making the autopsy. Printed books and forms for this purpose are to be found on the market, especially in England. This method of keeping notes has not in my hands yielded as satisfactory results as the one just described. I give, however, the following example of a post-mortem record, which was prepared in 1890 by Dr. Formad and myself and was in use for a number of years at the Philadelphia Hospital. The opposite (right-hand) page contained no printed matter, and could be used for more extensive descriptions or for the dictated record of the entire autopsy.

---

<sup>1</sup> *International Clinics*, vol. iv., Eleventh Series, 1902.

**BLOCKLEY ALMSHOUSE,  
PHILADELPHIA HOSPITAL—POST-MORTEM RECORD.**

No. of Autopsy,		Date of Autopsy,		Address,	
Name,		Age,	Sex,	Color,	
Ward,		Nationality,	189	Occupation,	
Admitted		Bed,	Died	hours after death.	Weather, { Cold, { Mild, { Hot.
History of { Alcoholic, { Syphilis, { Consumption, { Injury.		Clinical Diagnosis,	Autopsy,	Attending Physician,	
		Mode of Death, { Sudden, { Slow.	Autopsy ordered by	Resident Physician,	
				Performed by	

Cause of Death,	Weight of Body,	lbs.,	Height,
			{ Wounds, { Bruises.

I. <b>EXTERNAL APPEARANCES.</b> —Condition of Nutrition,	Scars,	Other Peculiarities,	P. M. Lividity,
Appearance of Skin,			
Deformities,			
Rigor Mortis,			
II. <b>HEAD.</b> —Scalp,			
Skull—size,	Thickness,	Injuries,	Eyes,
Fluid in Cranium, amount,	Thickness,	Injuries,	Configuration,
	Thickness,	Character,	
Dura Mater.—	Thickness,	Color,	Color, { Quantity, { Soft and thin, { Hard and thick.
Sinuses, quantity of blood,	Thrombi,		Adherent to Calvarium,
Pia Mater.—	Vascularity,		
Surface of Hemispheres,	Condition of Convulsions, { Flattened, { Asymmetrical.		Vessels at Base,
Brain.—Weight,	Color,	Puncta Vasculosa,	
Ventricles,—Size,	Contents,	4th Ventricle,	
Softening,	Blood Clots,	Tumors,	Other Lesions,
Base of Skull after removing Dura,	Areas of Degeneration,		
Deformities of Spinal Column,	Membranes,		Color,
III. <b>SPINE.</b> —	Thyroid,		Thymus,
IV. <b>THORAX and Neck.</b> —			
Tongue,	Pharynx,	Tonsils,	
Larynx,	Trachea,	{ Mediastinal Glands, { Bronchial	
Lungs.—			
Left Lung,	Pleural Cavities.—		
	Left, Adhesions,		
Right Lung,	Effusion,	{ Clear, { Purulent.	Blood,
	Right, Adhesions,		
	Effusion,		
Lesion,			

Heart.—Size, Pericardium—Adhesions, Heart opened in Situ shows Blood, { Fluid, Clotted. Ounces, Shape, { Globular, { Elongated, Hypertrophy of Cardiac Walls—None, Atrophy of Cardiac Walls, Valves and Orifices, { Chronic, Valvulitis, { Acute, Calcification, Aorta.—Arch, Thoracic, Abdominal, Other Blood-Vessels, Aneurism, Position of Diaphragm, Coronary Vessels, Tricuspid Valve,	Position, Ounces, Shape, { Globular, { Elongated, Left sided, Dilatation, Aortic, Mitral,	Effusions, Quantity, Red, Clots, { Chickens-fat, { White, tough, General, Fatty Metamorphosis, Mitral,	Character, Right Side, Right Sided, Tricuspid Valve,
<b>V. GENITO-URINARY TRACT.</b> —			
Kidneys.—Weight—Left, Ounces ; Lesion, Suprarenal Bodies, Bladder, Uterus, Contents,	Right, Ounces, { Color, { Shape, Ovaries, Peritoneum, Color, Cut Surface, Gall-Stones,	Ureters, Prostate, Testicles, Adhesions, Surface, Color of Bile,	{ Cyst, { Capsule, { Surface, Urethra,
<b>VI. ABDOMEN.</b> —Position of the Organs, Liver.—Weight, Ounces. Edges, Gall-Bladder and Ducts, Lesion, Pancreas.— Spleen.— Stomach.—Size, Congestion, Lesion, Small Intestines.— Vermiform Appendix, { Empty, { Full, Hernia, Large Intestines and Rectum.— Constrictions, Other Organs.—	Weight, { Empty, { Full, Mucous Membrane, Ulcers, Catarrhal Condition, { Empty, { Full, Hernia, Constrictions,	Thickness, Cancer, Ulcerations, Parasites, Intussusception, Dysentery, Hemorrhoids,	Peritoneal Lymph Glands, Oesophagus, Duodenal Ulcer, Contents, Obstruction, Volvulus, Contents,

**Specimens Preserved in Museum.—No.**



## CHAPTER III

### POST-MORTEM INSTRUMENTS AND HOW TO USE THEM

VARIOUS combinations of post-mortem instruments are found in the sets catalogued by dealers, but these, except for the systematic work possible only in hospitals and morgues, are more luxurious than necessary. The former wooden box with its plush lining is an abomination, owing to the impossibility of keeping it in a cleanly condition. The metal box is satisfactory, and one should be employed which can occasionally be sterilized by heat in its entirety. The ends of the box should be rounded so as to prevent any sharp edges from injuring the hands of the operator. If a box be used, all instruments should be thoroughly disinfected and returned to their proper places after each postmortem. It is annoying to take such a box to the place where the postmortem is to be held and then to discover the very instrument wanted to be missing. The instruments that are really indispensable for the proper performance of an autopsy are very few in number, as a complete examination may be performed in case of an emergency with a penknife and an ordinary wood-saw. Of course, in this field, as in surgery, ample opportunity has been offered for the exercise of mechanical ingenuity, and many instruments have been devised for facilitating post-mortem work that save much time and render greater neatness and exactitude possible.

The following list contains the instruments, apparatus, and chemicals most commonly used in the performance of an autopsy.

**KNIVES.**—*Section- or Cartilage-Knives.*—These should be made very strong, with a broad back, blunt rounded ends, and a bulge or belly at the outer third (Fig. 13), and should be narrower at the attachment of the blade to the handle. For general purposes the length of the entire knife should be from seven to seven and a half inches (about eighteen centimetres), the handle measuring about four inches. The Germans use knives even as long as eleven inches (twenty-eight centimetres). A separate rounded expansion for the index-finger found on the back of some section-knives is unnecessary (Fig. 14). The sharp-pointed knife should emphatically be condemned (Fig. 15). When the knives are sent to be sharpened, the



FIG. 13.—Section or cartilage-knife, with rounded end. (One-half natural size.)



FIG. 14.—Cartilage-knife with projection on back upon which the index-finger rests when making incisions. (Two-thirds natural size.)



FIG. 15.—Post-mortem knife with faulty point and without proper belly. (Two-thirds natural size.)




FIG. 16.—Coplin's brain-knife marked in centimetres on one side and in inches on the other. (Reduced.)




FIG. 17.—Bread-knife, useful in incising large organs, as the brain, the liver, etc. It comes in two forms,—with both sides meeting at the cutting edge like an ordinary knife, or with one side perpendicular and the other slanting for about three-eighths of an inch above the sharp edge, as shown near the handle in the illustration. (One-third natural size.)




FIG. 18.—Valentine's knife. (One-half natural size.)



FIG. 19.—Pick's myelotome. This little instrument is useful for severing the spinal cord in the removal of the brain. (One-half natural size.)



FIG. 20.—1, desirable form of forceps with spring; 2, saw with rounded end; 3 and 4, proper shapes of needles; 5, small saw with rounded end; 6, spring forceps; 7, box-jointed tenaculum forceps; 8, solid-headed hammer; 9, steel tape measure; 10, combined hone and strop; 11, scalpel with rounded hard-wood handle. (Reduced about one-half.)





FIG. 21.—A very desirable saw for post-mortem work; it is solidly constructed, and the teeth on the curved end are useful for sawing out the angles in the removal of the skullcap by the angular method. (Slightly less than one-half natural size.)




FIG. 22.—Butcher's saw, very useful for quick work in opening the calvarium. (One-quarter natural size.)




FIG. 23.—Hey's (Paré's) saw. (Two-thirds natural size.)



FIG. 24.—Metacarpal saw. (Slightly less than two-thirds natural size.)




FIG. 25.—Luer's double rhachiotome. This instrument is held in the right hand and steadied with the left by means of the handle attached to the fixed blade, the other blade being movable by clamps, so that the distance between the parallel blades may be varied at the will of the operator.




FIG. 27.—Strong scissors with short blades. (One-half natural size.)




FIG. 28.—Scissors with one rounded blade and with bent blade. (One-half natural size.)



FIG. 29.—Separate-bladed scissors, easily disinfected. The female blade is to the right; the male, to the left.



FIG. 30.—Proper form of enterotome. (One-half natural size.)




FIG. 31.—Improper left-handed form of enterotome, with pointed ends. (One-half natural size.)



instrument-maker should be cautioned not to grind them to a point. *Scalpels*, such as are used in dissecting. Those made of a single piece—i.e., without wooden, bone, or ivory handles—are to be preferred. The *brain-knife* (Fig. 96) should have a thin blade about ten inches (twenty-five centimetres) long, one and a half inches (four centimetres) broad, and blunt at the end like a table-knife. This instrument may also be used for incising the large organs and in opening the cavities of the heart. The brain-knife may be marked in the form of a rule and thus serve a double purpose (Fig. 16). An *amputation-knife* may be employed in place of a brain-knife, or in removing the brain through a trephine opening made in the skull. A *Waring bread-knife* (Fig. 17), which also does good work, may be used for incising the larger organs. A *Valentine knife* (Fig. 18), which has two parallel blades adjustable by screws to keep them the desired distance apart in order to cut at will thick or thin sections, is now rarely seen, but was much employed before the freezing microtome came into common use. Pick's *myelotome* (Fig. 19) is an instrument with a short blade bent nearly at right angles to the shaft, for cutting the spinal cord squarely across instead of in an oblique direction. A *curved probe-pointed bistoury* is used in cutting the dura mater, spinal cord, etc. A *razor* was formerly included in all lists of post-mortem instruments, but is now discarded.

**SAWS.**—The *saw* should possess a strong blade solidly attached to the handle (Fig. 21), as the two-piece jointed ones, kept in place by a screw, are very liable to become loosened. (Fig. 20, 2 and 5.) A butcher's meat-saw, which is arranged like a scroll-saw (Fig. 22) with its teeth pointing towards the front, its cutting surface measuring from ten to fourteen inches (twenty-five to thirty centimetres) for an adult and six inches (fifteen centimetres) for a babe, or a large cross-cut carpenter's saw, does the quickest work in removing the calvaria. *Hey's saw* (Fig. 23) is useful in sawing the angles when opening the skull. A *metacarpal saw* (Fig. 24) is often of service, especially in examining the femur of a babe for the detection of syphilitic osteochondritis. *Luer's double rhachiotome* (Fig. 25), employed for opening the spinal column, consists of two parallel saws with curved blades, the distance between which can be regulated by screws, and a very firm handle with a strong central support. Various forms of *dental and trephining engines*, usually driven by electricity, have recently been introduced and are useful in saving time

and labor. Among such engines may be mentioned those of Cryer <sup>1</sup> (Fig. 26), de Vilbiss, Wright, etc. These instruments are high-priced



FIG. 26.—Cryer's electrical surgical engine for cutting bone. A, spiral osteotome, with guard, for removing section of skull; B, spiral osteotome; C, trephine; D, guard for osteotome; E, electric motor; F, crank for hand propulsion; G, driving wheel for hand propulsion.

(from one hundred to three hundred dollars), on account of the infrequent demand for them. Hand-driven instruments may be purchased for twenty-five dollars and upward.

<sup>1</sup> *Medical News*, January 30, 1897.




FIG. 32.—Proper form of costotome; the handles do not meet by one-quarter of an inch and the ends are not pointed, but rounded. (One-half natural size.)



FIG. 33.—Improper form of costotome, with pointed blades and a catch, the handles meeting when the instrument is closed. (One-half natural size.)



FIG. 34.—Steel hammer with proper handle. (One-half natural size.)





FIG. 35.—Solid steel side chisel for breaking through any unsawed portions of bone in removing the calvarium. The pointed end is used as a pry and retractor for pulling out the sawed-off portion of the skull. (One-half natural size.)



FIG. 36.—Curved chisel, used for the same purposes as Fig. 35. (One-half natural size.)



FIG. 37.—Brunetti's left curved spinal chisel, of use in opening the vertebræ. (One-half natural size.)



FIG. 38.—Forceps.



FIG. 39.—Forceps.



FIG. 40.—Straight grooved director. (One-half natural size.)




FIG. 41.—Satterthwaite's calvarium clamp, closed and in use.






FIG. 42.—Iron clamp to be applied to the skull before the removal of the brain; especially used in dissecting-rooms.



FIG. 43.—Bigelow clamp for holding the head in the removal of the brain.




FIG. 44.—Folding iron head-rest.

**SCISSORS.**—One pair of scissors should be large and strong, with long handles and short, stout blades (Fig. 27); the other pair should have rounded ends with bent handles (Fig. 28). A pair with separable blades is frequently useful (Fig. 29). The *enterotome* is a scissors with one short and one long blade (Fig. 30), the latter being blunt and curved on itself at the end. Be sure that there is no sharp-pointed end, as this is the form usually supplied (Fig. 31). The *costotome* (Fig. 32) is an expensive instrument, with short, thick blades, the under one being curved and having a strong spring between the handles. Dangerous blood-blisters are sometimes produced by pinching the skin with the ends of the handles, which usually meet and fasten with a catch (Fig. 33). The ends should not meet and there is no necessity for the catch.

**HAMMERS.**—The most useful hammer is made of solid steel (Figs. 20, 8, and 34). One end of the head or striking portion is cuneiform, and there may be a hook on the end of the handle which is of service in springing off the calvarium. Lead filling in a hammer muffles the sound of its impact and prevents rebounding. A wooden mallet is preferred by some pathologists.

**CHISELS.**—There are chisels of various patterns devised for opening different regions. The *straight chisel* is the most serviceable, as it can be used in any region. The T-shaped chisel is also generally useful; it has one arm placed perpendicular to the other, and the arm which serves as a handle has one sharp and one blunt end so that it can be hammered upon. The chief use of the T-shaped chisel is in springing off the calvarium and in elevating the periosteum from it. Guarded, hatchet-shaped, and other chisels (Figs. 35 and 36) and spinal chisels (Fig. 37) are useful in opening the spinal canal, and a chisel with a guard about half an inch, or 1.25 centimetres, from the edge will not injure the brain while springing off the calvarium from the dura mater. The *raspatory of Chiara* has a broad, spoon-shaped end, four centimetres wide, with which the periosteum from a large surface can easily be removed; the other end is of the shape of a lance, one inch (2.5 centimetres) long, and is used for deep separation.

**FORCEPS.**—Dissecting forceps are indispensable when it is necessary to trace small structures; pointed, straight and curved forceps are the forms in use. (Fig. 20, 1 and 6.) *Bone-forceps*, large and strong and with rough handles, are necessary. One blade is blunt, so that it can be shoved against soft tissues without injuring them, as in

cutting the ribs. *Lion-forceps* of special type may be used when removing the bodies of the vertebræ. *Dura-tongs*, for pulling the dura mater away from the calvarium when it is adherent, may save the fingers from being injured by the bone. Other forms of strong forceps are seen in Figs. 38 and 39.

Grooved (Fig. 40) and curved DIRECTORS are frequently of use.

CHAIN HOOKS and a TENACULUM may be employed, but they are dangerous instruments. HOOKED RETRACTORS are more desirable than a tenaculum or chain hooks.

VARIOUS INSTRUMENTS.—A metal *catheter* and several flexible catheters, all of size number 8, may be needed for withdrawing urine. A *blow-pipe* with a stop or valve, a *trocar and cannula*, *probes*, some of which have eyes, and some form of *injecting syringe* are also useful. A *vise* is serviceable in firmly holding bone preparations in course of dissection, and in fixing a saw that is being sharpened. A *skull clamp* is considered by some to be of use in removing the calva-




FIG. 46.—Steel measuring stick marked in centimetres. (Reduced.)

rium (Figs. 41, 42, and 43). *Iron tripods* and other special devices for holding the head are shown in Figs. 44 and 45.

*Weights and measures* of various kinds are frequently found to be indispensable. These should include scales, a steel tape measure (Fig. 20, 9), graduated calipers, graduated glass cones, glass balls, and graduated measuring vessels of glass. The *scales* should have a capacity of twenty pounds, or ten kilogrammes, and be supplied with weights from a gramme upward. They are needed in weighing organs. The *steel tape measure* and the *two-foot rule* are marked both in centimetres and in inches. *Graduated calipers* or a measuring stick (Fig. 46) may be used in determining diameters. Fig. 47 represents an instrument




FIG. 45.—Cornell folding clamp for the secure holding of the head in the removal of the calvarium.  
(Specifications for the making will be sent upon application to the author.)





FIG. 48.—Metal pocket-case of instruments for finer dissection; very easily sterilized.  
(About one-half reduction.)








FIG. 51.—Various ways of wrapping linen twine, cut of proper length and ready for use.



FIG. 52.—Formad's leather pocket-case for holding the instruments usually employed in making a postmortem. (One-half natural size.)



FIG. 53.—Method of holding cartilage knife. The hand firmly grasps the knife, and the cutting is done with the belly of the blade.



known as a raspatory. A metal *pocket-case* for instruments is shown in Fig. 48. *Graduated wooden cones* (Fig. 49) and *glass balls* (Fig. 50) are serviceable for measuring orifices and canals. *Graduated measuring vessels* of glass are desirable. The larger vessels should be marked at every hundred cubic centimetres up to one or two litres, and the smaller for every two cubic centimetres up to a hundred. A *stomach-pump* is especially useful in withdrawing fluids from cavities. A *urinometer* is often of use. *Ladles* with a lip or spout, made of enamelled or agate ware, and with a capacity of half a pint, or two



FIG. 47.—Raspatory. (Reduced.)

hundred and fifty cubic centimetres, are needed in dipping fluid from cavities. A *whetstone* is useful, especially the form with a handle and a leather strop on the back of the stone. (Fig. 20, 10.) A *magnifying-glass* that enlarges at least ten diameters should be in the hands of every one making postmortems.

**OTHER SUPPLIES.**—*Enamelled trays* or *basins* are useful for receiving removed organs, and the basins are also required in cleansing the hands and instruments. *Blocks of wood* are needed to support the body. Metal supports are unwieldy and scratch the table. The blocks should be made of very hard wood, as of ebony, were it not for the cost, it being most difficult to secure boxwood of the proper size. Some of these should be prismatic in form, others excavated to fit under the neck during removal of the brain. All wooden utensils should be finished with oil so as to be non-absorbent. *Earthenware plates* or *wooden boards* are useful during the dissection of organs. *Needles* (Fig. 20, 3 and 4) and *coarse flax thread* or *fine twine* are needed in closing incisions made through the skin. The thread is also required in ligating the intestines before removing them. *Sponges* are a necessity readily procured, and should always be moist when in use. *Pins* are useful in fixing small structures in course of dissection. Special *tables* of zinc-covered wood, slate, iron, or glass are desirable in a pathologic department. Rotating tables are convenient, but weighing tables are expensive. The table should be constructed so as to carry off all fluids into a receptacle provided for them.

*Rubber gloves* that reach well up the wrist and *finger-cots* afford protection to the pathologist, particularly in cases where the danger of infection is great. The gloves are more readily put on and are preserved by dusting them freely with *ground soapstone* kept in a dusting bottle. *Quart museum jars* are useful for holding specimens to be preserved, and two-ounce, *wide-mouth bottles*, for microscopic specimens. A clean *glass bottle* with a glass stopper and *sealing-wax* to keep it closed are needed to receive the contents of the stomach in a case of poisoning. *Bromin* in a strong bottle with a ground-glass stopper that fits well serves a good purpose in disinfecting fresh wounds. Formaldehyde and distilled water should always be at hand.

*Pails* are needed as containers for water and to receive fluid removed from the body. *Cotton wool, sawdust, or tow* placed in the large cavities of the body before they are closed prevents the escape of fluid from them. An amusing story is told of a Philadelphia professor of pathology whose assistant used a self-raising buckwheat flour for the purpose; the formation of gas was so great that an explosion took place in the night. *Plaster of Paris and sand* serve a similar purpose in the cranial cavity. *Disinfectants and deodorants* should not be forgotten, as it is desirable to destroy or neutralize odors emanating from the body, and to disinfect and deodorize the hands of the pathologist after the examination has been completed. *Bellows* are occasionally useful in inflating viscera, as the lungs or stomach. A *hand-bag* which can be cleansed is often required for carrying instruments.

The *chemic, bacteriologic, and microscopic supplies* required in the work of the pathologist at the postmortem are red and blue litmus paper, turmeric paper, Lugol's solution, solution of sulphid of ammonium for detection of free iron derived from bile pigment, as in pernicious anæmia, Gabbett's solution, carbol-fuchsin, Loeffler's alkaline blue, absolute and commercial alcohol, ethyl chlorid or methyl chlorid, culture-tubes containing blood-serum, agar, and gelatin (bouillon is troublesome to carry), an alcohol-lamp, glass slides and covers for microscopic specimens, filter-papers three or four inches in diameter, an old scalpel which can be heated, a platinum wire three inches (or eight centimetres) long, set in a solid glass rod six inches (or fifteen centimetres) long, for making cultures (called an öse),<sup>1</sup> a microscope,

---

<sup>1</sup> Wetherill has recently suggested the use of an electric öse, the platinum wire being connected to a hand-battery of sufficient strength to cause the metal to glow when the current is turned on.



a freezing microtome, and easy access to an incubator. A Paquelin thermocautery or one of the simpler forms now so commonly used for wood burning is very useful for bacteriologic purposes.

In my own experience it has been found desirable or convenient to discard one instrument after another until now my satchel for private work weighs with its contents but ten pounds, and contains the following articles: two section-knives *in good condition*; a scalpel; a pair of medium-sized, strong scissors; a pair of bone-forceps; a dissecting forceps; a saw; an enterotome; a hammer with a hook on its handle; a pelvimeter; a new rubber catheter; gummed labels; various kinds of litmus paper; sealing-wax; a dissecting-apron and sleeves; a pair of rubber gloves, with plenty of ground soapstone in an iodoform dusting bottle; finger-stalls; a piece of thin rubber sheeting forty-five by thirty inches; a piece of oiled silk, or a rubber bag (sixteen by ten by four inches) from which fluid will not escape; two medium-sized bath-sponges; a quart museum jar graduated into ounces or cubic centimetres, into which some of the smaller articles are placed and which can be used for the removal of gross specimens later, if desired; a large needle and flax twine, cut and wrapped (Fig. 51) into three lengths, for sewing the body with single thread (forty-five inches), for sewing the head (twenty-five inches), and for tying the intestines (ten inches); some bromin in a strong bottle with a well-fitting ground-glass stopper; two per cent. iodoform celloidin solution; a small roll of cotton; four two-ounce, wide-mouth bottles for microscopic specimens, one of which should be filled with seventy per cent. alcohol, one with ten per cent. formalin, one with Müller's fluid, and the fourth with a saturated solution of mercuric bichlorid; two ounces of creolin; a cake of one per cent. bichlorid of mercury soap; an öse; an alcohol-lamp; several culture-tubes properly packed; incense powders; matches; pins, safety and ordinary; a steel tape-measure marked in inches and centimetres; a hand lens magnifying not less than ten diameters; spring scales weighing up to fifteen pounds; whetstone; and last, but not least, a note-book and several pencils, one of which will write on glass. To this list may be added other articles as the necessities of the case may demand.

For the private use of the general practitioner, a large section-knife, a scalpel, an enterotome, a saw, a chisel, a mallet, a pair of scissors, and a large needle may be purchased for about five dollars.

These should be kept rolled up in a piece of chamois-skin, preferably made with pockets into which the instruments fit, and if the latter be put away clean after use they are always ready for service; or a leather case (Fig. 52) may be employed.

The proper handling of post-mortem instruments is not acquired in a day, and the beginner will find that experience teaches many lessons which are not likely to be forgotten. A well-ground, keen-edged knife is a great desideratum, the advantage of a dull knife being simply that it is less likely to injure a beginner or careless operator and to disfigure the exposed portions of the body.<sup>1</sup>

In opening the body free incisions should be made by an easy, untrammelled movement, executed by the muscles of the shoulder rather than by those of the arm or hand. It is essential that the knife be grasped firmly (Fig. 53), and not held like a pen, as is a scalpel in the act of dissecting. Virchow says that the knife should be held in the whole hand, so that when the arm is stretched out the blade extends with it. The fingers and hands are fixed, if not absolutely, at least relatively, and the motion is executed with the whole arm, so that the movement is principally in the shoulder-joint and secondarily in the elbow-joint. Thus the whole strength of the arm and shoulder muscles is brought into play, and long, smooth incisions, so essential to proper inspection, are made. In cutting, pressure should be uniform, and the greater the pressure the quicker will the knife pass through the tissues. A clean cut made in the wrong place does less harm than a ragged one in the right place (Virchow). The portion of the blade near the handle should be used for work which dulls a knife, as cutting the ribs. This also applies to scissors, the part near the pivot being employed in all cases in which considerable force is required. When the knife is held as shown in Figs. 58 or 60, but preferably as in Fig. 60, the operator is sure to have a firm grasp of the knife-handle, so that there will be little likelihood of a dangerous slip. The actual cutting is properly and mainly done with the belly of the knife about one inch from its end, for which reason this part of the blade is always the thickest. The direction of the incision should invariably be from the operator, especial care being taken not to wound the left hand, and from those portions of the subject in which disfigurement would be most

<sup>1</sup> The method of holding and using the instruments will be seen illustrated by reference to the pages treating of the examination of the various organs.

likely to be noticed. Care must also be taken not to injure the assistants or those standing near. When the resistance of a tissue is unexpectedly overcome, the knife will sometimes travel a considerable distance before it can be stopped by an effort of the will.

The blade of the knife must be kept free from blood by frequent washing. This is especially necessary when incising organs, as the brain, in which incisions are made with much more satisfaction if the knife-blade be previously moistened. A pointed knife may be used for the removal of the tongue and the larynx, and a scalpel for fine dissection, as in tracing the spermatic or thoracic duct.

## CHAPTER IV

### THE CARE OF THE HANDS AND THE TREATMENT OF POST-MORTEM WOUNDS

Before beginning the autopsy, especially in a purulent case, the pathologist should carefully examine his hands; if these be not in good condition, the notes may be dictated by him while some one else is doing the actual cutting. All rings should be removed and the finger nails cut close, except possibly those of the thumb and index finger of one hand, which may be left slightly protruding in order to grasp certain parts, as the capsule of the kidney, to facilitate its removal, a feat often difficult of accomplishment without the use of an instrument, such as a knife, when rubber gloves are used.

Abscesses of the skin of the hands and forearms may be detected by the application of a ten per cent. solution of glacial acetic acid, which will at once reveal the location of such lesions by the smarting sensation that ensues. Slight wounds on the hands may be protected before beginning the autopsy by placing a small piece of absorbent cotton upon them and then applying the ordinary thick celloidin used in bacteriologic work, or the two per cent. iodoform celloidin already referred to. A varnish such as is used to coat pictures, liquid gutta-percha, or liquid resin plates may also be employed for this purpose.

It was once the custom for pathologists before starting work upon the body to cover their hands with some antiseptic solution, such as alcohol containing benz. coal. or germs to the extent of ten per cent. carbolic acid solution, or a solution of the balsam of Peru. If these be used they should be removed several times during the progress of the autopsy. It is however, much simpler to protect the skin surfaces grained by such applications by covering them with plaster. The fact that the body upon the face requires a slightly stronger antiseptic solution. This can be especially gotten by covering the left hand with the left hand, the right hand with the right hand, thus limiting the application of the antiseptic to these areas. The right hand is treated in the same place where an operation is being performed. The hands are protected by means of a bandage or the hands are covered with a rubber glove, which is discarded.



better. Of course, when digital investigations are necessary, as in exploring fistulous tracts, examining the vagina and os, and in certain kinds of peritonitis, antiseptic unguents are desirable; in such cases it is necessary to anoint only that hand or portion of the hand which comes in contact with the tissues under investigation. It is also an advantage sometimes to introduce vaseline into the crevices around the finger-nails.

An equally efficient and in many respects a much better safeguard against infection is the use of rubber gloves. Those should be selected which are neither too thick nor too thin, and not the old-fashioned thick, black, red, or white gloves. They should be provided with long sleeves, and should be purchased from a reliable dealer who has not had them too long in stock, as they markedly deteriorate by age. They fit snugly, and are especially useful when opening the stomach and intestines, as it is most frequently the intestinal contents which impart the odor that adheres so persistently to the hands. They do not prevent, though they to a certain extent hinder, the production of post-mortem wounds. After use they should be washed both inside and out with water to which a little washing soda has been added and scrubbed with a nail-brush until clean, rubbed lightly with a towel or absorbent lint until thoroughly dried, and then carefully dusted with powdered soapstone or with talcum powder. Either the weight of the water or pressure of the air may be used to force the everted fingers straight. With care rubber gloves may be sterilized in the autoclave. *They should never be put away moist or dirty.* Small openings may be patched in the same manner as a bicycle or automobile tire. Should the postmortem be upon a metallic poison case, a new pair of gloves should be used, and another pair from the same lot reserved unused for possible future examination by the chemist. Those of us who have made many hundreds of autopsies with our naked hands feel that we lose that delicacy of touch so desirable in post-mortem work when gloves are employed. Letulle,<sup>1</sup> in his recent work, enters into a tirade against their use. The coming generation will, however, undoubtedly wear them, or employ some such substitute as that recommended by Murphy,<sup>2</sup> of Chicago, who has recently suggested a method of dispensing with gloves in surgical opera-

---

<sup>1</sup> *La pratique des autopsies*, 1903.

<sup>2</sup> *Jr. Amer. Med. Assoc.*, Sept. 17, 1904.



tions. This consists in the application to the hands of a four to eight per cent. solution of gutta-percha in benzin or acetone, the former giving the better satisfaction in routine practice, as it is more lasting. The coating may have to be renewed during the operation, and when removed, by washing in benzin, leaves the hands soft and smooth. Rubber finger-stalls, especially the variety known as the seamless rolled finger-cot, which unrolls as it is placed on the finger, are useful if the operator have any hangnails or other abrasions of the fingers. They often break, however, during the performance of the autopsy. Blood, pus, or other fluid should not be allowed to dry upon the instruments used, upon the gloves, or upon the hands, for it not only impairs the delicacy of touch so desirable in this work, but may also cause unsightly stains upon the skin, which are difficult to remove, especially when certain preservatives have been employed in embalming the body.

The hands may usually be freed from odor by applying to them, while still wet, either a few drops of turpentine, formic aldehyde (from one to two per cent.), aromatic spirit of ammonia, listerine, paregoric, or mustard, and then washing them thoroughly with a good glycerin soap. Neelsen (quoted by Nauwerck) states that, if the odor can be removed in no other way, equal parts of fuming hydrochloric acid and glycerin may be used. The employment of equal parts of hypobromite solution (used in the quantitative estimation of urea) and of water, while severe, is also very effective for this purpose.

For disinfection of the hands after the postmortem one may use a creolin solution, made by placing about an ounce of creolin in a basin of tepid water;<sup>1</sup> a mixture of two teaspoonfuls of acetic acid, twice this amount of calx chlorinata, and a quart of water; bichlorid solution 1 to 1000; or a concentrated solution of potassium permanganate. The brownish discoloration of the hands may be removed by applying to them while still moist either oxalic acid or a concentrated aqueous solution of the bisulphite of sodium to which has been added a small amount of chlorin; or an antiseptic soap may be employed. Of the latter, I prefer the one per cent. bichlorid of mercury soap, or a ten per cent. lysol solution made with the tincture of green soap. Of course, any of the surgical methods in vogue for disinfection of

---

<sup>1</sup> Or, more exactly, a two per cent. creolin solution.

the hands may be employed. At the end a sodium bicarbonate wash and the application of a little lemon juice leave the hands in good condition. A brisk walk in the open air is also to be advised after the completion of the autopsy.

A post-mortem wound, as usually referred to, means not only a break in the continuity of the skin by an accidental incision, puncture, or other injury received at an autopsy, but also the inoculation therein of pathogenic bacteria from the cadaver, and their subsequent multiplication in the system, with the production of toxic symptoms. Wounds presenting similar appearances may, of course, be derived from many sources, as from surgical operations or from other post-mortem wounds. The intact skin of the hand is a perfect protective against the invasion of bacteria. In order that the organisms may infect the body, there must be both a point of entrance and a predisposition or lack of immunity in the individual affected. While any of the infectious diseases may be contracted in making a post-mortem, those most to be feared are tuberculous warts, syphilis,<sup>1</sup> gonorrhœal ophthalmia, tetanus, anthrax, glanders, plague, actinomycosis, typhus fever, yellow fever, cholera, and smallpox. I have seen septicæmia, general tuberculosis, ulcerative endocarditis, purulent meningitis, boils, whitlows, etc., follow post-mortem wounds. Several years ago one of my patients, who is now a justly celebrated veterinarian, suffered from a tuberculous wart which he had evidently contracted from a cow, thus adding another case to the list of those affected with bovine tuberculosis.

The results of a post-mortem wound depend very much upon the general health of the one affected, and experience seems to show that severer symptoms and slower convalescence may be expected in those who are habituated to the use of alcohol. Inoculations from serous surfaces are especially to be guarded against, as from some of the varieties of peritonitis due to criminal abortion, and other forms of septic peritonitis, meningitis, or pleurisy. Among other virulent kinds of post-mortem wounds may be mentioned those derived from cases of pyæmia, of septicæmia, of puerperal fever, of malignant œdema and diffuse cellulitis, of erysipelas, and of gangrene. Infection by the *Bacillus pyocyaneus* may cause long-continued high temperature

<sup>1</sup> J. DE LISLE (*Amer. Med.*, Sept. 19, 1903) writes, "Medical records furnish no instance of a specific contamination resulting from a wound received during the autopsy of a syphilitic cadaver."

with little local manifestation, as in my own case when I became inoculated with this organism from a case of cancer of the gall-bladder with secondary infection by this bacillus.

It is often asked why post-mortem wounds and injuries received in the performance of similar operations are more dangerous than those which are otherwise inflicted, though containing the same organism. Their greater virulence may in part be due to the fact that they are usually punctured wounds, in which the organisms are implanted deeply in the tissues, especially in cases of tetanus, which is due to an anaërobic bacillus. Again, it is well known that many organisms become more virulent by passing through successive animals, and, therefore, an organism which has overcome the resistance of the tissues and killed them is naturally more destructive than one which has not had such favorable opportunities for growth. It has been shown experimentally that bone-marrow possesses marked bactericidal properties. It is a well-established clinical fact that wounds produced by sharp spicules of bone are unusually severe. The reason given for this is that bacteria which have already overcome the increased resistance of the bone-marrow have now been introduced into the body.

Post-mortem wounds are generally caused in one of four ways: first, by the operator injuring himself with instruments used in the making of the autopsy, especially sharp-pointed knives and the saw; second, by scratches or punctures from ragged bones or calcified tissues, as the ribs or atheromatous patches of the aorta which have undergone calcareous infiltration; third, by inoculation of pre-existing wounds, abrasions, small eruptions, especially at the roots of the hair-follicles, hangnails, blisters, fissures in chapped hands, by infection from unsterilized instruments, by subsequent injuries received upon unsterilized hands, etc.; and, fourth, by cuts and scratches accidentally inflicted by the operator on his assistant, as in opening the head. Indeed, so frequently does the latter occur that a helper to steady the head should be dispensed with unless his hands be thoroughly protected by some covering, such as a towel. Some of the usual ways of producing wounds which are especially worthy of mention are by the operator cutting towards instead of away from himself or his assistant; by leaving a knife in one of the cavities and forgetting its presence; by placing his instruments in a dangerous position on the body, the table, or the ice-box; by the use of sharp-pointed



knives; by punctures from the needle made during the sewing up of the body; and by the too rapid passage of thread through the hands, producing a sort of brush-burn. Ragged wounds, such as those caused by the saw or by bones, are especially to be dreaded, for, being both punctured and lacerated, they are peculiarly prone to become infected.

The micro-organisms present at a postmortem made several days after death are apt to be less virulent than those encountered soon after dissolution, the saprophytes having now gained the mastery. Other things being equal, the more quickly the patient died after infection, the more dangerous will be the post-mortem wound; but the character of this lesion and the nature of the organism must always be considered. Undoubtedly, persons making many postmortems become immune to inoculation by the ordinary *Staphylococci* and *Streptococci*. When toxins are introduced along with the bacteria, the constitutional symptoms are apt to be more severe, as the toxins overcome a certain amount of vital force at the point of infection of the tissues which would otherwise aid in combating the micro-organisms.

As is well known, the bleeding of a wound is a considerable protection thereto; hence its immediate closure by the application of caustics or of celloidin is worse than useless. If the finger be wounded, it should be wrapped with a miniature Esmarch band and allowed to bleed freely under running water for at least five minutes, or the part may be washed in distilled water made alkaline by the addition of sodium bicarbonate, cleansed with equal parts of alcohol and ether, and then washed with an antiseptic solution. Sucking of the wound after cleansing has been practised. If a caustic be used, there is probably nothing better than glacial acetic acid, carbolic acid, or pure bromin. The employment of the actual cautery is advisable in some cases, but it must be so thoroughly applied that no pathogenic organisms are left behind, as otherwise the necrosed tissue affords a favorable medium for their growth. An antiseptic dressing of moist bichlorid of mercury gauze or dry sterile gauze is now applied, which should be renewed every twelve hours. On the slightest indication of pus or a deadish-gray appearance of the edges of the wound, it should be freely incised, thoroughly curetted, cleansed with a sterile salt solution, dusted with iodoform, and protected with a wet bichlorid dressing; or a solution of silver nitrate may be applied with benefit. I have seen no good effect from the local use of the



unguentum Credé (ointment of fifteen per cent. soluble metallic silver). The frequent application of hot flaxseed poultices containing a teaspoonful of Labarraque's solution is most grateful when the wound is discharging. Several inches above the wound a ring of iodine should be plentifully painted. Intravenous injections of antitoxic sera, collargol, formalin, and silver nitrate have been practised by some with alleged benefit in septic affections. Hume injects intravenously five hundred cubic centimetres of water containing one-half cubic centimetre of a ten per cent. solution of the nitrate of silver at a temperature of from 110° F. to 115° F.

Involvement of the lymphatics, as manifested by red lines running up the arm, usually on the inner surface, and tenderness in the axilla, indicates danger, and shows that the infection is no longer a local one. Inflammation of the lymphatics of the axilla may cause the glands in this region to become tender and enlarged, so that an incision is necessary; and in cases of axillary cellulitis, even though the wound of inoculation be small, early opening should be employed. Quinine is useful in these cases, and phosphoric acid and iron may be prescribed later. The affected arm should be carried in a sling, tonic treatment with changes of air instituted, and a surgeon consulted, who will treat the case according to the character of the wound, the nature of the infection, and the constitution of the patient. When healing has begun, massage has made many a serviceable finger or hand out of what would otherwise have been a stiff and useless one.

The anatomical wart is a local tuberculous lesion, often multiple, and is usually situated on the back of the hand or at the inner joints of the fingers. There is a warty thickening of the papilla of the skin, accompanied by a discharge of thin serous pus, but with no true ulceration. The sensation produced is similar to that caused by a splinter, which, however, subsides for several days after the removal of the fluid contents. The lesion sometimes heals spontaneously, but may give rise, as in one of my cases, to general tuberculosis. Wet dressings, combined with an application of equal parts of glycerin and extract of belladonna, may be employed, or the following mixture applied:

- R Salicylic acid, 10 parts;  
Extract of *cannabis indica*,  
Cocaine hydrochlorate, of each 1 part;  
Oil of turpentine, 5 parts;  
Glacial acetic acid, 2 parts;  
Collodion, 100 parts.

It would be interesting to try the hypodermic injection of tuberculin, or some of the newer forms of treatment of lupus of the face, as that of the concentrated rays of light recommended by Finsen.<sup>1</sup> In one of my cases treated by the X-rays in 1896 I thought that an anatomical wart was rendered worse by their use. When tuberculous warts have lasted several months, surgical treatment should be instituted, care being taken to remove them in their entirety without cutting into the diseased area. Guinea-pigs injected with such material usually linger a long time; in one of my cases over six months elapsed before the animal died from general tuberculosis.

Suppuration of the matrix of the nails can often be cured only by the removal of the nail, though frequent soaking of the finger in a hot saturated solution of boric acid or a strong solution of lead subacetate may be tried. Gwilym G. Davis recommends soaking the nail in a strong solution of silver nitrate—twenty grains or more to the ounce—and then wrapping the part in a moist bichlorid of mercury dressing. Dropping on a saturated solution of iodoform in ether has also been tried. Diffuse cellulitis should be treated by early and free incision and by the application of compresses. When the hand itself is involved in spreading gangrene, amputation should usually be practised. If tetanus is feared, the wound should be laid open and the area of contagion, if possible, removed, powdered antitoxin applied to the part, and the general health of the patient sustained. The subdural use of the antitoxin and, better still, the direct injection into the spinal canal have recently been employed.

If the knives used in post-mortem work were thoroughly sterilized after each necropsy, there would be fewer infected wounds. The making of autopsies is undoubtedly dangerous, and therefore those who perform them frequently should insure themselves in one or other of the accident companies which contain a clause giving a claim for benefits in case of wound-infection. As these companies generally show a marked disposition to dispute claims, every injury, no matter how slight, should be reported to them as soon as possible after its occurrence, so that any subsequent deformity or illness may thus be accounted for.

---

<sup>1</sup> BIE, *International Clinics*, vol. iii., Eleventh Series, 1902.

## CHAPTER V

### EXAMINATION OF THE EXTERIOR OF THE BODY

**SIGNS OF DEATH.**—The signs of death are of two kinds,—those which manifest themselves immediately upon the extinction of life and those which appear only after the lapse of a shorter or longer period of time, and vary at the time of dissolution according to the age, muscular state of the body, disease, presence of certain poisons, etc., and the external conditions surrounding the body. The later signs are the more positive, but the earlier ones are of more importance from a utilitarian point of view. Taken individually, each sign may be inconclusive, but when considered collectively they give a scientific authority to the generally only too apparent fact that death has taken place. Cases of trance and the Indian fakirs afford the best illustrations of suspended animation. Authentic instances of persons having been buried alive during suspended animation are almost unknown,<sup>1</sup> investigation of the newspaper accounts of such premature interments in almost every instance showing their falsity. In Munich the popular belief in such occurrences is so great that the bodies of those dying in the higher walks of life are kept for several days previous to burial in a specially prepared room, a bell being placed in the hands of the corpse for the purpose of summoning an attendant in case of resuscitation. Such notions usually originate from careless handling of the coffin, from the expulsion of a foetus by the formation of gases in the body of a pregnant woman, from real or apparent growth of hair, from conversion of bodies into adipocere, etc. In a judicial hanging the murderer is ordered to be hung by the neck until he is dead. The responsibility of fixing this time naturally evolves upon the physician.

The earlier or negative indications of death are first of all insensibility and inability to move, often preceded by the so-called death-rattle. There are loss of sensitiveness to stimuli, loss of reflexes, and the cessation of all tissue vitality, though in cases of sudden death spermatozoa will often be found in movement twenty-four hours after decease and atropine will dilate and physostigmine will contract the

---

<sup>1</sup> See ICARD, *Presse méd.*, No. 66, 1904.



pupils as long as molecular life exists in the ocular tissues. Loss of nervous and muscular irritability is determined by application of light to the eye, of snuff to the nose, or of cold, heat, force, electricity, or other irritants to the skin. Rosenthal considers the existence of electrical contractility in a dead body to be an indication that death has taken place within two to three hours. After the head is severed by the guillotine the eyes may open and close and in amputated limbs muscular twitchings may often be seen even after their complete removal from the body. Associated with the change in muscular tonicity is the facial expression. During the period of relaxation the visage is pale and flaccid, except in very rare cases, where the face even immediately after death has a red color and there is a drawn, contracted, painful expression, the so-called *facies hypocratica*.

Much more positive signs are the entire and continuous cessation of the respiration and circulation. As a rule, respiration stops a moment or so before the heart-throbs and ceases sooner in infants than in adults. The absence of breathing may be determined by auscultation and by the lack of motion of a down feather or small flame held before the lips or by the absence of the deposition of moisture on a cold mirror. A glass of water or of mercury placed upon the epigastrium will show a ripple on its surface if there be the slightest movement of respiration (Winslow's test). The X-rays have also been used to detect any activity of the heart and lungs.

Most of the minor signs of death depend upon the absence of circulation. This is determined first by observation. The skin of a dead person acquires almost immediately a leaden pallor or lividity and loses its transparency. The mucous membranes become pale and exsanguinated. The hands if viewed by strong transmitted light show no pink tinge where the fingers come in contact (diaphanous test). The palms and soles of the feet become more or less yellow in color. By palpation and auscultation the absence of pulse and heart-beat can be determined. A small artery, as the temporal, may be incised and examined; after death it will be found empty and its lining will be of a pale yellow color. The mercury in the manometer records zero. Scarification or cupping on a dead subject causes no flow of blood, while ammonia injected subcutaneously produces no congestion. In Icard's test fluorescein is injected subcutaneously; if life be present, local discoloration soon appears and the staining material may be chemically detected in blood abstracted at a distance from the point of



injection. A tight ligature around a finger, a limb, or the lobe of an ear will give rise to no reddening (Magnus's test). Pressure applied to a finger-nail will drive the blood away, leaving a white area, which will again be filled with blood if there is any circulation. If a flame or heat of any kind, as from melted sealing-wax, is applied to the skin, and a vesicle is formed, the blister on the dead skin will contain a non-albuminous fluid and the underlying cutis vera will remain dry and glazed, while in the living the contents will be rich in albumin and the cutis vera will be reddened. Caustics applied to dead skin will form no eschar, but may make the skin yellow and transparent. A steel needle plunged into a muscle after death has occurred will not be oxidized, even though it remain in place for many hours, but would tarnish in ten seconds if life were present (Glaister).

Besides these tests, there are changes in the eyes which are very important signs, and are in a great measure due to the loss of circulation. The fundus oculi is of a pale yellowish white and its vessels are empty or the column of blood in them is beaded by the presence of bubbles of gas. There is a marked loss of elasticity in the eyelids and in the globe. The eye collapses, sinks back in the socket, and appears flat and wrinkled; it loses its lustre and presents a glazed appearance. In some cases it is soft and flabby and may be covered with viscid mucus; more rarely, as in apoplexy or hydrocyanic-acid and carbonic-acid poisoning, it remains bright, full, and prominent for a considerable time. The conjunctiva quickly becomes cloudy and gray. The cornea may become opaque immediately after death or during the last hours of life; in other cases it does not change until the lapse of several hours. The cornea, iris, and conjunctiva lose their sensitiveness. In the last agony or shortly after death the pupils dilate, and again in about an hour they contract, as a rule unequally; this contraction lasts for three or four days. The loss of elasticity in the eye is very marked; the pupil can be made oval and will remain oval by synchronously compressing the globe (Ripault's test). This may, however, occur before death.

Another sign is complete loss of vital warmth. This occurs more rapidly at first than on nearing the temperature of the surrounding atmosphere. Clothing, fat, etc., cause the lowering to take place more slowly. Wilks and Taylor show that at an average temperature a nude dead body cools at the rate of about one degree Fahrenheit per hour. A body placed in water will cool more rapidly

than in air of a similar temperature, and refrigerants naturally reduce the temperature quickly. Nysten finds<sup>1</sup> that the bodies of those killed by lightning or by suffocation retain their heat longer than when death is due to other causes. Post-mortem cooling requires at least twenty-three hours for its completion. Brouardel says that the rectal temperature ordinarily is the same as that of the room in about forty hours. In the bodies of those who have died from some of the zymotic diseases, as cholera, tetanus, variola, etc., from injuries to the nervous system, or from certain abdominal disorders, the temperature may rise soon after death, and an elevation is also noted during the period of muscular rigidity. Where decease is due to some chronic affection chilling is slow, while after fatal hemorrhage it is very rapid.

The most positive sign of death is putrefaction, which appears after a longer or shorter time, and manifests itself first by a prominence of the superficial veins and by a greenish color in the iliac fossæ and the centre of the abdomen, later on the genitalia and thighs. Finally the whole body is involved. It becomes purplish red in color, due to the posthumous circulation, which is a displacement of the blood from the heart and large vessels by the pressure of the gases of putrefaction formed in the abdomen. Mummification and adipocere show that death took place some time ago.

POST-MORTEM OR CADAVERIC LIVIDITY; HYPOSTASIS OR HYPOSTATIC CONGESTION.—Unless drained of its blood by previous hemorrhage, a corpse usually shows a bluish-red to purplish-red discoloration on its most dependent parts, due to the cessation of the circulation and to the gravitation of the blood to those organs. The discoloration does not, however, appear upon those portions of the body upon which it directly rests. It will at once be seen that this fact may afford a basis upon which to form an opinion as to the position in which a body has lain after death. Post-mortem lividity rarely comes on before five hours; it reaches its maximum in the second day. It shows itself not only on the exterior of the body, but also on the dependent parts of such internal organs as the posterior wall of the stomach and the temporal lobes of the brain. Cadaveric lividity may resemble a bruise made during life, with all its various forms and shades of color. The distinguishing features are: (1) The discoloration in post-mortem

<sup>1</sup> *Arch. gén. de méd.*, June, 1862.

lividity disappears on pressure, while that due to a bruise does not. (2) A patch of post-mortem lividity will bleed freely when incised, because the vessels in the dependent parts are engorged with blood, while from a bruise there is little or no oozing, as the original hemorrhage is circumscribed and the discoloration is due to extravasated staining of the tissues and not to the actual presence of blood. An incision into the affected area should therefore be made in all suspected cases, especially in those of a medicolegal character. If the part be washed with running water, blood will appear again and again in hypostatic congestion. The two conditions are more closely simulated in those rare cases of hypostasis where we find a post-mortem oedematous infiltration and enlargement of the adjacent tissue. Should such exist or should the two conditions be combined, it is well to free the suspected area from the hypostatic congestion by turning the body on the opposite side for several hours before describing the bruise. As a rule, the more fluid the blood, as in cases of death from suffocation, the acute infectious fevers, poisoning by hydrocyanic acid, etc., the more marked will be the post-mortem lividity. In the latter case, as well as in poisoning by illuminating gas, the lividity may be of a characteristic rose-red color.

Bodies which have been kept for a long time (or a shorter time under unfavorable conditions) after death, especially during cold weather, present another form of cadaveric lividity which is characterized by a uniform reddish tint. This is caused by the diffusion of hæmoglobin from the blood-vessels into the surrounding tissues (imbibition). This form of lividity is most conspicuous along the course of the superficial veins and is not affected by pressure.

It is important to distinguish between post-mortem lividity and the greenish discoloration of commencing decomposition, usually first seen over the abdomen. According to some authorities, the greenish color is due to the precipitation of the iron of the hæmoglobin by the hydrogen sulphid arising from the decomposition of the tissues under the influence of bacteria, while others teach that it is due to chromogenic organisms themselves or a pigment elaborated by them of iron, potassium, and cyanogen. In one of my cases such discoloration was mistaken for the effects of personal violence, and serious allegations based upon this error were made against the husband of the deceased.

**POST-MORTEM RIGIDITY OR DEATH-STIFFENING.**—The involuntary muscles first show contraction, and it is doubtless this action



which sometimes causes the expulsion of a foetus after death. Post-mortem rigidity commences externally in the muscles of the lower jaw and spreads downward, disappearing in the same order, though Lacher, from an examination of six hundred bodies, found the condition to occur last in the arms. In ordinary cases it begins about two hours after death, is complete in from seven to eighteen hours, and ends as the stage of putrefaction comes on, in three or four days. Brown-Séquard quotes a case of typhoid fever where rigidity came on in less than four minutes, disappeared in a quarter of an hour, and putrefaction commenced in one hour after death. The stronger the individual and the shorter the duration of the fatal disease, the more prompt and marked will usually be the rigidity. The bodies of soldiers killed by being shot in battle after forced marches sometimes retain the position they occupied when they were hit, in certain cases even remaining erect when killed standing. The case of Captain Nolan at Balaklava is often cited in this connection, where it is alleged that he held a sword and with distended arm rode on horseback in the charge after death had ensued. The position of the hands produced by this muscular rigidity is also an important sign. The thumbs are usually flexed across the palms and the fingers flexed over the thumbs. Instantaneous rigor of the hand of a suicide may occur, a weapon being grasped tightly in the hand. This condition cannot be reproduced artificially, and shows high mental tension, nerve excitement, suicide, and not murder. Rigidity is marked, especially in the abdominal muscles, after death from cholera. The body of one who has died from tetanus, strychnine, or other spinal poison, as veratrum viride, may lie supported only by the head and heels, or when placed upright may stand erect with little or no support. In one of my cases of strychnine poisoning rigor mortis was present on disinterment twenty-four days after death. Suffocation causes long-continued post-mortem rigidity. Chronic alcoholism delays and prolongs it. The more muscular the individual, the slower is it in coming on and the longer in going off. Rigidity disappears more quickly in cachectic subjects, and is sometimes almost entirely absent after heat-stroke. The process does not depend upon the nervous system, but upon changes taking place in the sheaths of the individual muscular fibres. Section of one ischiatic nerve will delay, but not prevent, rigidity.<sup>1</sup> Laceration of muscles retards or may even preclude this

---

<sup>1</sup> BIERFREUND, *Arch. f. d. ges. Phys.*, 1888.



condition. The reaction of the muscles is at first acid, due to sarcolactic and other acids. As the rigor mortis passes off the parts become alkaline, a condition natural to a decomposing body. It should be remembered that in the preparation of the body by the undertaker the rigidity may have been overcome by force; this is especially true of the elbows. Be careful, on the other hand, not to be deceived by a previously existing ankylosis. Rigidity may be overcome by the use of hot applications, but when it has once disappeared it seldom returns and is never again so pronounced as at first. Rapid cooling delays rigidity, which, however, passes off the more quickly when the body is once more made warm. This condition must be differentiated from freezing, where on reduction there is a crackling sound. Cadaveric rigidity does not occur in the *immature* foetus.

DECOMPOSITION.—The bodies of infants decompose more quickly than those of adults. The process begins earlier in plethoric and fat adult bodies than in thin aged persons. It is more rapid after muscular activity and in those dead of acute diseases, fevers, heat-stroke, sepsis, suffocation by gases, etc., while it is longer delayed in cases where the system is exhausted and muscular irritability retarded, and in the bodies of those fatally poisoned by hydrocyanic acid, carbonic acid, sulphuric acid, etc. Arsenic may or may not prevent decomposition. At the same temperature a body which has been for one week in the air, one which has been two weeks in water, and one which has been eight weeks buried will show similar degrees of decomposition. (Brown-Séquard.)

Hofmann recommends in cases where decomposition is much advanced the removal of the brain in the ordinary manner, the making of some openings in the skin, the washing of the entire body in running water for twelve hours, and the further bathing of the corpse in a concentrated alcoholic sublimate solution or chlorid of zinc for an equal period. The green coloration due to decomposition disappears to a marked degree under this treatment.

The length of time which has elapsed since death has to be determined by the circumstances peculiar to each case. So many considerations may apply that in many instances it is dangerous to be too dogmatic.

EXTERNAL EXAMINATION OF THE BODY.—It is of great importance that we should not confine ourselves solely to the examination of the corpse, but should, in addition, carefully scrutinize the clinical

history, weigh the evidence derived from a personal survey of the surroundings, consider the circumstances under which death occurred, and question the persons who came in contact with the subject just before and after death. Data derived from such sources are of especial value in medicolegal cases or when a postmortem is to be performed upon an unidentified body; but the knowledge acquired by inspecting the surroundings and the exterior of the cadaver must in no way bias the operator before the internal examination is made, as the unexpected may happen here as well as elsewhere. The naked body is then to be minutely inspected, first as a whole for symmetry and then both anteriorly and posteriorly as to its component parts, proceeding in a definite and orderly manner. It would indeed be to our advantage in acquiring pathologic knowledge if the living body in the nude state were more frequently made the subject of careful study, for the information thus obtained is often of the greatest value to the clinician and surgeon.

**IDENTIFICATION OF THE BODY.**—Before a postmortem is begun, the remains should, if practicable, be positively identified to the obducent by one or more persons who knew the individual during life. If this be impossible, the one who found the cadaver or those who saw it in its original situation after death and those that removed it from one place to another may act as identifiers. Persons who have gone under several names should be recorded under their legally correct names, any aliases which had been used being also recorded. Certain details, such as articles of clothing, jewelry, and even pawn tickets, sex, age, height, weight, birth-marks, angioma, moles, tattoo markings, condition of the teeth, anomalies of the ear and eye, deformities, wounds, scars, or even the evidence of certain diseases, are of great importance and may often be the sole means of identifying the body. Should personal identification be impossible, a cast of the face, a photograph, and an accurate description of the body, with a full and clear statement of any peculiarities, should be made. In some cases a wax cast of the interior of the mouth, made afterwards in plaster, may be helpful. As the person whose body is being examined may have been a criminal and thus during life have had the Bertillon system applied for purposes of future identification, these measurements and finger-impressions should be secured in important cases. Skiagraphs of old osseous lesions, as well as a record of the teeth of the decedent and of their peculiarities, might also lead to identifica-

be estimated with any certainty, one may be able at least to designate the time of life as represented by the seven ages of Shakespeare.

**HEIGHT.**—The height is determined by measuring in a straight line from the vertex of the head to the centre of the external arch of the instep, the feet being flexed at a right angle to its plane of support. If a scale is not marked on the table and no other means of measuring is at hand, a piece of inelastic string or tape may be employed for this purpose and measured later. The writer suggests the use of a measuring apparatus modelled on the style of a shoe-measure. A simple form can readily be made by taking two one-foot rulers, or other sticks of about the same size, and attaching, one inch from one end, a seventy-eight inch tape-measure, which is made to run through a transverse slit one inch from the top of the other ruler. If a tape-measure of this length is not at hand, forty-two inches of inelastic tape may be sewed together and this attached to a measure of ordinary length. The first ruler is held close to the foot which is placed in a vertical position, and the other stick is held parallel to the first stick by an assistant standing at the head of the corpse, and the tape is drawn until it is taut. When not in use the tape-measure is wound around the sticks. Next measure the circumference of the head and shoulders. Should there be shortening of a limb, or atrophy, as in fracture and in infantile paralysis, full measurements of both limbs are to be made.

Certain abnormalities of stature are occasionally seen, such as: (1) Dwarfism, a condition which may be congenital or acquired. If acquired it may be either cretinoid or rhachitic, and is often associated with sterility, impotence, bone deformities, or atrophy of the thyroid. (2) Giantism appears in two types, infantile and acromegalic, which are intimately related. From the literature on this subject, it would seem that acromegaly frequently follows giant growth or even the period of excessive growth. This condition is often accompanied by an abnormal development of the genitalia and changes in the vascular glands, especially the pituitary and the thyroid.

If only part of a body is present, as in the case of Wakefield Gaines, where the trunk alone was found, the length may be approximated from various data. If the head and upper extremities remain, twice the length of the arm from the midsternal line to the tip of the middle finger, measured along the flexor surface with the arm in abduction, or the distance between the tip of the middle fingers along the flexor



surface, with the arms extended at right angles to the main axis, will about equal the height of the individual. If but one extremity is present, twice the length measured from the glenoid cavity plus one-half the distance between the glenoids measured between perpendicular lines, or nineteen times the length of the middle finger equals the approximate height. Other means of computing the height are: (1) The distance from the tip of the olecranon to the tip of the middle finger is five-nineteenths of the height. (2) The upper border of the symphysis pubis in an adult is the midpoint of the adult's height, but this is not trustworthy in women or persons with deformities. (3) From the head of the femur to the plantar surface of the heel is one-half the height, while the length of the femur is one-quarter the height. (4) Orfila has shown, however, that from one of the long bones alone the exact determination of the length of the body is impossible. Manouvrier in cases of extremely short or long bones multiplies by coefficients to secure the height, as follows: <sup>1</sup>

## MEN.

FEMUR.	TIBIA.	FIBULA.	HUMERUS.	RADIUS.	ULNA
Less than 392 mm.	less than 319	less than 318	less than 295	less than 213	less than 227
COEFFICIENT.					
x 3.92	x 4.80	x 4.82	x 5.25	x 7.11	x 6.66
More than 519 mm.	more than 420	more than 413	more than 368	more than 273	more than 293
COEFFICIENT.					
x 3.53	x 4.32	x 4.37	x 4.93	x 6.70	x 6.26

## WOMEN.

Less than 363 mm.	less than 284	less than 283	less than 263	less than 193	less than 203
COEFFICIENT.					
x 3.87	x 4.85	x 4.88	x 5.41	x 7.44	x 7.00
More than 478 mm.	more than 388	more than 376	more than 344	more than 250	more than 264
COEFFICIENT.					
x 3.68	x 4.42	x 4.52	x 4.98	x 7.00	x 6.49

<sup>1</sup> Rollet has prepared similar tables, which will be found in *Vibert's* work, 6th ed., pp. 561, 562.



(5) The length of the skeleton from the vertex to the calcaneum plus from four to six centimetres about equals the height of the individual, these figures being added to compensate for the loss of the interarticular cartilages, the intervertebral disks, and the coverings of the head and heel.

In many cases where homicide has been committed and the murderer has attempted to destroy the evidence of his guilt, or in destructive accidents, the *corpus delicti* has been proved by the finding of a part or member of the body or a portion of the clothing, as a piece of charred bone, a tooth, as in the Parker case, a ring, or a button. On the other hand, instances are on record where deluded individuals made confessions of murder which were proved to have been unfounded by the subsequent appearance in life of the persons said to have been killed. So important is this point that time and time again juries have failed to convict where the moral evidence was well-nigh conclusive. It is only upon irrefutable evidence that the fundamental principle concerning the *corpus delicti* is disregarded.

Where only a part of the body is available for examination, considerable difficulty is apt to arise as to the best method of procedure. The examiner will then need to possess a wide knowledge of comparative and pathologic anatomy and to exercise great ingenuity in order satisfactorily to demonstrate the identity of the parts submitted. Should the only proof of the *corpus delicti* be a skeleton or a portion of one, the expert may be asked to determine the age, race, and sex of the person, and the probable date at which death took place,—whether the bones are old or recent. With limitations, the age would be known by the condition of the epiphyses, whether united or not; by the cranial sutures, whether closed or not; and by the state of dentition. Race would be indicated by the different racial characteristics and peculiarities: thus, the negro by his splay-foot, projecting heel, and prognathous jaw; the Caucasian by his higher forehead, wider facial angle, and larger cranial capacity. Evidence of this character is not by any means conclusive. The determining of sex, after the age of puberty, presents less difficulty. In man the size of the cranium is greater and all the bony points are heavier and more prominent, the angle of the neck of the femur with the shaft is greater, and the lower jaw is heavier; in woman the bones are lighter and more compressed, the patella is smaller, and the articular surface of the femur and tibia is narrower. The characteristic differences are, however, found

in the broad female pelvis, the diameters of which are all greater with the exception of the vertical; the sacrum and coccyx are more curved and there is greater spread of the arches of the pubes.

The probable age of the bones would be indicated by their condition and appearance. The presence of the marrow and the periosteum is the most conclusive evidence of a recent state. The soft parts are usually destroyed within two years. Under ordinary conditions the body skeletonizes in about ten years, although this period is subject to wide variations, depending upon the cause of death, the chemical properties of the soil in which the body was found, and whether or not preservatives were used.

**WEIGHT AND NUTRITION.**—Next weigh the body, if this has not already been done, or at least estimate its weight. Particularly observe the state of somatic nutrition. If emaciation be present, note whether it is due to a deficiency of fat (*panniculus adiposus*), to muscular atrophy, or to a combination of both. This can readily be determined by picking up a fold of skin over a muscle and rolling it between the thumb and fingers. One may study upon his own person the differences existing in various parts of the body, noting especially the varying thickness of the integument in the front, the back, and the sides of the neck. The greatest emaciation occurs in phthisis, atrophic cirrhosis, muscular atrophy, and cancer of the upper digestive tract. The fat may be the corpulency of a high liver, of a gouty person, or of one suffering from cardiac affections. Women have a tendency to become obese as they reach the change of life.

An excessive deposit of fat may be due to *adiposis dolorosa*, in which condition there is a great increase of adipose tissue, not uniformly distributed, but occurring in lumps, the forearms, hands, legs, and feet often being without any or with but slight deposits of fat. In some cases there is also a thickening of the synovial membranes, with a tendency towards the formation of joint fungi and mice bodies,—probably a fatty infiltration. In one case reported by Dercum, in a man four feet eight inches in height, the patient weighed two hundred and six pounds.

**SKIN.**—Some of the bodies coming to autopsy are so filthy that no true estimate of the condition of the skin can be made until they have been cleansed. In vagabondism and alcoholism a distinct cutaneous discoloration often occurs due to the habits of the subject. Lice upon the body or head may be quickly and effectually disposed of by

saturating a towel with chloroform or kerosene and placing it upon the part affected for a few moments preparatory to its opening. Formalin may be used for this purpose if it can be applied several hours previous to the postmortem, so as to allow time for the evaporation of its fumes, which are so irritating as to forbid its application when the autopsy is to be made immediately afterwards.

The color of the skin is of great importance. It varies much in health and still more in disease and after the appearance of hypostatic congestion. Native Africans vary from yellowish brown to jet black. The children of negroes are usually creamy yellow when born, while it has been stated that the offspring of a mulatto mother and a full-blooded negro father is very dark at birth. The integument of a cadaver rarely possesses the rosy hue of health, but is rather of a grayish white, which shade is most conspicuous in cases of fatal poisoning by chlorate of potassium. The skin on those parts which have been exposed to the sun is generally more or less tanned, while in jaundice the color varies from the faintest tinge of yellow to a dark yellowish brown. Yellow color is also noted in chlorosis (yellowish green) and in pernicious anæmia (lemon-yellow). In the latter brown spots are also frequently found, usually situated on the abdomen, groin, buttocks, and thighs. Pallor is due to a primary anæmia or loss of blood, and is often so marked as to suggest the possibility of internal hemorrhage, as from the rupture of an aneurism or of the sac in extra-uterine pregnancy. The cachexiæ of cancer, argyria, etc., are at times peculiarly conspicuous in the dead body. The patches of bronzed skin, alternating with unaffected areas, seen in Addison's disease, may be scattered over the entire body, but are especially well marked on the abdomen; they are also sometimes found upon the mucous membrane of the mouth. This bronzing may occur when the suprarenal bodies are still apparently normal. Brown lines and a brown areola around the navel are observed during pregnancy; patches on the face may also appear. A white skin is found in albinism, vitiligo (where it occurs in patches), and in leprosy. Moles, tattoo marks, and certain cutaneous diseases, as leucoderma, cause characteristic discolorations of the integument. Redness of the skin is important. It may be simply a discoloration from some red clothing, or an erythematous inflammation, which, as a rule, however, disappears *post mortem*. In cases of asystolic cardiac disease a bluish-red cyanosis is often seen, but general reddening of the body is more likely to be a post-mortem



lividity, though it may be due to congenital or other forms of heart disease.

The breasts should be carefully examined in all cases, and the shape and size of the gland noted. The presence of any fluid in the breast should be detected by making pressure upon the gland and its character described after being placed under the microscope, especially in cases of abortion. Certain inflammatory conditions are found in the breast, the infection usually entering through a wound, abrasion, or fissure of the nipple. The infection may remain localized to the nipple, causing a simple ulcer or abscess, or produce a general affection. The abscess may be due to infection by the organisms of typhoid fever, tuberculosis, etc., the latter being miliary or diffuse. Several cases have been reported where the nipple became infected accidentally from vaccination. Atrophy of the gland is observed in the old, and hypertrophy is sometimes seen in the young. Supernumerary breasts and nipples occur; a well-formed breast with a nipple has been reported in the axilla and one in the groin. Tumors are very common in the female breast, but rare in the male. They are at times discovered by palpation, at other times are visible to the naked eye, and may, as in cancer, become great ulcerous patches. The most common tumor is cancer, which is most often found in the upper outer quadrant of the gland. Paget's disease is an inflammatory dermatitis of the areola and nipple, and is often a precancerous lesion. The glands of the neck, axilla, and supraclavicular region should be palpated and will often be found to be enlarged. Adenocarcinoma, adenocystoma, also known as chronic cystic mastitis, adenosarcoma, adenofibroma, adenocystic sarcoma, chondroma, myoma, myxoma, round or spindle-celled sarcoma, adenoma, lipoma, and fibroma have been found in these glands.

In the general survey of the body, relaxed abdominal walls, with the striæ of the multiparæ or of the patient who has had ascites, are to be described, also the enlargement of the superficial veins so often found in chronic heart, lung, and liver disease (especially in atrophic cirrhosis), tumors, aneurisms, thromboses of portal vein, etc. Gastroptosis and enteroptosis are often discernible on inspection. Œdema, general or local, is investigated especially as to its extent and the character of the pitting on pressure.

Now examine the skin for any abnormal marks, such as eruptions, scars, wounds, bruises, blood, dirt, discolorations, etc. The amount of cutaneous injury does not always afford a true index of the lesions



found internally. Thus, it is possible for a wagon or even a street-car to pass over a child without leaving any external trace other than a brush-burn, though upon opening the body the pelvis may be found crushed and the abdomen full of blood. As a rule, all eruptions and inflammations of the integument are pale and have a tendency to disappear at the postmortem. This is especially true of those on the mucous membranes and after the administration of certain drugs, as the purpuric rash from the use of quinine. The erythema produced by potassium iodid and mercury entirely disappears, though mercury may cause papules, vesicles, pustules, or even an exfoliative dermatitis, lesions which remain and can be studied.

Even an extreme eruption of measles may disappear *post mortem*, and in these cases, if a study of the lesion be desired, it is a wise precaution before death to mark the place to be remembered with a dermographic or anilin pencil or by the use of silver nitrate. In other diseases, as chicken-pox, smallpox, etc., the eruption is permanent. Chicken-pox, which may be coincident with smallpox, starts with vesicles that come out in crops. The eruption is very superficial, rarely umbilicated, and has no areola; it may become pustular, and often leaves scars. The lesions are most profuse on the trunk, especially the back, and as they dry up leave a black crust. Smallpox, on the contrary, begins as maculo-papules, which pass into vesicles, and lastly form pustules, all three conditions being often found at the same time in different parts of the body. The papules are deeply seated, indurated, and feel like shot when rolled between the fingers. The vesicles are multilocular and difficult to rupture. Smallpox may be associated with a pre-eruptive general purpuric rash. Malaria may be accompanied by urticaria, angioneurotic oedema, erythema multiforme, bullæ, herpes zoster, eczematoid eruption, and gangrene.

The number and variety of skin eruptions are legion, but eczema, acne, syphilis, alopecia, and psoriasis form 75 per cent. of all cases met with. In skin diseases certain definite lesions are found. Macules occur in syphilis, erythema multiforme, pityriasis rosea, pediculosis, measles, purpura, scurvy, rheumatism, peliosis rheumatica, extreme anæmia, typhus fever, and poisoning from snakes, mercury, antipyrin, etc. Brown macules include freckles, chloasma, moles, and nævus pigmentosa. White and pale yellow macules are seen in vitiligo, leprosy, morphœa, and facial hemiatrophy. Vesicles are found in herpes, especially around the eyes and the lips, in dermatitis venenata (ivy or

oak poisoning), impetigo, eczema, miliaria, and scabies. Blebs are seen mostly in impetigo, where they are flat and umbilicated, pemphigus, having no areola, dermatitis herpetiformis, and as clusters in syphilis. Pustules occur in eczema, acne, dermatitis herpetiformis, impetigo, varicella, ecthyma, smallpox, syphilis, scabies, and furunculosis. Papules occur in lichen, scrofulosis, prurigo, erythema multiforme, after the use of bromids, iodids, copaiba, cubebs, and tar, in eczema, miliaria, acne, scabies, syphilis, smallpox, measles, lichen ruber and planus. Ulcers are associated with syphilis, epithelioma, lupus, trauma, locomotor ataxia, bed-sores, etc. Large tubercles on the skin are associated with erythema nodosum, erythema multiforme, lupus vulgaris, syphilis, tinea sycosis, and leprosy. Crusting is found with eczema, seborrhœa, psoriasis, ichthyosis, syphilis, pityriasis, ring-worm, and after scarlet fever.

Besides these common forms there are other interesting lesions which should be looked for, as chimney-sweeper's dermatitis, ulcerating lesions of syphilis, actinomycosis and anthrax, scurvy and purpura, circumscribed keratosis, as in cutaneous horns, arsenical poisoning, keratoma, especially the senile seborrhœa of the French authors, which occurs in pinhead to dime-sized spots, more or less elevated, friable and slightly greasy or dry and hard, with yellow, brown, or black crusts which are firmly adherent to the skin and found especially on the exposed parts of the body.<sup>1</sup> These lesions, as well as nævi, may become epitheliomatous later on. In blastomycosis of the skin the lesions are mostly found on the hands, arms, face, and lower extremities. Tuberculosis of the skin is rare in this country. It appears in several forms, lupus vulgaris being the most common. The anatomical wart has been considered in Chapter IV.

Dermatomyositis sometimes occurs and may be described with the skin lesions. It consists of a swelling of the muscles, associated with an erythematous and pustular eruption and emaciation. The œdematous swelling may be followed by desquamation.

Certain occupations induce special affections, as the inflammations peculiar to those working in tar and paraffin, the necrosis of the jaw in match-makers, etc. Atrophy of the skin may follow injury to or inflammation of nerve-filaments. It at times accompanies pernicious anæmia. Induration of skin is seen in scleroderma, myxœdema, con-

---

<sup>1</sup> HARTZELL, *Jr. of Cutaneous Dis.*, Sept., 1903.

genital ichthyosis, and keloid. It accompanies œdema of subcutaneous tissues and scurvy, especially in the legs. Trophic affections of the skin, especially of the extremities, may be found in puerperal fever, gangrenous lymphangitis, diabetes, uræmia, ergotism, locomotor ataxia, etc., and is usually present in cases of angioneurotic œdema.

The yellowish deposits of xanthoma are among the striking and peculiar affections of the skin. They are found especially on the eyes and in the palmar creases. As one variety may occur associated with diabetes, this disease should be always borne in mind.

Skin eruptions are frequently found with Bright's disease. (1) Those of the early stage are pruritus, urticaria, and eczema. (2) Those of the final stage are universal erythema and bullous or desquamative eruptions. (3) Purpura or hemorrhagic eruptions may occur at any time during the disease, and affections due to marked œdema are also present in certain cases.

The tumors found in the skin are the wen or steatoma, lipoma, verruca or wart, nævus pigmentosus and nævus vasculosus, morphœa or keloid, molluscum fibrosum (which may cover the entire body), xanthoma, epithelioma (seen usually on the face), angiomas, which may undergo malignant change, adenoma, cancer, and sarcoma.

An entire chapter might easily be written on the significance and value of scars produced in various ways. Those made by the surgeon are often from their location self-explanatory, as the cicatrix after tracheotomy, trephining of the skull, or the mastoid operation.<sup>1</sup> It would, however, certainly facilitate matters, in this age of numerous hospitals and frequent operations, if the absence of organs removed by operation were indicated by some method which would be generally understood. Thus, the first letter of the Latin name of the part excised followed by the sign minus might be tattooed on the skin near the initial incision: *e.g.*, A— would show that the appendix had been removed, R— that nephrectomy had been performed, etc. The presence of scars may lead the obducent to think of herpes zoster, cupping, smallpox, chicken-pox, various skin diseases, as acne and syphilis, explosions, setons, certain occupations, previous application of croton oil, leeches, etc.

---

<sup>1</sup> The writer once desired to secure for a friend some fresh testicular tissue, and hurried to a recent suicide for the purpose of obtaining the testes. Finding scars on the scrotum, but no testicles, it was learned on investigation that these organs had been removed several years previously, and the young man, being in love, had hung himself because he felt that, being thus mutilated, he ought not to marry.



Scars made by the hypodermic needle in persons addicted to the use of morphine are usually found on the arms and thighs,—*i.e.*, in those situations which are hidden by the clothes and yet are easily accessible to the individual. Small abscesses containing pus are often present in these cases. Hypodermic injections by physicians shortly before death are usually made over the deltoid muscles or the breast, this region being selected owing to the quickness with which the drug is here absorbed into the general circulation. The puncture may be surrounded by an elevated white or reddish area similar to that produced by the application of cups. Recent saline injections (dermoclysis) also leave marks upon the skin. Exploratory punctures made by physicians at times end disastrously, as two of the several cases seen by me demonstrated. In one the trocar had penetrated the lung and given rise to abscesses which resulted in death. In the other case the exploratory needle used in searching for right-sided pleural fluid penetrated the liver and caused fatal hemorrhage.

Much discussion has arisen in regard to the means at our command for distinguishing a wound inflicted before and one made after death, and as to which is the fatal injury where there are more than one wound. On these and similar questions I have heard experts testify in court in a manner utterly unsupported by the facts of the case, and in a manner they would not do if talking before their county medical society. Great caution should, therefore, be used in the expression of dogmatic statements concerning such findings. Any blood found on the body should be accounted for and all bruises, injuries, etc., accurately located and described so as to be intelligible to the lay mind. In cases of death by electricity the points of entrance and exit of the current ought to be carefully sought for, and the shoes should be examined for the burns in the leather which are usually seen near the nails in the heels. In one of my eight cases of death from electricity a man carrying an umbrella with a steel handle, while looking into a shop, made connection with the arc light above the window and died instantly from the current thus transmitted.

Parchment-like spots are often seen on the body where the epidermis has been robbed of its protecting epithelium. Such areas are due to the drying of the part, and if produced during life there will usually be some ecchymotic spots around them. When seen about the mouth, they may have been caused by such agents as strong acids (especially carbolic) and alkalies.



The drawn-up and wrinkled appearance of the skin known as "goose-flesh," or cutis anserina, excoriation of fingers, mud, sand, water plants, etc., under the nails and in the hands, mouth, nostrils, and ears are especially conspicuous after drowning.

Note the presence of bed-sores and blisters, remembering that scalds found on dead bodies are sometimes due to carelessness in the use of hot-water bags or bottles during the final illness.

The region of the neck should be carefully inspected for finger markings, scratches, rope markings, etc. The neck should be rotated so as to ascertain if a fracture or a dislocation exists. In infants a sucking gland is found in each cheek. Cysts may be due to occlusion of the salivary glands.

Enlarged glands can be seen or palpated; they become conspicuous in cases of tuberculosis, leukæmia, cancer, etc. An enlarged thyroid should be measured and examined. A slight enlargement of this gland is often seen in pregnant women and women at term, the hypertrophy disappearing during the puerperium. Percussion may reveal a pleurisy, extensive infiltration of a new growth in the mediastinum, an ileocæcal abscess, ascites, enlarged organs, as a spleen or liver, etc. The presence of gall-stones or of an hydatid cyst may at times be elicited by manipulation.

Ascites can sometimes be detected by an elevation of the umbilicus, the finger being used to depress the part. Echinococcus cysts, encapsulated peritonitic exudates, ovarian cysts, primary carcinoma, and flaccid cysts of the urachus may all be found at the navel.

**DEFORMITIES.**—The body should now be carefully examined from head to foot and from left to right, and any variations from the normal, either bony or muscular, minutely described. Deformities may be congenital or acquired, single or multiple, symmetrical or asymmetrical. Babes<sup>1</sup> believes that there is a special centre, situated in the anterior base of the skull in the upper part of the face, which presides over the development of the limbs, and that disease of this region produces a tendency to the formation of symmetrical deformities.

*Congenital Deformities.*—These may be due to embryologic defects or to traumatic or pathologic intra-uterine causes. Injury during delivery is responsible for quite a number of these cases, Allis recently advocating that congenital dislocations of the hip are produced at the

---

<sup>1</sup> *Berliner klin. Wchnschr.*, 1904, vol. xli, no. 18.

time of birth. Those interested in the pathology of congenital dislocation of the hip will find an excellent illustrated article on this subject by Carl Ludloff, in *Klinisches Jahrbuch*, 1902, vol. x, no. 1.

According to Hirst and Piersol, the most common monsters are:

I. SINGLE MONSTERS: (1) Autositic Monsters. — *Ectromelus*, aborted or imperfectly formed limbs; *symelus*, a union of two limbs and imperfectly developed pelvis; *celosomia*, body cleft with some eventration, and with anomalies of limbs and genito-urinary apparatus; *exencephalus*, a foetus with malformed brain, part at least without cranium; *pseudencephalus*, bones of vault absent or very rudimentary and brain rudimentary; mouth a mere opening; *anencephalus*, a foetus without a cranium or brain; *cyclocephalus*, a union of the eyes, generally with an absence of the nose; *otocephalus*, lower jaw wanting; ears approach each other. (2) Omphalositic Monsters. — *Paracephalus*, imperfect extremities; imperfect head and face; lungs absent or rudimentary; heart often absent; one member of a unioval twin, sex feminine; *acephalus*, complete absence of head and upper extremities, rudimentary or absent heart, lungs, etc.; *asomata*, trunkless head, which is not well formed; no cord; *anideus*, shapeless mass covered with skin.

II. DOUBLE MONSTERS: Double Autositic Monsters. — (1) Terata Katadidyma: *Metapagus*, two foetuses united by their cephalic extremities; *pygopagus*, two foetuses united in the region of the buttock; *ischiopagus*, two foetuses united by the pelves, coccyges, and sacra, with a common umbilicus; *dicephalus*, with two distinct heads, usually separate necks; *diprosopus*, having a double face, body single. (2) Terata Anadidyma: *Dipygus*, double pelvis, lower extremities, and genitalia; *syncephalus*, division up to navel and imperfectly formed up to head; *craniopagus*, bodies joined at homologous parts of the cranium. Laloo was an illustration of a *Dipygus parasiticus*. (3) Terata Anakatadidyma: *Prosopothoracopagus*, twins united by the thorax, abdomen, and face; *omphalopagus*, united from umbilicus to xiphoid cartilage; *rhachipagus*, united at the vertebral column. The Siamese Twins were examples of monsters of the *xiphopagus* variety.

III. TRIPLE MONSTERS. In composite monsters there is a complete or partial union of two or more foetuses.

IV. DOUBLE PARASITES: *heterotyphus*, a parasitic foetus hanging from the anterior abdominal wall of the principal; *heteralius*, a parasite inserted at a distance from the umbilicus of its host and having no

direct connection with the latter's cord; *polygnathus*, ill-developed fetal parts joined to jaw of autosite; *polymelus*, duplication of lower extremities; *endocyma*, the greater part of the parasite within the body of the autosite.

*Spina bifida* is a defect in the union of the laminæ of one or more vertebræ, with more or less malformation of the spinal cord or its membranes. While spina bifida usually shows itself posteriorly, it may do so entirely alone or in combination with an anterior opening, and, at times, with increase of size due to the presence of a lipomatous mass. *Encephalocele* is a hernia of the brain, and *meningocele* a hernial protrusion of the meninges.

Various other defects and lesions may be found: as, *e.g.*, complete or partial absence of the *nose*; imperforate ala nasi; deviation of the septum, various abnormal shapes, as saddle-back in hereditary syphilis. *Palate*, cleft. *Cheeks*, fissures and fistulæ. *Ears*, absence of the helix; hæmatoma. *Mouth*, imperforate, abnormally large (macrostoma), abnormally small (microstoma). Atresia oris, besides being congenital, may be due to cicatrization from burns. *Tongue*, absent; cleft; atrophied (microglossus); hypertrophied (macroglossus); or it may be adherent to the palate. The frænum may be too short or too long. *Alveolar process*, absent; cleft; atrophied; hypertrophied. *Lips*, cleft (harelip); cysts. *Neck*, tracheal fistula; cysts on visceral clefts. *Fingers and toes*, absent; atrophied; hypertrophied; supernumerary; webbed; or clubbed. *Sternum*, absent, malformed, or fissured. *Ribs*, cervical ribs and various defects in their development. *Umbilicus*, skin insertion of the cord; abnormalities of the vessels; hernia. *Urachus*, persists and remains patent. *Bladder*, extroversion. *Penis*, the glans may be atrophied, hypertrophied, or phimosis or atresia may be present; often imperfectly formed in cretins. The entire penis may be absent, but this condition may be due to amputation, traumatic or pathologic, in which case the scar will be present. Scars on glans or prepuce are usually syphilitic. An elongated or fissured penis is often associated with calculi. Epispadias or hypospadias may exist. The penis may be found erected after death by hanging, injury to spinal cord, or drowning. It may be affected with gangrene or cancer, and the arteries may show arteriosclerosis, the latter condition being rare. *Testicles*, one or both may be absent from the scrotum or from the body altogether; they may not have descended, presenting the condition of undescended testicle, or may be



found in some abnormal position. There may be atrophy; hypoplasia, a condition often present in imbeciles; hypertrophy, congenital or acquired, or compensatory in one testis, as after the removal or destruction of its fellow; duplication or malformation. Abdominal, crural, cruroscrotal, or ilio-abdominal ectopia may occur. These organs are usually retracted in cases of drowning. *Scrotum*, absent, atrophied, or hypertrophied; gangrenous lymphangitis is seen sometimes in children. The hypertrophied condition may be acquired, as in elephantiasis. Other conditions may be cleft scrotum, hydrocele, hæmatocele, varicocele, and hernia. It may be contracted in cases of drowning. *Vulva*, absent; imperforate; atrophied; hypertrophied. The Bartholinian glands may become enlarged, forming retention cysts; abscesses; tumors, as fibroma, chondroma, lipoma, sarcoma, carcinoma, and myoma. Aphthæ occur as white spots on the mucous membrane; elephantiasis; herpes progenitalis; diphtheritic ulcers; acne; eczema, especially on the skin of the labia, the vulva, or the nymphæ, sometimes giving rise to atresia or stenosis; lupus; syphilis, as the chancre, mucous patch, or gumma; gonorrhœa, as a purulent vulvitis; chancroid; and venereal warts are found. Injuries are common after parturition and rape. Hæmatoma, rupture of varicose veins, lacerations, œdema, etc., are seen. *Clitoris*, absent; atrophied; hypertrophied, in which case it may simulate hermaphroditism; carcinoma. *Anus*, absent or imperforate, or may end in a blind sac. Fissures, indurated and irregular. Small polypoid growths may fringe the borders. All lesions around the anus are at times altered by the distortion of the part with the cotton introduced by those who have had the body in charge. *Rectum*, prolapsed. This condition is common in children. The rectum may contain congenital polypi; internal or external hemorrhoids. *Fistulæ*, internal or external, complete or incomplete, follow abscesses. *Hymen*, absent; imperforate; fimbriated. Its absence may be due to traumatic causes or rupture during menstruation. An *ovary* may lie in the canal of Nuck. *Urethra*, absent or occluded; atresia or partial phimosis. It may have abnormal openings, as on the penis, scrotum, perineum, clitoris, or rectum; in the last case forming a urethrorectal fistula. It may be cleft, presenting a condition of epispadias or hypospadias. It may be inflamed (urethritis), with or without *Gonococci*, and showing a bloody, mucopurulent, purulent, or altered spermatic discharge. Chronic urethritis occurs with thickening of the tube. It may be torn by the passage of a



stone or foreign body. Stricture may occur, in the male, as a rule, four to six inches from the meatus. Tuberculosis is extremely rare. Tumors, as fibroma, angioma, sarcoma, epithelioma, are seen. Condyloma or caruncle may be found.

Congenital hypertrophies may be confined to the big toe, and are, as a rule, associated with disturbances of the genitalia or a persisting thymus gland. In rare instances the enlargement is general, as in giantism, a condition not uncommonly acquired, when it is apt to be irregular and partial, affecting usually the bones of the face and skull (leontiasis ossea). Although appearing soon after birth, it more often arises at puberty and is due to an abnormal proliferation of the cartilages in the process of endochondral ossification. Acromegaly, a condition due to some lesion of the pituitary body, is often a cause of giant growth, the enlargement affecting the face bones and the distal ends of the long bones. Local hypertrophies due to inflammation and rhachitis are not at all uncommon. Rokitansky describes deposits of phosphates and salts of lime in the cranial and pelvic bones of pregnant women; these are the analogues of the "plaques choriales" of sheep, which probably contribute to the development of the fetal skeleton.

The opposite condition, atrophy, is much more common. The general congenital form is the microsomic dwarf (normal proportion), a rare condition, the stunting more often affecting only one part, and being due to ischaemia or inflammation *in utero*. It may affect a limb (agenesis), skull (microcephalus), pelvis, etc. Acquired microsomia (cretinism, etc.) is the result of absence or disease of the thyroid gland, which produces an arrest of development in the longitudinal growth of the cartilaginous bones and in the lateral growth of the membranous bones. Rhachitis and synostoses are other causes of stunted development. Partial atrophy, if congenital, is often confined to the head, tibia, fibula, or radius, and is, as a rule, associated with other deformity. There may be an entire absence of bones or parts, as in apodia, or the lack of a clavicle, scapula, or radius.

*Signs of Degeneration.* — Closely connected with malformations are the signs of degeneration, as misshaped ears and nose, asymmetrical face, deformed fingers, and some anomalies of the penis, vulva, and anus, which should be noted for their statistical value.

*Fractures.* — Deformities due to fractures are very common. Their character depends upon the location, the bones broken, and the age of the fracture. A recent fracture will exhibit crepitus, swelling,

increased mobility, and deformity due to contracture of the muscles; this contracture, however, may disappear *post mortem*. If the fracture is old there may be non-union, false union, union with deformity, false joint or a non-absorbed callus. Deformity due to fracture of the shaft of a bone can be more easily found by making comparisons between the same bones on both sides of the body, as the femora, humeri, and tibiæ. Likewise comparisons should be made between similar joints, especially the shoulder, elbow, hip, knee, and finger joints, to determine whether the deformity be a fracture or a dislocation. Fractures and dislocations of the neck are frequently overlooked, owing to lack of careful examination of these parts.

*Dislocations.*—Dislocations are also common sources of deformity. They may be: (1) Congenital, as illustrated in club-foot. (2) Traumatic, resulting from direct or indirect violence or muscular action, are the dislocations commonly met with. (3) Pathologic dislocations, due to degenerative changes in the joint, as occurs in tabes and Charcot's joint. Recent dislocations rarely show inflammatory changes, whereas older ones present evidences that such changes have occurred. Contractures are caused by nervous diseases, cicatrices with loss of skin and subjacent tissue, burns, and other accidents. These contractures are occasionally due to spontaneous dislocations. In old traumatic cases ankylosis is sometimes present. Ankylosis also occurs in pathologic conditions.

It may be important to examine the hyoid bone and laryngeal cartilages for fracture, dislocation, or laceration. In a case recently tried in the New York courts a patient was supposed to have received harsh treatment from an attendant in one of the hospitals there. The case hinged on determining whether a fracture of the hyoid bone had or had not been produced during life.

*Pathologic Deformities.*—Such distortions are due to diseases which may produce certain changes in the bone structure. The most common of these disorders are rickets, tuberculosis, syphilis, osteomalacia, acromegaly, and osteitis deformans. Rhachitis is a general cause of many varieties of deformity. In this disease the bones lose their tenacity and hardness, change in consistency (usually being thickened and spongy), and become distorted by the action of the muscles. These processes produce certain deformities: (1) Of the extremities, bow-leg (*genu varum* or *extrorsum*), knock-knee (*genu valgum*). (2) Of the sternum and ribs, pigeon-breast (*pectus carinatum*),

funnel-breast, Harrison's groove, beaded ribs. (3) Of the cranium, the square-box rhachitic skull, in the bones of which may be found spots of craniotabes. (4) The subject may be more or less dwarfed. These are the most common malformations.

Tuberculosis is often associated with a long, narrow chest, and is a source of common deformities, seen in coxalgia, Pott's disease, knee-joint disease (hydrops articuli), and the various grades of spinal curvature. Syphilis in its secondary and tertiary periods may produce nodes or cause great destruction of bone tissue. This process is present whether the malady be of the inherited or acquired variety. Osteomalacia causes bone softening, which may be followed by various deformities, especially fracture. This disease usually occurs in women after pregnancy. Osteitis deformans also produces changes in bones, usually those of the extremities. Certain chronic lung conditions result in contraction of one side of the chest and often a corresponding scoliosis; emphysema of the lungs is accompanied by a barrel-shaped chest; nervous diseases lead to more or less disfigurement of the body, as facial paralysis and spastic paraplegia; pernicious anæmia may either cause or be associated with spinal deformities; pleurisy may give rise to unilateral enlargement of the chest; aneurism may produce protrusion of the sternum. (For a further discussion of the changes found in bones and joints, see Chapter XVII.)

*Tumors* and other abnormal growths are also a common source of marked deformity. While deformities of the bones are the most frequent and conspicuous, yet there are other acquired deformities which are quite important. Congenital fissures of the neck which are tubular and go to the thyroid cartilage and the tuberculous perianal fissures should be followed out by careful dissection to their point of origin.

*Muscular Deformities.*—These are most generally due to muscular weakness. They may coexist with bone deformities and even cause or be caused by them; as deviation of the spine due to bone deformity destroys the harmony between the dimensions of the bones and the muscles, some muscles become elongated and others shortened.

*Acquired Deformities.*—Nature itself may produce deformities, as those arising from age, habits, and occupations. Notwithstanding that persons assume particular positions most constantly in certain occupations, they do not often acquire deformities.

*Tophi.*—These deposits occur in gouty persons, and are generally found in and about joint-cavities, ligaments, tendon-sheaths, arylte-



noids, epiglottis, vocal cords, skin of the penis, helix of the ear, tarsal cartilages, and margins of the eyelids. They contain a urate and a biurate, both of which dissolve in either acetic or nitric acid and give the murexid test for uric acid. If large and advanced they leave a white chalk-line when rubbed. Superficial tophi are movable and the skin over them is non-adherent, but as the process advances the mass adheres and may finally protrude. They are easily differentiated from sebaceous cysts.

**HERNIA.**—Hernia being of common occurrence and a frequent cause of death, the various situations where this defect may occur must be thoroughly inspected. The abdomen, inguinal canals, femoral openings, and umbilical region should be carefully palpated. The scrotum should be examined to determine the absence of one or both testicles, and when these are not found search for them should be made in the canal and elsewhere.

**EYES.**<sup>1</sup>—Inquire whether the eyes and mouth were open or closed when death occurred, and whether the expression was peaceful or the countenance distorted. While in life expression is manifested mostly by the eyes and the action of the facial muscles, in death the eyes lose their lustre and fail largely to influence the expression.

**Abnormalities.**—The eyes should be carefully examined in every postmortem, as abnormalities are quite common. The eyelids may be wholly or completely absent. They may not be divided or a fissure may exist involving one or both lids. There may be eversion or inversion. Ptosis, either acquired or congenital, may be present; if acquired, it may be due to specific causes. One or both eyes may have been replaced by artificial ones. The eye sometimes appears intact where sight had been absent for years before death, in which case there is always evidence that blindness existed. Abnormal prominence is usually caused by cavernous aneurism or exophthalmic goitre; the former may be associated with arteriosclerosis, the latter with enlargement of the thyroid gland. The presence of puffiness about the eyelids should be noted, as it occurs in Bright's disease, cardiac affections, arsenic poisoning, and traumatism.

**Position.**—Instead of presenting parallel visual axes, one or both eyes may be deviated inward, outward, downward, or upward, con-

---

<sup>1</sup> Much interesting material on this subject will be found in the *Pathology of the Eye*, 1904, by J. HERBERT PARSONS.



stituting one of the various types of strabismus, a condition which may aid in the diagnosis of toxic, cerebral, or nervous disorders. Conjugate deviation of the eyes occurs in apoplexy.

*Color.*—The color of the eyes is due to a blending of factors, varying in value in every case, depending largely upon the quantity of pigment present. The several races have, as a rule, characteristic colored eyes: the negroes and the Mongolians, black; the Australians and southern European races, dark; the Slavs, the Teutons, and the Scandinavians, blue. These peculiarities are worthy of note, as they may be of importance for purposes of future identification.

*Conjunctiva.*—Whitish patches, which may be congenital or acquired, are occasionally seen on the conjunctiva. If congenital they may be associated with moles on the face. The conjunctiva may be the seat of inflammatory conditions, which may be local or associated with some systemic disease. To note the variety of conjunctivitis present is of some importance, and if any of the severe forms should be suspected an effort should be made to ascertain whether or not it is specific. Ecchymosis of the mucous membrane may occur in cases of injury to various parts of the eye, traumatic conditions affecting the skull, dura, or brain, and even systemic disease itself. The conjunctiva is one of the seats of jaundice, and it is the place where jaundice shows itself most plainly in the negro.

*Pupils.*—Accommodation, convergence, and exposure to light, which during life produce alterations in the size of the pupils, after death do not affect it. In life, age, the color of the iris, and the character of the refraction also influence it. Under ordinary circumstances the pupils should be equal, but variations may occur, depending upon the conditions and cause of death, a few of which will be here mentioned. In fatalities due to chloroform the pupils may remain enlarged; in opium poisoning they often expand shortly before or after death; and in cerebral hemorrhage they are generally irregular, depending upon the location of the clot. The pupil can be measured approximately by holding in front of it a rule marked in millimetres and noting the number of spaces its width occupies. This method is somewhat faulty and only approximately measures the width of the eye, but an accurate measurement can be made with the pupillometer.

*Cornea.*—Note should be taken of the condition, curvature, and transparency of the cornea. Keratitis, ulcerations, and abscesses are common diseases of this locality. In old persons the arcus senilis is

usually present. Besides its liability to disease, the cornea may be lacerated, torn, or injured, with or without the lodgement of foreign bodies.

*Sclera*.—Examine its surface as to continuity and describe any lacerations or injury which it may have received.

*Iris*.—The iris should be inspected for color, condition, and quantity of pigment. As this is a common seat of disease and operations, it should be especially examined for the presence of a coloboma, one of its most common malformations, which may be either congenital or acquired. The congenital form is due to the failure of the ocular fissure to unite; it may be distinguished by the presence of the sphincter, which in the acquired form has been excised along the margin of the coloboma, as after an iridectomy. The fissure is usually situated in the lower part of the iris, and is often associated with coloboma of the choroid. The iris should also be examined for the scars of operations, for the information thus obtained is of value.

*Crystalline Lens*.—Luxation or subluxation of the lens should be looked for. If present it may be either congenital or acquired. Coloboma of the lens is accompanied by a similar condition of the choroid or iris.

*Optic Nerve*.—This portion of the eye together with the retina can best be examined with the ophthalmoscope.

*Growths*.—The most common growths of the eye and the parts generally affected are: Iris: angioma, metastatic sarcoma, usually from the ciliary body, granulomata, and cysts. Choroid: sarcoma, most common of all tumors, metastatic carcinoma, occasionally found, naevus, rare, cysts, rare. Ciliary body: sarcoma, common, adenoma, occasional, carcinoma, occasional, naevus, cysts. Retina: glioma and cysts.

Meningocele and herniæ of the brain containing cerebrospinal fluid may be found protruding from the sinuses into the orbit. Dermoids of the orbit are frequently discovered, especially near the eyebrows; those of the eye occur at the corneoscleral junction. The so-called carcinoma originating from the lachrymal gland is usually an adeno-sarcoma. Lipoma of the eye, which may be either congenital or acquired, occurs in the fatty tissue. Tumors originating from the bone are generally sarcomas or exostoses.

*Orbital Injuries*.—As injuries are frequently received in and about the orbit, careful examination of this region should be made. Frac-

tures through the orbit may cause, besides serious damage to the eye itself, grave cerebral complications.

*Orbital Diseases.*—Diseases of the orbit are quite common and may be important, for they often cause meningitis. Caries, necrosis, and cellulitis are generally preceded by periostitis.

After thorough examination of the eyes they should be carefully closed.

*HAIR.*—Examination of the hair may prove, especially in medico-legal practice, to be of importance. Not only the hair on the cadaver but also any hair found in the immediate vicinity of the body should be examined. Hair not belonging to the corpse demands inquiry as to whose it was and whence it came. In this way observations of value have been made and aided greatly in unravelling some of the world's deepest mysteries. The hair varies in color, length, quality, and quantity in different individuals, and also according to situation on the same person. The head of the new-born infant is covered with fine downy hair, a growth of the last two or three months of intra-uterine life. Shortly after birth it is shed and replaced by the true hair. The hair is one of the last tissues to yield to decay. The question of the growth of hair after death is a disputed one. Such apparent growth is most frequently caused by the retraction of the fat. In the new-born there is no medullary canal in the hair. (Vibert.) Human hair can be positively identified as such.

*Color.*—The color of the hair should be noted and described; also observe whether or not the color is uniform. As it depends mostly upon pigment, the color will vary in proportion to the quantity and variety of that pigment. Gray hair in adults is attributed to a diminution in pigment, and may be due to age, care, mental worry, sudden fright, burns, local inflammations, systemic diseases, nervous disturbances, hardships, or exposure to cold, as seen in Arctic explorers. Gray hair in the infant is congenital. Abnormal whiteness of the hair is a condition found in albinos. It may be complete or partial and is associated with loss of pigment in other organs. The examiner should not be misled by dyes.

*Length.*—The length of the hair should be observed and approximately measured. Long or short hair is characteristic of sex and of certain races. The longest hair is seen in the Indian, Chinese, and Malay; short hair in the Negro.

*Quality.*—Various races have hair of characteristic texture. The



Negro and the Bushmen have crisp, woolly hair; among the Anglo-Saxon, Germanic, and kindred races the hair is smooth; Australians have soft, smooth, wavy hair; the American Indian has coarse hair.

*Quantity and Distribution.*—Hairs normally may be present on all exterior parts of the human body, except the palms of the hands, soles of the feet, glans penis, mucous membranes, and the ball of the eye. Some races are prone to excessive hair growth, as the "hairy men" of the Island of Yesso. Loss of hair, complete or partial, may be due to depilatories, pregnancy, disease, or pressure. Baldness or moth-eaten appearance of the head, eyebrows, and moustache is seen in lues and myxœdema. Epileptic, idiotic, and insane persons generally have large growths of hair. Abnormalities of distribution have been recorded, and the examiner may often find either absence or overgrowth in certain localities. Under these conditions hair may be found in the interior of organs and neoplasms, especially ovarian dermoid cysts.

*Diseases.*—The hair of the body is subject to various diseases, and therefore its condition should be carefully noted. Various forms of alopecia, tinea, and fungi may attack the hair. These conditions may be due to local or constitutional disorders.

*Injuries and Tumors.*—Located under the hair tumors and various injuries may be present. Therefore pass the fingers through the hair of the scalp, and if it be at all thick part it, which aids in the discovery of wounds, hæmatomata, and tumors which it may conceal. Should any be discovered, cut or shave the hair so as to examine them more carefully. If the head has been injured, it will usually be found that the hair has already been removed by the surgeon.

*NAILS.*—When examining the nails attention should be paid to the material found under them, as it quite frequently is of medico-legal importance. The growth of the nail is regarded as one of the diagnostic signs of fetal maturity.

*Anomalies.*—The nails may be absent, atrophied, hypertrophied, brittle, discolored, cracked, etc. Congenital absence of one or more nails is usually associated with other malformations. Acquired anonychia may be due to trauma, nervous diseases, pyogenic infections, scar-leaving affections, as syphilis, and blood-stasis, as in cyanotic conditions. Another anomaly often met with is imperfect nail formation, resulting from dystrophia, in which the nail is usually opaque, discolored, brittle, and fissured. Certain diseases are sometimes noticed about the nails, as abscess, eczema, psoriasis, paronychia, syphilis, and



professional dermatitis with paronychia. Traumatism confined to the nail or surrounding tissues is frequently met with and its nature and extent should be noted.

**TEETH.**—The teeth should be examined as to anomalies, condition, and disease.

*Anomalies.*—One or more teeth may be permanently absent, or supernumerary teeth may be present to such an extent that there are two dental arches in either one or both jaws. The teeth may be irregularly placed, often beyond the alveolar process. In examining the teeth notice should be taken of the condition of the palatal arch.

*Condition.*—Much can be learned from the condition of the teeth, as to care, neglect, habits, and disease. The teeth should be examined to see whether they are artificial. It is a noticeable fact that people from certain countries have particularly fine teeth regardless of the care taken of them. Caries of the teeth and the extent to which it involves the bone should be noted and, if possible, the cause determined. Among other conditions phosphorous poisoning produces necrosis of the teeth and maxilla. Hutchinson's teeth are frequently seen and the condition is one of considerable importance. It consists in a single deep notch of the permanent upper central incisors, but the deformity is sometimes present in the molars when it is absent in the incisors. Dental tumors, such as epulis, sarcoma, osteoma, odontomata, or dentigerous cysts, are occasionally found.

•






FIG. 54.—Scheme to show the lines for incising the trunk and exposing more fully the abdominal cavity. The primary incision, *A B*, from the interclavicular notch to the symphysis pubis, goes to the left of the umbilicus at *C*, or it may pass in a straight line to the left of the navel. The transverse incision, *D E*, and the cutting of the rectus muscles at *F G*, are for the purpose of enlarging the abdominal




FIG. 55.—Elliptical incision for examination of the abdominal cavity; sometimes useful after abdominal operations and in infants who died from inflammatory conditions of the umbilical vessels.




FIG. 56.—Incision sometimes useful after death following operation in the abdominal region.



FIG. 57.—Method of making the initial incision over the sternum, as seen from above.



FIG. 58.—The same incision as in Fig. 57 somewhat extended, as seen from the side.





FIG. 59.—Method of raising flap on right side so as to expose sternum and ribs, as seen from above.



FIG. 60.—Same incision as in Fig. 59, seen from the side.



FIG. 61.—Method of opening the abdominal cavity. The index and middle fingers being stretched apart, the abdominal wall is lifted off the viscera so that the cutting may safely be done from above downward.



FIG. 62.—On the right side the cartilages have been cut through near their junction with the osseous portions of the ribs. On the left side the fifth rib is about to be cut through, the knife being introduced below it and the cutting done upward, using the sixth rib as a fulcrum.



FIG. 63.—Method of separating the sternoclavicular attachment. The articulation is discovered by steadying the sternum with the right hand and moving the clavicle in various directions with the left hand. Unless marked ankylosis exists, incision with the knife is easy after the discovery of the articulation.



FIG. 64.—The incision through the first rib is here shown. The previous incision seen in Fig. 63 has been carried almost to where the thumb is located, the clavicle elevated, and the incision made through the first rib.



## CHAPTER VI

### TECHNIC OF OPENING THE ABDOMINAL CAVITY AND THE TOPOGRAPHIC EXAMINATION OF THE PARTS CONTAINED THEREIN

AFTER the completion of the external examination of the body, all necessary instruments are placed in order upon a board or tray,<sup>1</sup> and the operative part of the autopsy is begun.

The operator should stand so that when facing the body the incisions from above downward can be made with the greatest ease. This condition is best fulfilled by right-handed operators standing on the right side of the supine object, and left-handed operators on the left side. The body should be drawn well to the side of the table nearest the operator, the head resting on the top of the table and, if preferred, the shoulders supported by a block.

With the knife held in the manner previously described (page 36), as nearly horizontal as possible, a clean incision (Fig. 54) should be made by a single sweep of the arm from the interclavicular notch (*A*) to the symphysis pubis (*B*), passing to the left of the umbilicus (*C*) in order to avoid the round ligament and any vessels going to and from the navel, care being taken not to penetrate the abdominal cavity and thus injure the contained viscera, or to extend the incision to the external genitalia so as to disfigure them (Figs. 57 and 58). On the chest this primary incision goes down to the sternum, whereas on the abdomen it penetrates only to the muscle-sheath. In Europe the initial incision is usually made at the middle of the chin,—*i.e.*, starting at the symphysis menti and ending at the symphysis pubis; for there, as a rule, only the poor who die in the hospitals reach the post-mortem table, autopsies being seldom performed on the bodies of persons belonging to the upper classes, who would naturally object to the disfigurement entailed by this method. In this country the longer incision should be used only when great haste is necessary, as in cases of contagious diseases, such as diphtheria,

---

<sup>1</sup> A towel may be laid over the external genital organs and the upper parts of the thighs, upon which the instruments to be used in the performance of the autopsy are placed, with their handles towards the obducent. An ordinary stool is frequently used abroad for this purpose.



or when the body is not to be seen again by relatives or friends. If the mentopubic incision be employed, it should not injure the thyroid gland, the omohyoid being the only muscle to be cut through. The knife now follows underneath the skin, fat, and fascia along the lower border of the inferior maxillary bone, until the digastric muscle and the submaxillary gland are seen. The gland is then incised, the muscles being left *in situ*. The common carotid artery, internal jugular vein, and pneumogastric nerve are now readily exposed. Care should be taken to avoid wounding the vein, as the resulting hemorrhage hinders subsequent manipulations; such injury is especially to be guarded against on the left side in those cases where search is to be made for the entrance of the thoracic duct into the subclavian vein.

The initial incision over the thorax is now carried down to the sternum in any place where this has not already been done, and the layers of skin, fascia, and muscles of the right side are quickly dissected away close to the ribs, freely exposing the costochondral articulation and some three inches of the sternal end of the clavicle. To do this the attachments of the soft tissues are made tense by drawing them away from the median line with the left hand (Figs. 59 and 60), while long sweeping incisions are being made downward and laterally with the large section-knife. The left side may next be similarly treated, though in practice this is more frequently done after the abdominal incision has been completed. Some permit an assistant to prepare the left side while the operator is uncovering the right, but the time saved by this procedure is small and the danger of injury to those taking part is great.

Should a pneumothorax have been diagnosed during life, the thoracic pocket made by elevating the skin-flap on the side of the pneumothorax is filled with water, and a puncture is made at the bottom through the costal pleura at the intercostal space between the sixth and seventh ribs at the axillary line. If a pneumothorax be present, bubbles of air will escape through the water, the visible supply of which will rapidly diminish. If the head be lowered and enough water be used, this will finally escape from the mouth. It should be remembered, however, that a cavity in the lung opened accidentally by the knife would give the same result as that arising from a pneumothorax.

A note is now made as to the situation and character of any blood which may escape. The condition of the fat (*panniculus adiposus*)

is considered, and its thickness noted at the nipple-line, beneath the xiphoid, and again three inches below the umbilicus. In atrophy the color of the fat becomes darker, changing to orange or reddish yellow. As a rule, the older the individual the darker in color is the fat, varying from straw-color in children to the dark yellow seen later on in life. Different species of animals have different colored fat, depending upon the difference in its chemical composition. The tissues here are often oedematous, as in general dropsy or erysipelas.

The mammary glands may now be examined from behind, and, if desired, the glands of the axillæ may be exposed by continuing the dissection of the pectoral muscles from beneath. After the reflection of tissue over the ribs, the mammary gland on each side may be opened by four or five parallel perpendicular incisions, followed by several transverse ones, if deemed necessary. When the gland is quiescent, it will be found to consist of dense, whitish, fibroid tissue, in which are a few pin-point, grayish-red spots (breast-grains). During lactation it is larger, and the secreting tissue grayish red in color and granular, looking like a salivary gland. (Orth.) Fibrous tissue is found only near the nipple. Pressure may cause an outflow of a yellowish creamy colostrum (which is not altogether unlike pus in its appearance); this should be at once examined under a low-power of the microscope. The following conditions of the gland should be thought of:

1. Deformities: (*a*) depressed nipple; (*b*) mamma infantilis (hypoplasia); (*c*) supernumerary glands or nipples. (2) Congestions and anæmias. (3) Burns. (4) Inflammations: (*a*) inflammation of nipple; (*b*) fissures; (*c*) mastitis, acute; (*d*) abscess of connective tissue: (*a*) between skin and mamma; (*β*) between muscle and mamma; (*γ*) acini and ducts. (*e*) Fistula: (*a*) soft edge may mean milk fistula; (*β*) indurated edge may mean mammary abscess; (*γ*) if broken and caseous, it is more likely tuberculous. (5) Granulomata, gummata, tuberculosis, caries of ribs. (6) Changes of nutrition. Remember in this connection that tumors of the genitalia or pseudopregnancy may cause an hypertrophy of the gland with a true secretion of milk, and that a similar result may occur in rare cases in the male apparently from certain psychic influences. (7) Tumors, fibroma, carcinoma, echinococcus cysts, etc.

The muscles now to be examined are those of the neck, chest, and abdomen. The external examination notes any marked changes in

bulk. Both a transverse and a longitudinal section are necessary to a complete study, and the general characteristics should be observed. The muscles may be atrophied or hypertrophied. Trophic change induced by affections of the anterior horns of the spinal cord may show itself in muscular atrophy and may be either inflammatory or degenerative, as in infantile paralysis, progressive muscular atrophy, and amyotrophic lateral sclerosis. The lesions may chiefly affect the peripheral and intermuscular nerves, as in lead paralysis with atrophy and in certain atrophies following diphtheria and other exanthemata, or the muscles may be primarily affected, as in the juvenile form of Erb's paralysis and pseudohypertrophic paralysis, or reflexly, as in the Charcot joint-affections. (Dreschfeld.) Hypertrophy due to exercise increases the number of muscle-cells; when due to an increased blood-supply, the individual fibres are increased in size. Muscle is a highly organized tissue, and as such does not reproduce itself with ease after injury.

As the color of a muscle is due to hæmoglobin or some modification of it, its appearance will vary according to the state of the blood. Normally muscle is a bright red, but in anæmia it becomes paler, at times a grayish red. In general it may be said that the color and consistence of the muscles bear a distinct relation to each other: pale muscles are usually soft, while the darker muscles are more firm. The muscles are dry when much fluid has been carried off by the alimentary canal, as in typhus fever and cholera, and moist after the occurrence of disturbances of the circulation. Zenker in 1864 described a form of colloid degeneration resembling the flesh of fish in the flat muscles of the abdominal walls, occurring especially in enteric fever, though found in tetanus, scarlet fever, smallpox, and near sarcomatous tissue. In diseases where the muscles have long been inactive, a similar grayish translucent appearance is at times observed. The dark meat of the fowl undergoes decomposition sooner than the white. The flesh of different animals possesses characteristic odors. Embalming fluids containing zinc bleach the muscles, while arsenic preserves their natural color. Formalin hardens them.

The general characteristics having been observed, the pathologic conditions to which these muscles are subject are not liable to escape detection. The more important morbid lesions are:

(1) *Hemorrhages*.—These may result from trauma, wet cups, hypodermic injections, etc. The outflow from the cut veins often



gives a good idea as to the color, fluidity, and odor of the blood. A special form of bleeding into the rectus may occur in typhoid fever, the so-called "hæmatoma recti abdominis."

(2) *Inflammations*.—Among these are included: (a) *Acute Myositis*.—This is often suppurative, and may be primary, from trauma, or more usually secondary, in the muscles of the chest, to pleural affections, or, in the muscles of the abdomen, to pelvic suppuration. This inflammation does not, as a rule, produce true abscesses, but infiltrations in the muscle and separation of its fibres, which undergo a fatty or hyaline degeneration. Hæmatogenous inflammation is by no means uncommon, perfect examples of miliary tubercles often being found if searched for in suitable cases. (b) *Chronic Myositis*.—The interstitial connective tissue is increased so that at times it is visible to the naked eye, the muscle-fibres are atrophied, the color becomes a grayish red, and the muscles feel solid. This condition is generally associated with diseased states of the neighboring parts,—e.g., affections of the ribs, pleuræ, cervical glands, etc. There is a syphilitic form of fibroid myositis. Glanders and actinomycosis may affect the muscles. (c) *Parenchymatous Myositis*.—The muscle is paler than normal. All the various forms of degeneration—as cloudy swelling, hyaline or fatty—affect muscle, and microscopic examination, as in acute primary polymyositis, is necessary in order to determine their presence. (d) Bony formations are sometimes found, as the "drill" bones in the shoulder muscles and the "riders" bones in the adductors of the thigh. (Ziegler.) Progressive ossifying myositis is a rare disease, running a chronic course, which especially attacks young people; those thus affected are sometimes exhibited in dime museums as "petrifying" persons.

(3) *Parasites*.—The most important parasite is the *Trichina spiralis*, which is found most frequently in the muscles of the neck and in the intercostals near the attachment of the diaphragm, and in old cases the calcified capsules may be easily recognized as small, white, oval bodies, which when present in large numbers look and feel like grains of sand. In the muscle itself the site of election is close to the spot where the tendon unites with the muscle proper. In order to see the parasite the capsule should be dissolved with hydrochloric acid. In its early stages the animal is not readily discovered, and its detection is made easier by pressing a teased portion of muscle between two glass slides and observing it by transmitted light with a hand lens.



In all doubtful cases the aid of the microscope should be evoked. Measles and hydatids may also be found in the muscles.

(4) *Tumors*.—Primary tumors are rare: they usually originate from the connective-tissue septa.

The abdominal cut is now deepened between the umbilicus and the xiphoid cartilage until a small portion of the peritoneum is exposed.<sup>1</sup> This membrane should then be carefully opened and if it be desired to determine the presence and character of any gas present in the abdominal cavity, the incision is made down to the peritoneum, either two inches above or the same distance below the umbilicus, and the abdominal walls are elevated with the fingers or a tenaculum so as to form a pouch, into which water is poured. A test-tube is then filled with water and inverted over the pouch, and a small incision is made through the peritoneum under the mouth of the test-tube so as to allow any gas escaping to go into it. The test-tube is tightly closed before all the liquid has run out of it, by pressing a thumb or finger up against its mouth, and placed in a shallow dish containing sufficient water or mercury to seal the open end of the tube. It is then handed to the chemist for examination. If a lighted match be held close to the point where a knife is pushed into the chest, any gas escaping deflects the flame. It should be remembered that certain gas-forming organisms may be the cause of the gaseous collection in serous cavities. The recent discovery of hitherto unknown elements in the air makes the study of aggregation of gases here an extremely interesting one. If the gas has an acid odor, an opening in the stomach is to be suspected.

If fluid be present, the abdomen usually protrudes, the sides are flattened, and the superficial veins much dilated, a caput Medusæ forming about the umbilicus. Ballotement will often reveal its presence, and when found a mental note should be made carefully to examine the œsophageal veins, as a fatal hemorrhage may occur from their rupture, a fact which I have more than once personally substantiated *post mortem*. In ascites just enough of the fluid should be removed to facilitate the determination of the height and location of the diaphragm, which may be done by introducing the hand, palm upward, or a steel sound, into the abdominal cavity and following the under surface of this muscle as far as possible. When the tips of the fingers or the end of the sound reach the point of least resistance, this spot

<sup>1</sup> For the technic of a bacteriologic examination see Chapter XXIII.

should be sought for with the other hand from without. The vault of the diaphragm extends to the fifth rib on the left side and to the fourth rib or fourth interspace on the right. Both sides are measured in the line of junction of the ribs with the costal cartilages. The greater height on the right is due to the liver, which forces the diaphragm upward, and in excessive hypertrophy may cause it to reach even as high as the level of the second rib. Increase in the abdominal contents, as by tumors, pregnancy, hypertrophy of the spleen, etc., elevates the diaphragm, while augmentation of the thoracic contents naturally pushes it downward. Along with the depression is a sense of fluctuation in cases of hydro- or pyothorax. The position of the diaphragm in a new-born child helps to determine whether or not it has breathed. Before respiration has occurred, the summit is found on a level with the fourth rib on the right side and on a level with the fifth rib or the fourth intercostal space on the left. After full expansion of the lungs has taken place, the summit is found at the fifth or sixth rib on the right and at the sixth rib on the left (Orth).

The opening may now be somewhat enlarged and additional fluid removed with a syringe, cup, or large pipette, measured, and its character noted. The remaining portion may be collected from the various folds and pouches in the peritoneum with a sponge or small cup. The amount of fluid normally present is very small, not usually exceeding a teaspoonful; it may be lemon-yellow, red, or brown; icteroid or milky; watery, thick, gruel-like, or even semisolid. The removal of liquid at this stage of the operation prevents its admixture with blood, as from an accidental incision into the liver while cutting the costal cartilages, or with other fluids of the body, such as those from the pericardium, the pleura, the bladder, or various portions of the intestinal tract. Ascites is especially associated with Bright's disease, chronic heart disease, chronic lung disease, anæmia, portal obstruction due to cirrhosis of the liver, chronic passive congestion of the liver, inflammatory adhesions, etc., tumors, displaced or hypertrophied viscera, as the enlarged spleen of malaria or leukæmia, and peritonitis, especially when tuberculous. The serous membrane is apt to be lustreless, whitened, and thickened, especially the capsules of the spleen and liver, if the disease has lasted any length of time. The intestines are frequently matted together by fibrous adhesions, and the uterus and adnexa often show a similar condition, especially when the peritonitis is of a tuberculous nature.

In cases of increased amount of fluid it is of importance to distinguish between a serous transudate and an inflammatory exudate. When large amounts of pus and fibrin are present, the differentiation is easy, as a transudate contains neither. Difficulty arises when a clear watery fluid is found in which minute flocculi are seen, as these may be either small flakes of fibrin and pus-cells or collections of washed-off endothelial cells. The differential points are as follows:

TRANSUDATE.	EXUDATE.
Fluid clear and watery, though it may form a spontaneous clot.	Fluid thick, ropy, and at times foul smelling.
Alkaline reaction.	Acid reaction.
Flocculi are thin, veil-like, and of a transparent gray color.	Flocculi are thick, opaque, and of a grayish-white color.
Specific gravity usually below 1016.	Specific gravity generally over 1016.
Albumin usually below two per cent.	Albumin may exceed three per cent.
No bacteria or their products present.	Contains organisms, toxins, globulicides, etc.
Urea <sup>1</sup> and cryoscopic index low.	Urea and cryoscopic index high.
Under the microscope the flocculi are seen to be made up of flat cells with large nuclei and lymphocytes.	Microscope shows the flocculi to consist of fine threads and polynuclear leucocytes, the nuclei of which appear more distinctly on the addition of acetic acid.

Milky exudates are of two kinds, fatty and chylous. The former excretion has been found in connection with peritoneal and mesenteric cancer, and is recognized by the fat-globules seen on microscopic examination. Slight amounts may be due to the fact that the patient was on a milk diet or was suffering from lipæmia, a dyscrasia also found in diabetes (Osler). A chylous exudate results from the perforation of the thoracic duct or the receptaculum chyli.

Suppurative exudates, due to perforation of the intestine, are thick, yellowish, and contain much fibrin, which is deposited on the peritoneum and bowel in layers. The odor, which is peculiarly nauseating, may be due to the *Bacillus coli communis*; the process is usually acute.

A hemorrhagic exudate or fluid may be non-inflammatory, as that arising from trauma (rupture of the liver or spleen or extra-uterine pregnancy), from cirrhosis of the liver, from cancerous and tuberculous peritonitis, etc., or it may be inflammatory. Pure bile, most frequently mixed with blood, may be found in the abdominal cavity after injury to the gall-bladder or the bile-ducts.

---

<sup>1</sup> ULRICI, *Centralbl. f. innere Med.*, no. 16, 1903.

One finger is now introduced into the opening previously made in the abdominal cavity, the flap of the skin is elevated, and the incision is somewhat lengthened. Next the index and middle fingers of the left hand, held V-shaped (Fig. 61), palm upward, are thrust under the abdominal wall in order to raise it above the intestines so as to prevent injury to them in the subsequent incision, the fingers acting as a director while the cut is continued to the pubes. Then a similar incision is made up towards the xiphoid cartilage. If there be much meteorism, the index-finger of the left hand can be introduced and held against the parietal peritoneum. If scissors be used, the lower blade may be guarded by the fingers of the left hand when the cut is made. Another method is to make the incision while the part is well elevated above the intestine by strong traction upward on the right abdominal flap. The cutting should preferably be done from within outward, great care being taken not to puncture or injure any of the abdominal viscera, especially the stomach and bladder. After noting the location and distribution of adhesions as pointing to previous inflammatory conditions, it is well to break up such adhesions with the fingers. Should the intestine be accidentally opened, it is best to stop at once and tie both above and below the opening in order to prevent the escape of the contents of the bowel into the peritoneal cavity.

If it be desirable to enlarge the opening in the abdominal wall (Fig. 54), a second incision (*D E*) may be made, at right angles to the first one and about three inches above the umbilicus, or the rectus muscle on one or both sides of the body may be divided subcutaneously a little above Poupart's ligament (*F* and *G*). Should there be a penetrating wound of the abdomen, as from a dagger or a previous *cœliotomy*, the abdominal incision may be changed at will (Figs. 55 and 56).

When the contents of the stomach are found in the peritoneal cavity, care must be taken to determine whether their escape was due (*a*) to post-mortem digestion, or *autopepsia*, (*b*) to trauma, (*c*) to perforation from a gastric ulcer or from chemical erosion of the coats of the stomach by poisons, etc., (*d*) to the presence of (*b*) and (*c*), with the factor (*a*) as the real cause. In the first case the ingesta are usually widely distributed throughout the abdominal cavity, though most plentifully in the immediate neighborhood of the perforation, the rent is large and irregular, and the walls are soft and slimy; while in disease the opening is apt to be small and circular and surrounded by evidences of hemorrhage and peritonitis. Undigested food enters



the peritoneal cavity through a breach in the gastric wall; when digested food or fæces are present, the seat of injury is the bowel or duodenum, and, if the latter, the material is usually stained with bile. Autodigestion is especially frequent in cachectic children. Intestinal worms may escape into the peritoneal cavity through perforations in the bowel. From the fact that the autopsy is usually performed a considerable time after death, the appearances presented by a gastric ulcer are slightly different from those seen at an operation during life, for the serous wall of the ulcer may have undergone a certain amount of post-mortem digestion.

The suspensory ligament of the liver may be studied at this time. Should there be an omphalomesenteric duct, it should be carefully followed out to its diverticulum in the ileum.

Foreign bodies, which may be calcified, are sometimes found free in the abdominal cavity; they are derived from torn-off appendices epiploicæ or polypoid tumors. Surgical instruments and appliances, such as sponges, artery-forceps, scissors, and gauze compresses, have been discovered in the abdomen after the performance of operations, v. Neugebauer<sup>1</sup> citing 195 such instances. In Europe severe punishment has been meted out to surgeons for their forgetfulness in this respect.

The abdominal cavity being thoroughly exposed, the most striking abnormalities therein are to be noted. Transposition of the viscera would at once be observed. The most marked displacements of abdominal organs seen by the writer were in cases of Pott's disease and diaphragmatic herniæ.

The omentum ordinarily comes first under observation. Normally the omentum is non-adherent to the intestines except at its point of attachment; in purulent peritonitis it may be markedly adherent to the peritoneum covering the intestinal tract, creamy or plastic lymph appearing in streaks throughout its structure. The omentum may form a part of every variety of hernia found in the abdomen; it may be present alone in the hernial sac, or the intestines may become strangulated by passing through an opening in it. Trauma or atrophy of the connective tissue may produce such openings, some of which may be of large size. The amount of fat deposited between the layers of the omentum varies considerably, being in some cases practically

---

<sup>1</sup> *Centralb. f. Gynäkologie*, 1903, vol. xxvii, no. 8.

absent and in others measuring as much as half an inch in thickness. During health the omentum is rich in fat, which disappears early and rapidly in emaciation. Normally the layers are readily separable, and when spread out form a beautiful thin, transparent membrane, with irregular deposits of fat, and showing the blood-vessels partly filled with blood. It is a common seat of fat necrosis, tuberculosis, and generalized cancer; in the last two conditions it may be so contracted upon the transverse colon or the greater curvature of the stomach as to be hardly visible, and separable therefrom only with the greatest difficulty. Enlarged glands, encysted parasites, infarcts, pins, supernumerary spleens, etc., may be found.

The serous covering of the stomach and intestines should be minutely inspected, as the play of colors is very varied and the information gained from this examination is often of great importance. In thrombosis of the mesenteric vessels the gut may be gangrenous for ten feet or more. Miliary tubercles are found opposite tuberculous ulcers and extend along the lymphatics; they are also seen on all the other portions of the peritoneum, often being wide-spread in tuberculous peritonitis. Small yellowish, creamy collections of lymph, with dilated lymphatics, are seen if death occurred several hours after eating; these are physiologic and not pathologic products, but I have known them to be mistaken for miliary tubercles and even for carcinomatous growths. The presence of typhoid ulcers may be recognized by a congested area along the length of the intestine. The location of the vermiform appendix should always be noted, and Virchow's dictum (first published in 1875, though practised long previously) should be remembered: "At least in every case of inflammation of the peritoneum the appendix is to be carefully examined." In the female an inspection should be made of the uterus and its adnexa. The mesenteric glands, especially those near the ileocaecal valves, are to be carefully looked at; they are greatly enlarged in typhoid fever, in which they sometimes undergo suppuration, and in children dying from inanition, where they appear as red nodes, often running together into conglomerate masses.

The transverse colon may assume odd shapes and positions; thus, it may be bent like the letter U and extend as low as the bladder; it may or may not drag down the stomach. In some cases it forms peculiar S-shaped curves; in others the hepatic and splenic flexures may be markedly deficient. These malpositions are supposed by some

to be especially common in the insane. Cotton which has been inserted in the rectum or vagina by the nurse or undertaker to prevent the escape of fecal or other matter may be mistaken for a foreign body and may possibly have caused displacement of neighboring parts.

The stomach is subject to marked changes in size and situation, as from hour-glass contracture, tumors, ulcerations, etc. In the babe its situation is nearly vertical. This viscus is often filled with gas formed after death; a peculiar sound may sometimes be heard when the gas is expelled by pressure from without. One does not realize the extent to which the stomach may be distended by food and drink until he has made post-mortem examinations of the viscera of inebriates and persons accidentally killed soon after they had eaten hearty meals. The capacity of the stomach may be estimated by filling it with water and measuring the amount; but the method is not accurate and may destroy the appearance later on of a gastric ulcer.

All the openings in which herniæ are apt to occur are next to be examined, the most common varieties being inguinal hernia in the male and femoral and umbilical herniæ in the female. Other forms of rupture are those into the canal of Nuck, the obturator foramen, or the sciatic notch; into the various fossæ about the cæcum or the fossa jejunalis; into new fossæ formed by bands of adhesions, as from extra-uterine pregnancy; from solutions of continuity in the mesentery; crural; diaphragmatic, which is often congenital, but may be due to traumatism; between the rectus muscles and through Petit's triangle; after operations, especially those on the appendix, etc.

Volvulus and invagination are not infrequently seen. True invagination is to be distinguished from a form which often occurs in children just previous to death; in the latter cases multiple lesions (sometimes as many as fifteen or twenty), produced during the agonal period, are found. There is a peculiar form of invagination in which the ileocaecal valve draws the ileum down into the caput coli; this condition when extreme may even cause the ileocaecal valve to appear at the anus. Philipowiez<sup>1</sup> finds that volvulus of the sigmoid occurred in one-third of all such cases reported.

Note if the gall-bladder is distended or contracted; see if it extends below the liver, and, if so, to what extent. Feel it gently and note if any gall-stones are contained therein. Should a bacteriologic

---

<sup>1</sup> *Arch. f. klin. Chir.*, 1903, vol. lxx, nos. 3 and 4.

examination of this part be wanted, it is now to be made. Follow with the hand the upper surface of the liver, first of the right lobe and then of the left, in order to determine their extent, noting the height and the distance to which they extend below the ribs. The tips of the right and left lobes of a large liver almost meet at the vertebral column. The left lobe may extend downward like a beaver's tail, and as a result of tight lacing the whole organ may be divided into an upper and a lower portion by bands of connective tissue containing the biliary vessels and a few liver-cells. Extra lobes are very common; some of them even take the form of supernumerary livers. This condition may be congenital, but it is more frequently due to syphilis. In one of my syphilitic cases the liver was made up of more than thirty lobes, in shape resembling a bunch of flattened and distorted hydatid cysts. The liver should next be slightly raised, the pylorus examined, and the tips of the fingers used to determine the presence of calculi in the bile-ducts and gall-bladder.

When no extensive pathologic lesions exist, the situation of the pancreas may readily be determined by remembering the close connection of its head with the concavity of the duodenum.

During this superficial examination of the abdominal cavity any needful departure from the ordinary routine may be planned. Thus, in a case of cancer of the head of the pancreas it may be advisable later on to remove this organ along with the stomach, the duodenum, or even the liver. Again, in the case of a child or when there is not time for a careful dissection, all the organs of the abdominal cavity may be removed *en masse*. One must always be on the lookout for supernumerary organs, for they occur in the most unexpected places, as pulmonary tissue below the diaphragm, adrenal tissue in the liver, and pancreatic tissue on the wall of the stomach.

To repeat, the relative positions of all the tissues should be observed and any departure from the normal noted, and a careful search made for foreign growths, attachments, anomalies, etc., none of the parts being at this time removed from the body or their relations so disturbed as to prevent further examination.



## CHAPTER VII

### TECHNIC OF EXPOSING THE THORACIC CAVITY AND THE CRITICAL EXAMINATION OF THE PARTS CONTAINED THEREIN

**METHOD OF OPENING THE THORAX.**—After the superficial examination of the parts contained in the abdomen, the organs of the thoracic cavity may be exposed to view in the following manner. The second to the tenth costal cartilages on the left side are cut through, one by one, from above downward, at a point close to the attachments of the osseous portions of the ribs. For this purpose a heavy cartilage-knife is employed, which should be held as nearly parallel to the chest surface as possible, so that as the blade cuts through one cartilage it strikes the next one, thus preventing injury to the organs beneath. In order that the knife may not be dulled by this procedure, the cutting may be done by the part of the blade near the handle. Or the knife may be introduced into the intercostal space beneath the rib that is about to be cut, using the next lower rib as a fulcrum and cutting from within outward (Fig. 62). As the incision proceeds downward the ribs are severed more and more towards the axillary line, thus making the opening in the chest larger and larger. In cases where the cartilages are calcified it may be best to use a costotome or a saw for their division, in which event the ribs might as well be cut outside the costochondral junction in order to allow more room for subsequent manipulations. The second to the tenth ribs on the right side are now severed in a similar manner.

The right clavicle is next separated from the sternum. As its head articulates with the latter bone and the cartilage of the first rib, the collar-bone is grasped with the left hand and its inner end is moved to and fro, or an assistant may produce the same result by moving the whole arm. In this way the line of articulation is easily made out, and permits the part to be disarticulated by cutting downward and slightly outward until the first rib is reached (Fig. 63), thence continuing the incision outward along the under border of the clavicle and the upper part of the first rib for at least two inches. The first rib, which is generally calcified, is next cut through with a knife



FIG. 65.—Method of incising the first rib and the sternoclavicular articulation with the costotome.



FIG. 66.—All the ribs of the right side have been severed, the sternoclavicular attachment to the first rib remaining intact on the left side. The lower portion of the sternum is elevated and traction made on the diaphragm, which is cut as close as possible to the lower border of the sternum.



FIG. 67.—The lower border of the sternum having been freed, the breastplate is elevated and pulled upward and towards the left. The left sternoclavicular attachment is thus easily discovered, and is cut through. The first rib is then detached. Care is especially taken on this side not to injure the subclavian vein, not only on account of the blood escaping upon adjacent parts, but also owing to the difficulty after disturbance of the parts in finding the entrance of the thoracic duct into the vein.



FIG. 68.—The sternum is here practically ready to be removed from the body. The knife is cutting any attachments which may not previously have been severed in the neighborhood of the left sternoclavicular articulation.




FIG. 69.—Breastplate after its removal from the body. This is what is known in France as the chondrosternal plastron. Should the ribs have been cut through, it would be called the sternocostal plastron. The two internal mammary arteries will be found on the under surface. The retrosternal glands are also to be examined.



FIG. 70.—The skin flap is placed over the projecting margins of the right clavicle and ribs in order to protect the operator's hands from injury. On the left side this has not been done. On the right side a transverse incision, FIG. 64, *Z*, has been made, while on the left side the rectus muscle has been incised as in *C* of the same figure.





FIG. 44.—Method of opening pericardium. The left hand supports the right flap of the pericardial sac, while the knife cuts the pericardium up to its attachment to the great vessels coming off from the heart.



FIG. 75.—Method of removing the heart from the body. The index-finger is placed in the left ventricle and the thumb in the right ventricle, and the ventricular septum is grasped. The heart is then raised upward and towards the chin, placing on a stretch the blood-vessels which enter the heart. These are cut, starting with the lower pulmonary vein and going from left to right in a circular direction until the upper pulmonary veins are reached, or the initial incision may be made at the inferior vena cava and end with the pulmonary veins.

from below outward or from above inward (Fig. 64). Or the costotome may be employed for this purpose (Fig. 65).

The next procedure is accomplished by making traction on the breastplate upward and towards the right. Beginning below on the left side and keeping close to the lower border of the sternum, the underlying tissues, diaphragm, and mediastinum are cut through with short transverse strokes of the knife (Fig. 66), the sternum being elevated more and more as the tissues are separated. All the lower attachments having been cut and care having been taken not to open the pericardium, the breastplate is now elevated and any uncut tissues of the mediastinum and of the right side are incised. The elevated sternum is now to be pulled towards the left. Any sternocostoclavicular attachments on this side being made tense are easily discovered and severed, the knife passing along and beneath the upper part of the sternum (Fig. 67). The increased room and the greater leverage afforded by torsion of the sternum upon the left sternocostoclavicular attachment make this part easier of removal than on the right side. The tissues made tense by raising the sternum are divided, the bone being pushed more and more to the left and slightly rotated. The right first rib is cut through and the left clavicle is now disarticulated from below (Fig. 68). This procedure usually requires very little use of the knife, the force applied by the rotation often being sufficient for this purpose. The breastplate, after its removal from the body, is shown in Fig. 69. If an aneurism or tumor be found adherent to the ribs or sternum, its point of attachment is preserved by sawing through the bone at some distance therefrom.

In removing the sternum great care is necessary in order to avoid cutting the innominate or internal mammary veins which lie beneath its upper end and the clavicle. In Bavaria and Württemberg, in order that these vessels may not be injured and the part bathed with blood, the regulations for the performance of medicolegal autopsies direct that the lower end of the sternum, when freed, shall be strongly elevated, and the sternoclavicular connection and the first rib cut from the under side, or the breast-bone may be sawed through below the attachment of the first rib, leaving it and the sternoclavicular articulation intact. The writer does not approve of the method often used, after cutting the ribs, of breaking the sternum by turning it backward just below the clavicular attachment. Though it avoids injuring the veins, it leaves an ugly place from which to receive scratches and does

not give as much room for the subsequent examination of the thoracic cavity and neck. Some careless operators do not even remove the bone, but while still attached turn it back over the face. Do not forget to return the sternum to its proper place in the restoration of the thorax, an error often made and not discovered until the body is sewed up.

In order to protect the hands of the operator from future injury, the skin flaps are now wound around and beneath the exposed clavicle and ribs (Fig. 70), or these may be covered with a strip of antiseptic gauze held in place by a stitch around an upper and lower rib. Cotton should not be used for this purpose, as portions become detached and become adherent to the tissues of the body, thus interfering with future manipulations.

**STERNUM AND RIBS.**—The examination of the sternum and ribs may now be undertaken. Their shape is often altered, as in Pott's disease, pigeon-breast, emphysema, perforated sternum, rickets, occupation deformities, such as pressure atrophy in shoemakers, caused by holding the shoe against the breast, etc. Tuberculous caries of the sternum, often secondary to caseation of the mediastinal lymph-glands, or metastatic tumors may be present, or an aneurism may cause pressure atrophy (erosion) or even perforation of this bone. It is this form of saccular aneurism which is now treated by wiring and by electrolysis. Fracture is not common, but may occur between the second and third costal cartilages,—*i.e.*, near the junction of the manubrium with the gladiolus. The ensiform appendix of the sternum is sometimes curled upward and outward like a hook in cases of hepatic hypertrophy or tumor. Where this condition is present with atrophic cirrhosis of the liver, it indicates a previous enlargement of that organ (Suchard). The marrow of the sternum (best exposed by a longitudinal opening), which is normally of a slightly reddish, lymphoid appearance, may present the changes characteristic of leukæmia, anæmia, tuberculosis, etc. In the last stages of carcinoma the sternum and ribs are at times so infiltrated with cancerous deposits, especially when the breast is affected, as to break readily. The ribs may show evidence of rhachitis by the presence of the rhachitic rosary, in which case a section of the rounded enlargements, especially where the cartilage joins the bones, will show the changes peculiar to rickets. In old persons the entire cartilage may be calcified or even ossified. The central substance of the ribs sometimes undergoes atrophy and absorption, leaving a large canal filled with blood. The cartilage may contain



cystic cavities. Coplin has recently shown that the intercostal muscles are apt to show dissociation of fibres, leucocytosis, infiltration, granularity, etc., in disorders affecting the lungs and pleura.

The clavicle may now be freed almost to its acromial attachment, the arm extended at a right angle to the body, and the region of the subclavian vessels and brachial plexus readily exposed and studied. In one of my cases I found that the brachial plexus had been ligated by a competent surgeon in mistake for the artery. Should it be desired to remove these vessels, the vein may be tied and the whole incised beyond this point preparatory to their removal *en masse*. The ending of the thoracic duct on the left side may now be studied, or this may be done after the removal of the heart and lungs.

**MEDIASTINUM.**—The condition of the mediastinum is to be noted, especially as to emphysema in the areolar tissue, tumors, usually secondary, the ductus arteriosus, the thymus, and the peribronchial and other lymphatic glands. Except in the young the latter are pigmented, and for this reason have more than once been mistaken for melanotic sarcoma. They are often tuberculous, and may be infiltrated with cancerous or sarcomatous matter. Emphysema is most often produced after death by the removal of the sternum during the autopsy or by decomposition. When the lung is lacerated, the emphysema is more extensive and may even extend into the neck. Hemorrhage into the mediastinum may be due to trauma, to phosphorous poisoning, or to acute yellow atrophy of the liver. An abscess may be found, or a chronic mediastinitis, marked by fibrous thickening and density of the connective tissue. The latter usually occurs in conjunction with a fibrous pericarditis (mediastino-pericarditis), and is of importance on account of its influence upon the heart action (Orth).

**THYMUS GLAND.**—This weighs about 13.75 grammes at birth, and increases in size until the end of the second year, when it weighs about 26.2 grammes. It then gradually diminishes and after puberty is normally absent, though it has been observed in acromegaly, myxedema, exophthalmic goitre, and many other pathologic conditions. The gland should always be sought for, as it is not infrequently present in the adult, when it is more of a yellow color than the normal grayish red. Even when the thymic tissue itself has disappeared, the place of its former situation can usually be made out by the increased amount of fibro-adipose tissue. Hemorrhages are often found in the thymus glands of stillborn babes. Pus may sometimes be present.



Abscesses are seen at times in syphilitic children. Mistakes have been caused by the altered appearance of the normal juice after it has undergone post-mortem change; hence great care is necessary in making the diagnosis of suppuration. Sudden death in infants may be due to pressure symptoms from an unduly enlarged thymus. Sarcoma, endothelioma, angioma, and dermoids are found.

**THYROID AND PARATHYROID GLANDS.**—Both lobes of the thyroid may at this time be examined *in situ*, or, if preferred, the gland, together with the tongue, velum palati, epiglottis, oesophagus, trachea, parathyroids, carotids, intercarotid bodies, etc., may be removed in a single piece (see page 110) and studied subsequently detached from the body. The pyramid of the thyroid (Lalouette's pyramid) is a slight cone-shaped extension from the upper part of the gland to its point of attachment by loose fibrous tissue to the hyoid bone. The thyroid body may show enlargement due to parenchymatous or interstitial changes, or a combination of both, or associated with hypertrophy of the thymus and dilatation and hypertrophy of the heart. The colloid material may be considerably increased in amount and deposits of lime sometimes occur. Congenital goitre is now and then observed. The colloid goitre may become cystic and form cystic adenoma, into which hemorrhage might later occur. Thyroiditis is found in some of the infectious fevers, as diphtheria. Myxædematous degeneration, or cachexia strumipriva, is due to disease or removal of the gland. In cretinism the body is small, the head large, the countenance heavy, the abdomen protruding, kyphosis is often present, the lips are thick, the skin and mucous membranes dry and pale, and the hair is coarse and lustreless. Primary malignant tumors of the thyroid are seen, a mixed-celled sarcoma, at times angiomatous, being more common than cancer. There may be accessory thyroid glands, as at the base of the tongue, where goitre may occur.

The parathyroid glands,<sup>1</sup> which were discovered by Sandstroem, are four in number and are histologically different from the thyroid. They are usually unaffected by changes in the thyroid, colloid material being only rarely present. The superior parathyroids are situated behind the junction of the upper two-thirds with the lower one-third of the posterior thyroidal body and near the cricoid cartilages. The

---

<sup>1</sup> *El policlínico*, 1902, no. 21, fasc. 3; *Ziegler's Beiträge*, 1902, vol. xxxi, p. 142; *Virchow's Archiv*, vol. clxxiv, no. 3.

inferior group is posterior to the lower part of the thyroïdal lobes. The parathyroids probably develop from the third and fourth branchial clefts, those from the lower cleft eventually becoming the higher ones. They are of epithelial structure and furnish an internal secretion. They possess duct-like passages, probably analogous to the thyroglossal duct, and often become cystic. The parathyroids bear a distinct relation to the larger thyroid vessels and their shape varies considerably. As age advances the amount of fibrous tissue and fat increases. The vessels enter at a slight depression which may be regarded as a hilum. Tumors sometimes develop in the parathyroids and their removal may give rise to symptoms of myxœdema. Graves's disease is probably due to partial aparathyroidism, notwithstanding the fact that the disease is benefited by section of the cervical sympathetics.

**SUPERFICIAL EXAMINATION OF THE LUNGS.**—The appearance and situation of the presenting portions of the lungs are now observed. The normal color of the lungs at birth is a pinkish-white; in adult life, a dark slate-color, mottled in patches; as age advances this mottling may become nearly or quite black, owing to the deposit of carbonaceous material. Changes in color may be due to differences in the amount and character of the blood present or to some pathologic process. When the thorax is opened, the normal lung retracts, on account of its own elasticity. This contraction of the lung may not occur, because of the absence of elasticity, because of emphysema, because of pleural adhesions, because the alveoli are full of solids or fluids, the result of inflammation, or because stenosis of the larynx, trachea, etc., may prevent the egress of air. In cases of alcoholic intoxication and suffocation the lungs are generally found to be markedly expanded (Orth).

Next note the amount of fluid contained in the pleural cavity, whether or not it is clear, bloody, turbid, or of an inflammatory nature, and whether or not adhesions are present. The remarks made upon the peritoneal fluid apply with equal force to that found here and in the pericardium. As a practical point it is well to remember that serous membranes when normal are barely visible to the naked eye, being smooth and glistening, but when inflamed their appearance depends upon the nature of the inflammation; the membrane will then be found roughened and more or less opaque, especially if examined by an oblique light. The situation and extent of any lesion present should be noted.

If for any reason a pneumothorax be suspected, after carefully removing the fluid present, fill the pleural cavity with water and inflate the lungs with air by means of an intubation tube connected with a pump by a piece of rubber tubing. The rising air bubbles will reveal the situation of the laceration in the lung. In examining the pleural cavities inspect the left one first.

**PERICARDIUM.**—Note the position and condition of the pericardium, whether or not it is distended with fluid and to what extent it is covered by the lungs. When there is much distention of the pericardial sac, the direction and length of its principal diameters and borders—the latter, it should be remembered, are anatomically the reverse of those of the heart—should be noted before any fluid is allowed to escape. To open the pericardium it should be grasped near its centre by the fingers or a tenaculum, and a longitudinal fold elevated before it is incised in order to prevent injury to the heart and the escape of any excess of fluid which may be present. A small incision is then made at the highest point, under strict precautions if a bacteriologic examination is to be made, and the fluid present drawn off with a syringe or pipette into a graduated glass and its quantity noted. The opening in the pericardial sac may now be enlarged sufficiently to admit two fingers, which are then spread apart, thus elevating the pericardium and protecting the heart while the pathologist cuts between them. With a knife or a pair of scissors two incisions are made—one downward and to the right, the other downward and to the left—as far as the diaphragmatic attachment. The right flap is then drawn strongly forward away from the heart and another cut is made in an upward direction to the point where the pericardium is reflected around the vessels coming off from the heart (Fig. 71). The phrenic nerves are now plainly seen on the lateral inner surfaces of the pericardium and the anterior portion of the heart is exposed to view.

**Hydropericardium.**—Normally there are from one to two teaspoonfuls of clear citron-colored fluid in the pericardial cavity. In certain renal, cardiac, and pulmonary diseases this may be increased to several pints, the greatest amount being seen in general anasarca. The fluid is clear, watery, wine-colored, and may coagulate on standing. The serosa is smooth, glossy, and transparent. Later it may become sero-fibrinous, hemorrhagic, or purulent.

**Hæmopericardium.**—The presence of pure blood in the pericardiac sac is usually a sequence of rupture of an aneurism of the heart or of



the aorta, trauma, etc. Hemorrhagic exudates may be the result of inflammatory diseases, as in cancerous and tuberculous pericarditis, rheumatism, of chronic nephritis, of hypertrophy of the heart, etc. In the first instance the blood is present in large amounts and is clotted, while in the latter case it is derived from newly formed vessels in the inflammatory tissue.

*Pneumopericardium.*—The presence of air in the pericardial sac is nearly always caused by perforation from without, as in cases of stab-wounds, though it may be due to openings into the lungs, œsophagus, or stomach. Or it may be consequent upon decomposition, especially of an exudate. Its pathology resembles that of pneumothorax.

*Pericarditis.*—The pericardium, normally transparent and glistening, may lose its lustre, become rough and hyperæmic, and be covered with a more or less dry fibrinous exudate. When there is but little fluid and abundant exudate, the latter is thrown into villous projections by the movements of the heart, and the characteristic buttered surfaces, or *cor villosum*, may be found. Newly formed granulation tissue may follow a fibrinous exudate, with the formation of a productive pericarditis, and the later plastic adherence of the visceral layers of the pericardium, thus causing a complete obliteration of the sac. Suppurative pericarditis shows pus in the sac, and may be the result of trauma, or secondary to suppurative mediastinitis, cancer of the ribs, extension from pulmonary or pleural affections, or a general infection. Minute hemorrhages are seen, with flocculent or curdy collections in the dependent parts of the sac, and erosions are sometimes present. In some epidemics of pneumonia I have found that pericarditis was the immediate cause of death in nearly all the fatal cases.

*Other Lesions.*—Cancer, usually secondary, may be met with; in a specimen of melanotic sarcoma of the heart and pericardium at the laboratory of morbid anatomy in the University of Pennsylvania, the lesion was secondary here as well as in the lungs, the diagnosis of sarcoma being made by Prof. Guitéras from the discovery of pigmented cells in the sputum. Foreign bodies, gummata, cysticerci, echinococci, and trichinæ have been described. Tubercles may be seen along the course of the vessels or old cheesy tuberculous deposits may be found in chronic cases.

*Injuries.*—Wounds of the pericardium and heart may be caused by stabs, broken ribs, and foreign bodies in the œsophagus. If the main axis of the muscle fibres have been cut, the pericardium will be



full of blood; if the injury be parallel to its long axis, there may be no bleeding and the wound of the heart may heal spontaneously. Suturing of the heart muscle is now a well-recognized surgical procedure. Foreign bodies, like bullets, have been found encapsulated in heart muscle.

If an aneurism have been discovered, it is usually best not to separate the aorta from the heart, but to remove the aneurismal sac and the heart together. The aorta is not to be opened until the heart has been examined. In endocarditis vegetations are sometimes present in the arch of the aorta, and might easily be overlooked if not especially searched for. To discover air emboli, the thoracic cavity is filled with water and the heart opened under water *in situ*. Gas may arise from decomposition of the blood.

HEART.—The heart is to be observed before it is touched. Its normal position may be altered by fluid in the pericardium or in the pleuræ, by cardiac hypertrophy, in which case the apex may reach to the anterior axillary line, or by tumors of the mediastinum. The heart is about as large as the right fist. It measures from base to apex from 85 to 90 millimetres in men and 80 to 85 millimetres in women between the ages of twenty and sixty years; its greatest transverse diameter varies from 92 to 105 millimetres in men and 85 to 92 millimetres in women; it is about 35 or 36 millimetres thick in men, and from 30 to 35 millimetres in women. Any displacement is determined by the situation of the apex and the base, which are anatomically described especially in relation to the ribs, sternum, nipples, and median line of the body. Cardiac enlargement may be due to heart disease or secondary to disorders of the lungs, kidneys, aorta, etc. The color of the surface of the heart depends very much upon the condition of the epicardium and the underlying fat. The auricles, especially when well filled, are dark blue, while the color of the ventricles differs with the condition of the muscle. The consistence of the various portions of the heart depends upon the degree of contraction of its muscular tissue, as well as upon the amount and composition of its contents (Orth).

The contraction (systole) and the relaxation (diastole) of the two auricles and the two ventricles are considered in relation to the amount of blood contained within them. The amount of blood, especially if it be fluid, does not afford a criterion of the quantity therein during life, owing to the free communication of the vessels and cavities of the heart. After death from asphyxiation the right chambers of the

heart are distended with dark fluid blood, while after death from digitalis the left ventricle is contracted. Overfilling of the left ventricle is found when death was caused by cardiac paralysis. For bacteriologic examination or chemical analysis the blood is usually taken with the sterilized pipette, as described on page 347, from the cavity which is most distended by it, unless, of course, for some reason blood from a special cavity or side is desired. The circulation of the lymph and its deposits should be carefully studied.

The epicardium and the amount of subepicardial fat are to be carefully observed, as well as milk spots. In cachexia the subpericardial fat may be transformed into a soft, transparent, gelatinous mass, which becomes whitish on the addition of acetic acid. This is the so-called mucoid change of the subepicardial fat. Small lipomata may be found near the apex; and small subpericardial ecchymoses—so-called spots of Tardieu—are of medicolegal importance, as they are frequent in cases of death due to suffocation, particularly in the new-born, but may occur in the infectious fevers, as in diphtheria.

The situation and condition of the coronary arteries should be noted, and they, more especially the anterior one (the left), should be palpated, to ascertain whether or not they are "pipe-stem" in character. The interior is to be examined when they are opened later on. The coronary veins are easily distinguished from the arteries by the relative thinness of their walls as well as by their course. Overfilling of the larger veins indicates an obstruction to the outflow of blood from the right auricle (suffocation, etc.), unless it be confined to the posterior parts, in which case it is due to hypostasis. The large opening seen in the superior vena cava just before its entrance into the right auricle is the termination of the great azygos vein.

The interior of the heart is now to be examined, and here again, to secure the best results, it is expedient to adhere to a definite plan of procedure. There are several so-called "methods" of opening the heart, but all have the same object and all accomplish it more or less completely,—viz., that of exposing the cavities and valves with the least possible interference with the septa and the parts subsequently to be examined, and in such a way as to permit of the organs being reconstructed, or returned to their original shape and relations. The method adopted and described by Virchow for use in the Berlin Charité is undoubtedly the best, although the others may, if thoroughly understood and properly executed, yield very satisfactory results.

Ordinarily it is advisable that certain incisions be begun while the organ is still *in situ* and completed after it has been removed from the body. As each cavity is opened, careful note should be made of the quantity, color, and consistence of the contained blood and of the size and character of any clots that may be present. If the opening is occupied by a clot, this should be removed.

*Primary Incisions.*—After breaking up pericarditic adhesions, if present, the heart should be gently rotated on its long axis by slight pressure between the index-finger and thumb of the left hand, at the same time that slight traction is made downward and to the left of the body. This will bring the points of entrance of the superior and inferior venæ cavæ into view; midway between which the first incision is begun and then carried downward in the direction of the right ventricular ridge until the right auriculoventricular septum is reached (Fig. 72, *A B*, and Fig. 77). Next make an incision in the right ventricle, just below the auriculoventricular septum, passing down the right ventricular ridge to the interventricular septum, which is a little to the right of the apex (Fig. 72, *C D*). On the left side make an incision in the auricle, beginning in or slightly below the lowermost pulmonary vein and continuing in the direction of the left ventricular ridge as far as the auriculoventricular septum (*E F*). Open the left ventricle along the entire length of the left ventricular ridge, and, as this ventricle normally forms the apex of the heart, the incision will be carried to and through that point before the ventricular septum is reached (Fig. 72, *G H*, and Fig. 78). This incision must not join that of the other ventricle, but should be separated by an interval of about one-half inch. From the fact that these incisions are made while the heart is still *in situ*, they may be called primary incisions.

In cases of sudden death in which an embolus of the pulmonary artery is suspected, it is best to open that blood-vessel before removing the heart. This assures the finding of the embolus, which might otherwise be obscured in cutting the pulmonary artery for removal of the heart. By this method, also, the ductus arteriosus and congenital heart lesions in infants may be investigated.

*Removal of the Heart from the Body.*—To remove the heart, introduce the index-finger and thumb of the left hand into the left and right ventricles respectively, grasp the ventricular septum near the apex, and elevate the heart sufficiently to make slight traction on the great blood-vessels (Fig. 75). Then, if no aneurism be pres-






FIG. 72.—Primary incisions for opening the heart, usually made while this organ is still in the body.




FIG. 73.—Secondary incisions for opening the heart.




FIG. 74.—The completed incisions for opening the heart.



1

2

3

4

5

ent, sever all the normal attachments as near their point of passage through the pericardium as possible, and in the following order,—viz., the inferior vena cava, the superior vena cava, the pulmonary artery, the aorta, and lastly the pulmonary veins. Avoid injury to the œsophagus during the removal of the heart from the body. Or, the heart is drawn outward preparatory to severing the vessels, as may be seen illustrated in Fig. 76.

*Measuring and Testing the Valves.*—Immediately upon the removal of the heart from the body, the blood and clots should be carefully removed from about the valves. The valvular openings are then to be measured. Their size is usually estimated by the number of fingers that the ostium will admit. Normally the mitral ostium will admit the index and middle finger, whereas through the tricuspid opening the index and middle finger of one hand and the index-finger of the other hand can be introduced. This method is, of course, convenient, but is very unscientific and inaccurate and should be superseded by the use of a constant unit of measure. Graduated cones, or balls of definite sizes placed on rods (Figs. 49 and 50), answer the purpose very well. They are gently inserted in the direction of the blood-current, and the exact size of the opening can then be given in millimetres or inches. Vegetations upon the valves may be injured by careless handling. An equally scientific method is to measure the attached margins and to determine the diameter by dividing by 3.14 ( $\pi$ ).

The competency of the valves should now be tested. To do this, trim the great vessels down so that the valves may be seen. The heart is then evenly supported by each of the vessels in turn,—i.e., held in air and in such a way that the semilunar valves will be as nearly horizontal as possible, at the same time receiving no unnatural support from beneath. Water or mercury is then to be poured in by a second person until the vessel is filled, and note is made of the action of the valve. In case no one else is present, the heart is to be held under water and then quickly taken out, and the valve being tested observed. If there is any leakage from the aorta, make sure that it is not from a cut branch of one of the coronary arteries. The best result of the water-test is seen in the semilunar valves, the competency of the auriculoventricular valves not being accurately determined by this method, which has of late rather fallen into disuse.

Should it be necessary or desirable to ascertain the competency of the auriculoventricular valves, the primary incisions above described

are not made until the heart has been removed from the body, and the test is begun by cutting a transverse slice from the apex and exposing the ventricles. The heart is now everted and each ventricle is filled separately with liquid. This method of removing the organ before opening is also useful in examining the heart of a child or when it is desired to make a bacteriologic examination of the valves. In the latter event no water should previously be used, lest some of the vegetations be washed out or other bacteria than those present be introduced, thus creating more or less serious confusion.

Hamilton advises the use of air for testing the competency of the valves, and gives the method as follows:<sup>1</sup> "An incision is first made into the left auricle, and any post-mortem clots are carefully removed from the left chambers through it. Another incision large enough to admit the nozzle of a half-inch tube is made into the ventricle near its apex and in the line of that required for laying it fully open. The tube is joined to a bellows, and air is driven intermittently into the ventricle by means of it, the aorta having been meanwhile closed. The valve will be seen to open and close, according as the air is aspirated or driven out of the bellows. A like procedure is adopted for the demonstration of the tricuspid. To test the aortic valve, the incision before described as necessary to lay open the left ventricle is continued up as close to the valve as possible without injuring it. The tube is tied into the aorta, and the action of the valve is watched from below. The same method is used to test the competency of the pulmonary artery valve. As a matter of fact the tricuspid, in the human heart, will always be found more or less incompetent."

*Secondary Incisions.*—Place or hold the heart with its posterior surface downward. This can be told by the situation of the pulmonary artery, which is situated anteriorly. Insert a pair of probe-pointed scissors or the blade of the enterotome (now a cardiotome) in the incision in the right ventricle, and cut from the centre of that incision through the centre of the attachment of the two anterior leaflets of the pulmonary artery (Fig. 73, *I J*, and Fig. 79). The point of junction of the anterior leaflets can usually be seen from the outside, but, if not, it can very easily be determined by looking into the vessel or feeling it with the index-finger. This incision is to be continued until it opens up the entire portion of the pulmonary artery which has been removed

---

<sup>1</sup> HAMILTON, *Text-book of Pathology*, vol. i, p. 9.



FIG. 76.—The pulmonary veins are placed on a stretch and are ready to be incised. There are two veins on the right and four on the left side, six in all.





FIG. 77.—Method of opening the right auricle; an incision is made down to the auriculoventricular septum of the right side. This incision is usually made while the heart is *in situ*, but for the sake of clearness is here shown as being made outside of the body.



FIG. 78.—Method of opening left ventricle. The heart is being opened outside of the body. The left hand steadies the heart while the knife cuts along the left ventricular ridge, starting just below the auriculoventricular septum and ending at the apex.



FIG. 79.—The pulmonary artery is made tense with the left hand, while from the centre of the right ventricular incision the anterior portion of the right ventricle is cut in the direction of the thumb and middle finger which mark the junction of the two anterior pulmonary semilunar cusps.



FIG. 80.—The left auricle and ventricle are fully opened, exposing the mitral valve, chordæ tendineæ, papillary muscles, endocardium, etc.



FIG. 81.—Completed incisions of the heart, the organ having been reconstructed after the examination of all its cavities and parts. The coronary artery has not been dissected out. This is done for several inches with the scissors, and then transverse incisions may be made with the knife about three-eighths of an inch apart as the artery becomes smaller and branches.



from the body. Some pathologists advise making this incision towards the left of the pulmonary artery, so as to cut between the left anterior and posterior cusps. The right ventricle is now exposed so that the condition of the pulmonary valves, endocardium, myocardium, chordæ tendineæ, etc., of this side of the heart may be noted. Now dissect away the connective tissue binding together the pulmonary artery and the aorta.

In opening the left ventricle, cut the anterior wall as near the ventricular septum as possible, starting from the apical extremity (*H*) and stopping at the point overlapped by the left auricular appendix (*K*). Then, using the cardiotome, the incision is completed (either from the aorta or from the ventricle) by cutting between two leaflets (*L K*). In the aorta there is but one anterior leaflet, consequently the incision should be to either one or the other side, but preferably as close as possible to the curves of the pulmonary artery. After examining the valves, myocardium, aortic intima, etc., dissect out the coronary arteries with probe-pointed scissors.

Lastly, unite the auricular and the ventricular incisions of each side by cutting through the auriculoventricular septa (Fig. 74). In Fig. 80 is shown how well the auricle and ventricle of the left side may be examined after the completion of this incision. The knife is best introduced from the auricle to the ventricle with the edge of the blade down and then turned, the cutting being done from within outward. It will be noticed that the instruments are passed through the valvular openings in the same direction as the blood flows. The valves will not be injured by this method and the entire heart can be folded together so as to show its original contour (Fig. 81). In extreme mitral stenosis it is often advisable not to complete the left auriculoventricular incision.

A simple method of opening the heart, and one which yields fair results, is to place two fingers on the anterior ventricular septum, which is recognized by the situation of the anterior coronary artery, and make two parallel incisions into each ventricle. The pulmonary artery and the aorta may then be opened.

The heart, freed from blood and clots, is now to be weighed.

The gross appearance of the heart, as well as the thickness, color, and consistence of the various parts of the cardiac muscle, can now be observed. The wall of the right ventricle is normally from 2 to 3 millimetres thick (in women slightly less than in men) and may



pathologically measure from 7 to 10 millimetres. The thickness of the wall of the left ventricle is from 7 to 10 millimetres, and may be increased to 25 millimetres or more by pathologic changes. The estimation of the weight of the heart is one of the means of determining whether or not a true hypertrophy is present. The normal heart weighs about 250 grammes in women and about 300 grammes in men; but when hypertrophied it may weigh over a kilogramme. In the puerperal state the heart is normally increased in size, the right side often dilated (Letulle), the subpericardial fat increased, and hemorrhages may occur.

The color of the heart muscle varies according to the amount of blood it contains, but is always lighter and more grayish red than the skeletal muscles. It may be of a brownish red or even brown, as in anæmia and brown atrophy of the heart. In the latter condition the tortuous vessels and mucoid covering form a striking picture. The scattered yellowish patches seen throughout the muscle appear in bands, making a sort of net-work (wren's breast or tiger markings). This yellow streaking is often most conspicuous on the papillary muscles of the left ventricle. When this condition is at all extreme, the endocardium and pericardium will be found greatly thickened.<sup>1</sup> In septic conditions the heart is of a dirty-red color and very friable. Light-gray spots or streaks indicate the formation of fibrous tissue. The consistence of the heart muscle varies with the color: brown hearts are hard and dense, while those of a yellowish tinge are apt to be soft and flabby. The fibroid heart is always hypertrophied. After dilatation of an hypertrophied heart sets in, the muscle becomes softer by the process of fatty degeneration. The heart muscle is very soft in sepsis, and in cases of heart weakness developing after infectious diseases, especially after typhoid fever and diphtheria.

Hypertrophy and dilatation are usually associated with each other. In concentric hypertrophy the walls are thickened and the cavities are smaller than normal. As this condition is often due to post-mortem contraction or to marked systole, the heart should be soaked in tepid water before the measurements are taken. One may also distinguish simple hypertrophy, where overgrowth of the walls is found associated with normal cavities; eccentric hypertrophy, or hypertrophy with dilatation; and pure dilatation without hypertrophy. The highest

---

<sup>1</sup> E. BEER, *Jour. of Path. and Bact.*, December, 1903.

degrees of hypertrophy occur in cases of double aortic disease, where, too, moderator bands are sometimes found. The conical shape is often lost by the broadening of the apex and deepening of the muscles.

In the examination of the auricles an aperture in the foramen ovale may be overlooked if the heart is so held as to put the auricular wall too much on the stretch. As the communication between the auricles usually takes place by openings most frequently coming off from beneath the former edges of the valve, all suspicious cracks, orifices, or slit-like communications should be searched for with a pointed probe while the heart is relaxed, care being taken not to tear or puncture the tissue or to mistake the ending of the coronary veins, which empty near the obliterated opening, for a patulous foramen. In the left auricle the pulmonary veins rarely come out intact; should they do so, a V-shaped incision is made between each pair so as to expose them. The "dog's ears" are opened by cutting towards their tips, with an extra incision transverse to this should it be required. Softening clots may then be discovered which otherwise would escape attention.

If it be desired to follow out the subclavian vessels by careful dissection, the entire clavicle of that side should first be removed.

If a sound be used for finding the opening of the thoracic duct into the vein, care must be taken not to injure the valve which is present at this point. It is much more difficult to find the entrance of the lymphatic vein of the right side at the junction of the jugular and the right subclavian veins, as the parts are correspondingly smaller on this side of the body.

The situation of the mitral and pulmonary valves can be easily remembered by the mnemonic *Martin Luther, The Reformer*,—*mitral* on the *left* side, *tricuspid* on the *right*. That there is but one *posterior* cusp to the *pulmonary* valve and one *anterior* cusp to the *aorta* affords an easy way to recall this oft-forgotten point.

**REMOVAL OF THE LUNGS.**—To remove the lung the left hand, palm inward, is introduced along the costal curve until the under portion of the upper lobe can be elevated without undue pressure being made upon the pulmonary tissue. Should there have been no antecedent inflammation and consequent adhesions, a condition especially liable to be found at the apices, this procedure is readily accomplished, but sometimes, when the adhesions are very strong and cannot be broken down by the hand, a probe may be used for this purpose, or it may be necessary to dissect away the costal pleura and even the ribs and

remove them along with the lung. When this condition is found in the performance of routine postmortems, the examination of the affected lung may be accomplished by making the incisions while the organ is still in the body. The upper lobe is now carried away from the median line of the body, anteriorly and downward, thus exposing the structures forming the root and giving a fine picture of the arch of the aorta. Then, separating the index and middle fingers of the left hand, the root of the lung is surrounded so that the upper lobe rests on the palm. In this way pressure can be made downward and away from the spinal column. Next a perpendicular incision should be made in the direction of the spinal column and the bronchus severed. The advantage of this procedure is that it enables the operator to observe the character of the fluid in the bronchus, avoiding its (otherwise very probable) contamination with blood. When the character of the fluid is noted, the rest of the structures, including the intercostals arising from the aorta, may be severed with a few horizontal incisions, care being taken to avoid cutting the aorta, the œsophagus, the large azygos vein, and the thoracic duct. It is well to remember when cutting these vessels that the left bronchus, which is considerably longer and smaller in diameter than the right, is situated *below* the left pulmonary artery, while the right undivided bronchus is entirely *above* the right pulmonary artery. The left lung should be removed first, and, as it has usually two lobes while the right has three (I have seen this condition reversed but twice), there is no necessity of adopting any method of distinguishing them after they have been removed from the body. Then, too, the left lung has a depression in its anterior border for the apex of the heart, it is longer and narrower than the right, not quite so heavy, and, as already stated, the arrangement of the bronchus and artery is different on the two sides. If, however, it is deemed necessary to do this, a single cut in the apex or bronchus of the left lung and two in the right will afford a ready means of distinguishing the one from the other. Normally, the lungs are darkened from inhalation of pigmented material, the deposit of pigment often assuming a mosaic or net-work appearance, corresponding to the situation of the lymphatics in the external lobules of the lung. Examine the visceral pleura for fibrinous deposits, exudates, adhesions, etc.; note the color, which varies with the age, the quantity of contained blood and air, minute hemorrhages, excessive pigmentation, cicatrices, spicules of bone, emphysematous spots, miliary tubercles, calcified tubercles with cheesy



FIG. 82.—Method of opening the left lung. The organ, lying on its posterior surface, is held steady by slight pressure with the left hand on its upper portion, while a long, clean cut is made from the apex to the base of the lower lobe. In opening the right lung the incision is best made in the opposite direction, —*i.e.*, from the base to the apex.





Fig. 1. Lung from this case was emphysematous and contained a large cavity. It was found to contain in the lung in its examination.



FIG. 84.—Method of opening the branches of the pulmonary vein. The artery is thicker and more elastic than the vein. The veins are best opened after the primary incisions shown in Fig. 83 have been made. It is naturally impossible in the same lung to make a complete dissection of the bronchi, artery, and veins, owing to cutting of vessels not belonging to the system undergoing dissection.



FIG. 85.—Method of opening the bronchi and their ramifications.



interiors, nodules, patches of consolidation, hemorrhagic and anæmic infarcts, tumors, infectious granulomata, etc. The lungs should be weighed at this time, before they are opened for further study.

Each lung is then carefully and lightly palpated from above downward throughout its entire extent by running the fingers over its surface, the fissures being separated and the anterior and posterior edges examined. By gentle pressure between the fingers crepitation is now produced. What this is like in the normal lung can be learned only by actual trial. In marked emphysema the crackling sound of the larger blebs as they break can sometimes be heard across the room. The presence of liquid naturally decreases crepitation. In hepatization the pulmonary parenchyma may break down even under gentle pressure. After squeezing a crepitant portion of the lung, there is ordinarily enough air left in the tissues to cause the pieces to float upon water. An interesting experiment in cases of atelectasis, infarcts, etc., portions for microscopic study having been previously removed, consists in blowing air forcibly through the bronchus by means of a cannula connected with a bellows.

Placing the lungs upon their posterior surface on a board, rather than upon the more slippery stone table, the lower lobe of one lung is grasped with the thumb, the remaining fingers seizing the upper lobe;<sup>1</sup> in this way the organ may be firmly held (Fig. 82). With a single stroke an incision should be made from apex to base, commencing at the lateral convexity and passing to the entrance of the large vessels in the direction of the bronchi, the lung being now laid open like a book (Fig. 83). In the case of the left lung the base had better be turned towards the operator, while in the right it will be *vice versa*, requiring an extra incision to open the middle lobe. *Immediately note the color of the cut surface.* The normal color without blood is light gray, while with different quantities and qualities of blood the shade ranges from light red or brick-red to dark, black, or blue-red. In heart disease the color of the pulmonary tissue is apt to be brown; in anthracosis it is black. The amount of hypostatic congestion, and the character of the fluid which exudes on lightly squeezing with the fingers areas not intended for microscopic study, are now determined. A microscopic examination of the scrapings collected by passing the

<sup>1</sup> If the index-finger be introduced into the fissure between the lobes (and this method holds the lung very securely), care must be taken not to cut the finger in the subsequent procedures.



knife-blade over the cut surface should be made. The appearance of the surface after removal of the liquid is determined, and any unusual spots more carefully examined. Next it is necessary to examine the substance of the lung for cavities, to observe the shape and position of areas of consolidation, and to ascertain the specific gravity of consolidated areas in cold water. In pneumonic cases the entire lung may be placed in water to determine the portion containing air. A hemorrhagic infarct or a portion of an apoplectic lung will sink in water, as well as the lung of croupous pneumonia. Cubes of normal lung may sink in fluids having a low specific gravity.

**EXAMINATION OF THE PULMONARY VESSELS.**—Now is the time to open the pulmonary veins (Fig. 84), artery, and bronchi (Fig. 85). Parallel or transverse incisions may be made, but care should be taken not to make them so deep as to detach any portions of the lung. The pulmonary arteries resemble the veins in character, though they are thicker, more elastic, and whiter than the latter. By following the pulmonary artery up on its anterior aspect from the heart there is no danger of mistaking one for the other, this error most often occurring when the dissection is not started until after the lungs have been removed from the body. Again, there is but one artery for each lung, while there are two veins.

**REMOVAL IN ONE PIECE AND SUBSEQUENT EXAMINATION OF TONGUE, ŒSOPHAGUS, TRACHEA, AND ADJACENT STRUCTURES.**—It is frequently advisable to excise as one piece the tongue, œsophagus, thyroid gland, trachea, epiglottis, etc., so that a minute examination of these parts may be made while they are exposed to good light in a convenient situation. For this purpose, in those cases where disfigurement of the body is of no importance, the primary incision over the thorax may be extended up to the symphysis mentis and the parts dissected out with ease. Orth's method of doing this is as follows: The skin is first reflected. Then by the use of the cartilage-knife an incision is made into the mouth at one angle of the jaws as close as possible to the bone, cutting with a sawing motion to the chin and then back on the other side to the angle of the jaw, severing the geniohyoglossus muscle. The tongue, after being separated from the jaw, is pulled down with the forceps held in the left hand, after which the soft palate should be separated from the hard by the use of a knife, including in the operation the tonsils. A cut should now be made as high up as possible to remove the pharynx, trachea, and œsophagus



FIG. 86.—Method of removing tongue, tonsils, œsophagus, trachea, etc., in a single piece, without incising the skin more than is done in the primary cut.




FIG. 87.—Examination of the organs of the neck. The arrows show the direction in which the incisions in the tongue and in the posterior wall of the esophagus are to be made. (After Nauwerck.)



FIG. 88.—Method of opening trachea posteriorly. The incision starts from above and extends downward.



FIG. 89.—Examination of trachea and vocal cords. The incision may be made anteriorly, thus leaving the walls between the two parts intact.





FIG. 91.—Method of tying the intestines preparatory to their removal. They are tied in two places, a foot or so above the ileocecal valve.



FIG. 92.—Bucket method of opening and cleansing intestines, especially useful in private cases.

from the spinal column and the deep pharyngeal muscles. This should be done with small perpendicular incisions on the spinal column through the retropharyngeal and retro-œsophageal tissue, the tongue being pulled strongly forward. The parts may now be left intact for later dissection, or the œsophagus may be cut off just above its entrance into the stomach and the trachea below the vocal cords. These parts may be removed in a single piece, however, without the incision being extended to the chin, as by careful manipulation the hand can tear the skin away anteriorly from its attachments by working from beneath, and a knife may be introduced from below into the centre of the tongue (through the geniohyoglossus muscle) posterior to its frænum, thus leaving the tip *in situ* in case an examination of the mouth is to be made. (Fig. 86.) By a circular incision of the muscles, fasciæ, etc., with the knife beneath the skin, keeping as close as possible to the bony walls of the jaw, to the carotids, and to the bodies of the vertebræ, the pharynx, larynx, trachea, and œsophagus may be separated, drawn forward and downward, and removed. The tonsils are either torn out bodily with the fingers from below, or else incised while in the body and examined from above or below by reflected light. The velum palati may also be removed by cutting the hard palate in front and dissecting it away from its bony attachment. This should always be done in cleft palate. The edges of the tongue may be examined for injuries, such as wounds made by the teeth during a fatal convulsion. The mucous membrane is flattened in syphilis, and the tongue may be the seat of lymphangioma. The vessels of the arm should now be cut and the arch of the aorta detached from the œsophagus and bronchi. The whole aorta may be examined later, if it is deemed better, or this vessel as far as the diaphragm can be removed with the œsophagus. This procedure is particularly useful when disease of the latter is suspected, as a cancer of the gullet may rupture into the aorta or an aneurism of the aorta break into the œsophagus. The aorta is from seven to eight centimetres in diameter at its commencement and gradually narrows to forty-five or even thirty-five millimetres in the abdominal cavity. The œsophagus and the trachea are preferably opened up posteriorly throughout their entire extent. (Figs. 87, 88, and 89.) Carefully examine the vocal cords; see if there are any tumors, syphilitic or tuberculous ulcerations, inflammation, malformations, foreign bodies, diphtheritic membrane, etc. The condition of the mucous membrane of the trachea and



the elasticity of its cartilages should be carefully tested. I have seen the whole transformed into a rigid tube by infiltration with lime salts. The trachea and larynx may show abrasions and injuries, as by cut throat, deformities, stenosis, pressure defects, etc. The œsophageal veins frequently carry on a large part of the collateral circulation in cirrhosis of the liver, and the rupture of one of them may cause death from hemorrhage, much blood being found in the intestinal tract. The œsophagus may show peptic or typhoid ulcers, and diverticula of various kinds and degrees are seen. Ulceration from cavities, as well as those just mentioned, may on healing result in stricture, which as time goes on may be the starting-point of cancer.

In cases of strangulation, as by hanging or otherwise, the examination of the vessels of the neck is of great importance. For this purpose the incision behind the ear, made for the removal of the brain, may be extended down the neck, and the skin, fat, and superficial fascia of the face dissected away, thus making easy the exposure of the jugulars, carotids, etc. The tearing of the intima of the carotid indicates hanging or strangulation; marks produced by pressure of the rope, in the form of parchment-like skin at the sides of the neck and hemorrhages into the tissues, are also found after death by hanging. Emboli of the carotid may cause sudden death, and thrombophlebitis of the jugular in cases of thrombosis of the lateral sinus is to be searched for. Aneurisms are sometimes seen. Hemorrhages into the sympathetic nerves may occur in cases of fever with delirium and of heat-stroke; pigmentation and fatty changes also take place in the cachexias and in fevers.

The finding of the carotid body is facilitated by the removal of the common carotid artery along with about one-half inch of the internal and external carotids. The ganglion is located usually at the angle formed by the bifurcation of the carotid, but it may be situated at any place in this neighborhood. It closely resembles the superior cervical ganglion and has the size of a grain of rice, is somewhat oval in shape, vascular, and of good consistency. It is attached to the artery by means of a small band of connective tissue; bands from the capsule divide it into nearly equal parts and again into lobules. Funke<sup>1</sup> has tabulated fifteen tumors arising from the carotid body, his own case being a perithelioma.

---

<sup>1</sup> *Amer. Med.*, July 16, 1904.

## CHAPTER VIII

### DISEASES OF THE HEART, BLOOD, BLOOD-VESSELS, AND LYMPH-VESSELS

ANOMALIES.—Abnormalities in the development of the heart are frequent and varied, occurring especially in non-viable infants. Thus, one may find acardia, double hearts, two- or three-chambered hearts, rudimentary hearts, malformed blood-vessels, premature closure or patency of fetal passages, as of the foramen ovale and ductus arteriosus, imperfections of the septa, increase or decrease in the number of semilunar leaflets, stenosis and atresia of the pulmonary, aortic, and arteriosal conuses, transposition of the primary arterial trunks, ectopia, with deformity of the anterior part of the chest, etc. Dextrocardia may be part of a general congenital *situs inversus* or be confined to the thoracic organs alone. Acquired malformations are considered under the separate diseases of the heart.

BLOOD.—Many of the changes which the blood undergoes are macroscopic and can be studied *post mortem*, though it should be remembered that numerous factors tend to alter its composition and color in a dead body. A pocket spectroscope and a Tallqvist or a Wetherill (Plate III) hæmoglobin scale are useful for studying the characteristics of the blood. Reproductions of the spectra of hæmoglobin, reduced hæmoglobin, carbon-monoxid hæmoglobin, etc., will be found in many of the works on physiology and toxicology. In using the Tallqvist's blood-color scale due allowance must be made for the difference in the color of the blood after death and for the abstraction of the water. At my request, Dr. Wetherill has kindly prepared a post-mortem color scale for showing the percentage of hæmoglobin in the blood twenty-four hours after death.<sup>1</sup>

The method of using the scale, which for the sake of convenience has been placed upon the inner back cover (Plate III), is quite simple.

---

<sup>1</sup> The writer is at present engaged upon the preparation of a scale designed to show the changes of color in human blood twenty-four hours after death resulting from the more common poisons. He will be thankful to receive from the reader reproductions showing any such alterations except those induced by hydrocyanic acid and illuminating gas.



A drop of blood taken from one of the large veins, such as an innominate, or from the right side of the heart, is placed upon or absorbed by a small piece of filter-paper or white blotting-paper and held in the centre of the disk, so as to compare the color with the scale in daylight at an angle (by preference) of  $45^{\circ}$ . As soon as the colors are nearly matched, the paper is run out along the black dividing line, thus affording an opportunity of forming an intermediary judgment, which can be approximately determined to within 2.5 per cent. The black lines separating the colors and the outer white paper best bring out by contrast the various shades of color.

While printing this scale, the opportunity was afforded of supplying a copy of Dr. Wetherill's moisture scale, which is naturally of more use clinically than at the postmortem. The change in color depends upon the action of moisture on filter-paper soaked in a four-tenths of one per cent. of a saturated solution of the chlorid of cobalt. To measure the moisture in a room or upon the surface of a dead or living body, the disks are dried over a burning match or in a desiccator and placed under celluloid or glass to exclude the atmospheric conditions, if the latter procedures be desired. After ten minutes a comparison is made with the scale and the percentage determined. In uræmia and diabetes the skin is dryer, as well as upon the side affected by a recent paralytic stroke. The scale may be used for determining the amount of moisture in the air at the place where the body is found or the autopsy performed.

The specific gravity of the blood is best obtained with the specific-gravity bottle, there being no difficulty here, as during life, in securing the desired quantity of blood. Hammerschlag's method of mixing benzol and chloroform of a known specific gravity until a drop of blood remains stationary may also be employed. The specific gravity of the blood in health is normally from 1057 to 1059, and varies directly as the amount of the hæmoglobin present. In leukæmia the specific gravity is high, while after drowning it differs according as the person has been drowned in salt or fresh water. The cryoscopic index is  $-0.57^{\circ}$  C., showing practically no variation during health.<sup>1</sup> For its use in drowning, see Chapter XXVI.

---

<sup>1</sup> For a recent article on cryoscopy, see CATTELL, *International Clinics*, April, 1904. Revenstorff (*Jahrschr. f. gericht. Med.*, vol. xxii, 1903) determines by this method the length of time which has elapsed since death.

There may be observed at the time of the postmortem all degrees of coagulation of the blood, from an almost absolutely fluid condition, as in poisoning by hydrocyanic acid, to a hard and dense fibrinous clot,—the so-called heart polyp,—which contains almost no red blood-corpuscles. The firm, yellowish “chicken-fat” clots, seen so frequently in pneumonia, adhere to the walls of the heart, and indicate slow death, with gradual paralysis of the heart’s action. When all the coagula are rich in fibrin, some acute inflammatory process has caused an increase in the leucocytes and blood-plaques, the generators of fibrin. In the left auricle they at times assume polypoid or spherical shapes, and may even become attached to its wall and undergo organization. The ordinary post-mortem coagulum is the red clot, the so-called currant-jelly clot, which is not attached to the endocardium, though it may adhere to the muscular interstices of the heart. *Hyperinosis*, or increased capability for fibrin formation in the blood, is at times met with in certain anæmic affections and infectious diseases. *Hypinosis*, or decreased capability for fibrin formation, occurs in leukæmia, hydræmia, certain of the acute exanthemata, hemorrhagic diathesis, obstruction of the biliary tract, and in cases of suffocation or intoxication with certain poisons, as carbonic acid. Methæmoglobin, found in cases of poisoning by chlorates, nitrites, toadstools, etc., gives a brownish tinge to the blood. In putrefaction, if the blood be left standing, the clear serum separates and leaves a yellowish-green sediment. Under the microscope shadows of red cells are seen. The blood is normally alkaline, but at the end in Asiatic cholera it may be markedly acid or its alkalinity much diminished.

The following diseases may be diagnosed by their agglutinative reaction: Typhoid fever, paratyphoid, dysentery, Malta fever, cholera, plague(?), tetanus (in the horse, but not in man), psittacosis, tuberculosis(?), and pneumonia. The reaction with the proteus may be used to distinguish between invasion after death and infection during life. Auto-agglutination of the erythrocytes has been observed in Hanot’s hypertrophic cirrhosis. Flexner believes that the liability of the red corpuscles to become agglutinated in the blood-serum is the cause of thrombi which so frequently occur in certain of the infectious fevers, as typhoid.

*Pathologic Conditions.*—(a) *Plethora Vera*.—A condition in which all the elements of the blood are proportionately increased. (b) *Plethora Serosa*.—A marked increase in the watery constituents, ap-

pearing after transfusion, increased ingestion of liquid, acute cardiac failure, etc. (c) *Olygæmia*.—A diminished amount of blood; occurs only as a temporary condition. (d) *Hydræmia*.—Abnormal increase in the watery portion of the blood; seen in cardiac, pulmonary, hepatic, and renal diseases. In hydræmic plethora there is an absolute increase of serum. If relative, it is called *oligocythæmia*. (e) *Anhydræmia*.—Here there is a concentration of the cellular elements of the blood due to an abstraction of its watery constituents, the blood becoming thick and even tarry, as in cholera. It is seen at times after tapping, as for ascites, and in starvation. (f) *Hæmolysis*.—Destruction of red corpuscles; occurs after burns, certain poisons, infectious fevers, etc. (g) *Lymphocytosis*.—An absolute and relative increase of lymphocytes; commonly associated with hereditary syphilis, scurvy, chlorosis, pernicious anæmia, Graves's disease, splenic tumors, pertussis, pneumonia, and lymphatic leukæmia. (h) *Eosinophilia*.—An increase in the number of eosinophiles; occurs in asthma, fibrinous bronchitis, acute and chronic skin diseases, especially in trichinosis, ankylostomiasis, also after acute infections and with malignant tumors (moderate increase). It is a compensatory reaction with diseases of the spleen. Eosinophiles are decreased after castration, and in the febrile stages of pneumonia, grippe, typhoid, diphtheria, and sepsis. (i) *Myelocytosis*.—An increase in the number of myelocytes is always pathologic, and is seen in greatest degree in myelogenous leukæmia, pernicious anæmia, acute infections, as typhoid, mania, Basedow's disease, syphilis, tuberculosis, osteomalacia, etc. It may occur in all diseases with marked anæmia. (j) *Polycythæmia rubra* is an absolute increase in the red cells; seen in cured cases of anæmia,—usually associated with engorged organs,—in new-born children, and in persons living at high altitudes or suffering from chronic phosphorus and carbon monoxide poisoning. (k) *Anæmia*.—A diminution in one or more of the constituents of the blood. (l) *Primary, Essential, or Idiopathic Anæmia*.—An anæmia, the cause not definitely known, usually attributed to the blood-making organs, and characterized by a *disproportionate* reduction in the elements of the blood. (m) *Secondary, Simple, or Symptomatic Anæmia*.—An anæmia due to a definite cause, as an *Anchylostomum duodenale*, *Anguillula stercoralis*, *Bothriocephalus latus*, *Ascaris lumbricoides*, *Tricocephalus dispar*, trichinosis, infectious fevers, etc., and characterized by a *proportionate* reduction in the elements of the blood; shown by changes in the specific gravity.



color, size, and shape of red cells, etc. (*n*) *Poikilocytosis*.—Alteration in the shape of red corpuscles (crenated, reniform, and pyriform are most common). (*o*) *Macrocytosis and Microcytosis*.—The red cells are respectively increased and diminished in size. (*p*) *Leucocytosis*.—Increase in the number of white blood-cells without alteration of the relative numbers of each variety; marked in the new-born, in pregnant and parturient women; usual at postmortem; appears with many infections, suppuration, malignant disease, and hemorrhage. (*q*) *Leucopenia*.—A diminution in number of white blood-cells; seen most characteristically in typhoid fever, starvation, cancer, grippe, measles, tuberculosis, typhoid fever, Hodgkin's disease, and phthisis; normal in pregnancy, obesity, alcoholism, nephritis, icterus, typhus, malaria, and cardiac and pancreatic diseases. (*r*) *Lipæmia*.—Fat in the blood, giving it a milky appearance; seen in leukæmia, diabetes, alcoholism, phthisis, etc. (*s*) *Uræmia*.—The presence in the blood of an excess of chemical compounds, as urea, which should be eliminated by the kidneys or other excretory organs. There may have been during life an increase in blood-plaques, but they are most difficult of demonstration even shortly after death. In uræmia, besides the macroscopic lesions of œdema of the brain, a condition of chromatolysis of the cells in the central nervous system may be demonstrated. The destructive changes are especially found in the motor cells, and may be followed later on by degeneration of the motor tracts.<sup>1</sup>

*Blood-Stains*.—When any suspicion of violence occurs, look carefully for blood-stains. If in doubt, treat all suspicious findings as if they were such, unless some special reason exists for not doing so. *Such stains should be most critically examined in the privacy of the laboratory* before expressing an opinion as to whether or not they are consistent with human blood. Try to ascertain: (1) Their connection with the body under examination or with the person suspected of the crime. (2) Their extent, using great care in determining the nature of the substance stained. (3) Conditions,—whether fluid or clotted, wet or dry, cracked or caked, etc. (4) How made,—whether by smear, by splash, by flow, by soaking up, as in cloths, etc. (5) Connect, if possible, the amount, shape, and condition of the stains with their probable source, and note any peculiarities. When practicable, preserve parts or all of stains. It is often well to saw off an entire

<sup>1</sup> EDITORIAL, *Jr. Amer. Med. Assoc.*, April 23, 1904.



step or remove a panel of a door in order to produce the same as evidence in court. In the present state of our knowledge it is not practicable to state from what part of the body the blood came and the age of the stain, though the more recent the blood the more soluble it is.

Two illustrations from my case-book will show the importance of this line of research. A man committed rape on a seven-year-old child and murdered her. Blood was seen on the fly of his trousers by his room-mate. In order to divert suspicion from himself, the murderer accused his room-mate of the crime. The trousers of both men were sent to me for examination. In the pair of pants belonging to the perpetrator of the crime the lining of the fly had been cut away and neatly sewed, but there remained a few telltale threads containing blood, which was found to possess the characteristics of human blood. On the trousers of the other man was found a red substance, which examination showed to be lumberman's red chalk, the crime having been perpetrated in the backwoods. In the second case blood splashes on a white curtain were stated by a murderer to be red paint which one of his children had put there with a paint-brush.

The presence or absence of blood is determined by (1) the physical examination; (2) chemical tests; (3) spectroscopic examination; (4) microscopic examination, and (5) the hæmolytic serum test.

The agglutinative reaction or antiserum test for the diagnosis of human blood has been applied in a number of recent trials. It would seem to afford positive proof of the special source of the blood under examination, though Robin secured the reaction from the blood of a monkey and Linossier, Nuttall, Dieudonne, and others have demonstrated the reaction in pus, nasal mucus, saliva, urine, pleural exudate, and sweat derived from the human body. Uhlenhuth<sup>1</sup> was put to a severe test by the German Department of Justice. Various objects stained with the blood of man and of different animals were sent to him, the nature of the blood being known to the Department of Justice but not to him. When the blood was furnished in sufficient quantities, his results in each case were positive.<sup>2</sup> One method of preparing the

---

<sup>1</sup> *Deutsche med. Wchnschr.*, September 11 and 18, 1902.

<sup>2</sup> For an account of trial cases see PATEK and BENNETT, *Amer. Med.*, September 6, 1902, p. 374; WHITTIER, *Amer. Med.*, January 18, 1902; FERRAR, *Bolletino della reale Accad. medica di Genova*, 1901, no. 7; OGIER and STICKIS, *Soc. de méd. légale*. P. and BECHTEL case tried at Allentown, Pa., in 1904.

antiserum and applying the test is as follows: Ten cubic centimetres of defibrinated human blood, as that freshly obtained from the human placenta, are injected into the peritoneal cavity of a rabbit at intervals of six days, and after five such injections an effective serum should be obtained. Butza<sup>1</sup> prepares the animal by injecting from ten to twenty cubic centimetres of a centrifugated human pleural exudate intraperitoneally into a rabbit for five or six successive days. The animal need not be killed, but bled again and again, as in the preparation of the diphtheritic antitoxin. The blood should not be brought to too high a point of efficiency, as it will then require too high a dilution for practical purposes. The blood to be tested is diluted with water, one to one hundred, and filtered. Of this clear, slightly red solution two cubic centimetres are placed in a small tube and mixed with an equal quantity of 1.6 per cent. salt solution; six to eight drops of the serum of the rabbit are then added to each tube to be tested, but all will remain perfectly clear except the tube containing human blood. The reaction is extremely delicate and can be obtained with very slight traces even of old dried blood. The clouding should occur within thirty minutes in the proportion of 1 to 30, and a precipitate within two hours; other samples of blood remain clear after six hours. The test should always be repeated. Bordet,<sup>2</sup> Deutsch,<sup>3</sup> Wassermann and Schütze,<sup>4</sup> and Dieudonné<sup>5</sup> describe practically the same method as Uhlenhuth, and have obtained similar results. Bordet and Deutsch each claim to have been the first to use this method.

Corin<sup>6</sup> believes that the active principle of the serum in the biologic differential diagnosis of the blood is paraglobulin, for not only may blood-serum be used for the purpose, but also transudates containing globulin. The paraglobulin in an ascitic fluid was precipitated by magnesium sulphate, dried, and injected into animals in an aqueous solution. In like manner the paraglobulin can be precipitated from the blood of the animal experimented upon and preserved in pulverized form. This powder when wanted for use is dissolved in water and employed in testing the blood under examination. Biondi<sup>7</sup> finds

<sup>1</sup> *Spitalul.*, 1902, vol. xxiii, p. 377.

<sup>2</sup> *Annales de l'Institut Pasteur*, 1899, pp. 225 and 273.

<sup>3</sup> *Orvosik Lapja*, 1901, no. 11.

<sup>4</sup> *Berl. klin. Wehnschr.*, 1901, vol. xxxviii, no. 7, p. 187.

<sup>5</sup> *Münch. med. Wehnschr.*, 1901, vol. xlviii, no. 14.

<sup>6</sup> *Vrtiljschr. f. gerichtl. Med.*, 1902, vol. xxiii, p. 61.

<sup>7</sup> *Vrtiljschr. f. gerichtl. Med., Suppl.-Heft*, 1902, vol. xxiii, p. 1.

*post mortem*, are: (1) Marked reduction in the number of red corpuscles (to one million or less per cubic millimetre). (2) Alteration in their shape,—poikilocytosis. (3) Alteration in size,—microcytes, macrocytes, megalocytes. (4) Nucleated reds,—normoblasts, megaloblasts. (5) Increase of neutrophilic whites. (6) Hæmoglobin markedly decreased, but color-index usually high and blood of a raspberry-red color. (7) Blood-plates absent or scanty. At the post-mortem the skin is, as a rule, lemon-yellow in color. The skin and the serous membranes commonly reveal hemorrhages, which may, however, be present only in the retina. Certain brown discolorations are often found, especially on the abdomen and buttocks. The subcutaneous fat is well preserved and of a light-yellow color. The muscles resemble horse-flesh and are often degenerated. The heart is usually large, flabby, empty, intensely fatty, and tawny-brown in color. The other organs exhibit fatty changes. The stomach may be normal or the disease may be associated with chronic gastritis, gastric carcinoma, or atrophy of the gastric tubules. Iron is deposited in excess in the lobules of the liver, especially in the outer and middle zones. The spleen and hæmolymp glands show a marked leucocytosis and excess of iron pigment. The spinal cord may show extensive posterior sclerosis with hemorrhagic foci, due to the action of the toxins and nerve-fibre degeneration. The lesions are usually most marked in the part of the tract farthest from the trophic centre. Changes in the ganglion-cells of the sympathetic system have been reported. The marrow of the long bones is reddish, resembling that seen in the infant. In pernicious anæmia there is incomplete formation of serum. It is stated that the clot of pernicious anæmia and that of anæmia secondary to cancer may be distinguished by the contraction of the cancer clot and the resulting squeezing out of the serum.

*Chlorosis*.—Chlorosis is a primary anæmia which occurs usually in girls between fifteen and twenty years of age, and is characterized by a marked diminution in the percentage of hæmoglobin, by alterations in the number, shape, and size of the red blood-corpuscles in severe cases, and sometimes by hypoplasia of the circulatory and generative organs. The white blood-cells rarely show much variation. Cases of simple chlorosis rarely come to autopsy. Subcutaneous fat is usually well preserved or even increased in amount. The skin is pale and of a greenish hue, and other evidences of anæmia may appear. Areas of pigmentation, particularly about the joints, occasionally



occur. The internal organs will be found pale and flabby. The heart, large blood-vessels, and generative organs may show insufficient development. (Virchow.) The heart and large veins are often filled with a greenish clot. Thrombi, at times multiple, are common, especially in the femoral vein and the longitudinal sinus. Pulmonary embolism has been observed.

*Leucocythæmia (or Leukæmia).*—A primary anæmia characterized by a great increase in the number of the white corpuscles, by an alteration in the relative proportions of the various white corpuscles the one to the other, and by marked structural changes in the lymphatic glands, spleen, and bone marrow. It may be (*a*) splenic, (*b*) medullary, (*c*) lymphatic, or (*d*) mixed. As a rule, the body is apparently well preserved, but in some cases emaciation may be extreme, while in others the amount of adipose tissue may be increased and of a peculiar punctate appearance, owing to the presence of petechial hemorrhages. The skin is waxy and has a peculiar lemon-yellow color. The mucous membranes are blanched and œdema is often present. The blood is pale, even grayish, in color, the hæmoglobin being often reduced one-half or more. It rarely clots with any degree of rapidity, and in the clot red cells settle, leaving a white film above. At the postmortem the heart and large veins may be found distended with large, greenish, pus-like blood clots. In splenic anæmia microscopic examination of the blood shows that the increased white corpuscles are largely myelocytes, while in the medullary form they are lymphocytes. The white corpuscles are enormously and permanently increased, so that one white to twenty red, or even one to one, is not uncommonly found. The organs in general are pale; the heart is flabby and frequently fatty in appearance. The liver, spleen, and lymphatic glands are usually markedly hypertrophied, while the thyroid may be normal or but slightly enlarged. The thymus gland has in several instances been found enlarged. Lymphoid masses are seen in the lungs. The Peyer's patches are often increased in size.

(*a*) In splenic anæmia, which is a comparatively rare form of the disease, the spleen is markedly enlarged, somewhat firm in consistency, and of a reddish-brown color. The capsule is thickened and the whole organ is bound down by adhesions. The Malpighian bodies are frequently obliterated, their place often being taken by grayish-white, circumscribed tumors throughout the organ. The hyperæmia in some cases is so excessive that rupture of the spleen is said to occur



from this cause. The vessels at the hilum are enlarged. Dropsy from pressure on the abdominal viscera may result. As in other forms of leukæmia, the bone-marrow may show decided changes, especially in the long bones. Instead of fatty tissue there may be splenization, or the marrow may resemble the consistency of the matter which forms the core of an abscess. (b) Medullary leukæmia very seldom occurs as an inflammatory process. Where the marrow changes are excessive, the flat bones—as, *e.g.*, the sternum—undergo alterations similar to those occurring in the long bones. There is a hyperplasia of the red marrow; this may resemble pus or be of a dark-brown color. There may be localized swelling of the bone. (c) In lymphatic leukæmia the lymphatic glands throughout the body, especially those of the neck, the axillary and inguinal regions, also the glands of the mesentery and the intestines, are swollen, pale in color, firm to the touch, but seldom suppurate or show any tendency to run together. The spleen, liver, and lymphatic glands, as the tonsils, lymph-follicles of the tongue, pharynx, and mouth, often show marked thickening of their capsules. On section the glands are somewhat resistant, and often exhibit nodule-like bodies, which are firm in consistence and largely composed of proliferating leucocytes and connective tissue. The liver as well as the spleen is enlarged and may exhibit marked structural changes. This form may be associated with, or most difficult to differentiate from, lympho-sarcoma.

*Von Jaksch's Anæmia.*—This is a primary anæmia of infancy, closely resembling leukæmia, but without the visceral lesions. The red cells are diminished, though many of these are nucleated. The spleen, liver, and lymph-nodes are enlarged, and the number of leucocytes is increased.

*Osler's Disease (Chronic Cyanosis).*—A chronic disease of people usually past middle life and not associated with dyspnœa, kidney, lung, or heart disease, but characterized by marked blueness of the skin. There is a marked polycythæmia, the red cells varying from 10,000,000 to 12,000,000 per cubic millimetre and the hæmoglobin being increased as much as fifty per cent. above the normal. At the postmortem the heart and spleen are found enlarged and the internal organs markedly congested. At times small hemorrhages are noted.

*Hodgkin's Disease (Pseudoleukæmia; General Lymphadenoma).*—A disease characterized by a progressive hyperplasia of the lymph-glands, by anæmia, and sometimes by secondary lymphoid growths in

the liver, spleen, and other organs, but with no severe leucocytosis. The lymphatic glands most frequently enlarged are those in the cervical, axillary, and inguinal regions, though the mediastinal, thoracic, and abdominal glands, especially the retroperitoneal, are often affected. In the early stages the glands are moderately enlarged, soft and elastic, isolated, and freely movable. Later they increase in size and tend to run together, become stony hard, and are surrounded by a dense capsule. The capsule may perforate, and the growth invade the surrounding structures. On section the tumor appears grayish-white; it is smooth and the interior may be firm and dry or soft and juicy. Suppuration sometimes occurs when the growth reaches the skin. Emaciation at the time of death may be extreme. The spleen and liver are usually somewhat enlarged, but rarely greatly so, and on section show lymphoid tumors varying in size from that of a pea to a walnut. The Peyer's patches frequently show enlargement. Pleural effusions are not uncommon. The skin may be the seat of adenoid growths. The glandular enlargements may be due to simple inflammatory hyperplasia, lymphadenoma, or lymphosarcoma. The bone-marrow may be converted into a rich lymphoid tissue. The blood-changes are those of a distinct anæmia of the simple type. The red cells are less numerous and are slightly smaller; the hæmoglobin is always diminished; the leucocytes are normal or decreased in number.

*Hæmophilia*.—An hereditary constitutional disease characterized by a marked tendency to excessive hemorrhage from very slight causes. It is transmitted through the females of a family to the males. Little regarding its morbid anatomy is definitely known, and therefore any opportunity for the study of a case *post mortem* should be taken advantage of. The vessel-walls are unusually thin, brittle, narrow in calibre, and do not readily contract. In some cases the blood itself presents marked alterations. Owing to the ease with which the joints are injured, hemorrhages are often found about the capsules of joints, with inflammation of the synovial membranes. In a few cases increase in the number of red cells and diminution of white cells and blood-plates have been noted. Geier<sup>1</sup> finds the cytoglobin, which is produced by the destruction of the red blood-cells, to be markedly increased in hæmophilia.

*Purpura*.—This is characterized by extravasations of blood into

---

<sup>1</sup> *Med. Obozrainie*, Mosk., 1904, vol. lxi, no. 1.

and from the skin, by great debility, evidences of anæmia, and often multiple arthritis. Infectious purpura is seen in pyæmia, septicæmia, malignant endocarditis, typhus fever, etc. The forms are (a) purpura simplex, (b) purpura hæmorrhagica, (c) purpura rheumatica, (d) iodic purpura, (e) Henoch's purpura, (f) neurotic purpura, (g) mechanical purpura, (h) toxic purpura (seen in snake poisoning, after the use of certain medicines, etc.), (i) cachectic purpura. The blood clots slowly and imperfectly; leucocytosis may or may not be present; the blood-plates may be scanty; and the red cells are often reduced in number. There is a large percentage of lymphocytes and an increase in the eosinophiles. The skin is dry and pale, except for the blotches of extravasated blood, which vary from one to four millimetres in diameter, are bright red in color, later become dark, and finally remain as brown stains. The hollow viscera and serous cavities may contain considerable quantities of blood-stained serum. The serous membranes and solid organs may also reveal hemorrhages varying in size from a pin's head to the palm of the hand. Congestion and œdema of the lungs are frequently present. There is generally an acute diffuse nephritis. Ulcerations of the intestines with enlargement of the solitary and agminated glands are sometimes found. In one of my cases, in which the purpuric blotches were unusually large and widely distributed over the body, a husband was accused of beating his wife and thus causing her death.

*Scurvy.*—Scurvy is a constitutional disorder characterized by anæmia, great debility, spongy gums, and tendency to hemorrhage. This disease is by no means so frequent as formerly, owing to better hygienic conditions and to the proper feeding of those in ships, prisons, work-houses, etc. The blood is dark and fluid; there is a decrease in the number of the red cells, many of which are pale and distorted; microcytes are present; there is no leucocytosis. After death decomposition sets in rapidly. There is very little wasting of the subcutaneous fat or of the muscles. The hemorrhagic patches observed in the skin during life are often obscured by post-mortem lividity; œdema is common. The subcutaneous tissues, especially those of the lower extremities, contain a blood-stained fluid, with here and there discolored patches, some black and others of a pale color. About the back of the thigh and knee the muscles and tendons may be embedded in a thick, firm clot, and themselves contain numerous hemorrhagic foci. Occasional hemorrhages occur within the joints, or into any of



the serous and mucous membranes or internal organs, especially the kidney and bladder. The gums are swollen and may present fungous appearances; they are sometimes ulcerated, and the teeth may have fallen out. Rarely there may be ulcers in the intestines. Hemorrhagic infarcts are at times seen in the lungs and spleen, the latter organ being enlarged and soft, while fatty changes are quite constant in the liver, kidneys, and heart.

*Scurvy, Infantile (Barlow's Disease).*—Usually associated with improper food, such as too much malted or condensed milk. Cases, however, have been reported in breast-fed children. The most important lesions are increased vascularity and extravasation of blood between the periosteum and the bone and into the cavity of the long bones, especially those of the lower limbs, producing tumor-like swellings. Epiphyseal fractures are not uncommon. In fact, in the majority of cases there are bone changes analogous to those of rickets, and the disease often develops in a rickety child.<sup>1</sup> Deep-seated extravasations may give rise to muscular swellings and in some cases to extravasations in the joints. Smaller extravasations are observed in the pleura, lungs, spleen, intestines, and kidneys. The gums are spongy, sodden, distended with serum, and sometimes covered with blood. One of the most characteristic lesions is extravasation of blood into the orbital cavity, causing displacement of the eyeball downward and forward.

*Diabetes Insipidus.*—A constitutional condition characterized by continued secretion of large amounts of pale urine, of low specific gravity, containing neither albumin nor sugar, attended with excessive thirst and at times with emaciation; usually the patient looks well nourished. It occurs most often in young males and is usually hereditary. The urinary system may show merely signs of the passage of an abnormal amount of liquid,—enlarged and congested kidneys, dilated pelves, dilated ureters, and an hypertrophied bladder.

*Gout.*—A constitutional disease characterized by excessive formation of uric acid and the gradual deposition of its salts, especially sodium urate, in and around the joints of the extremities, producing an acute arthritis. Anatomical changes are found most frequently in the great toe, though the disease shows a marked tendency to involve the smaller joints, both of the feet and the hands. In acute stages there are notable hyperæmia and round-celled infiltration and diffusion

<sup>1</sup> *Lancet*, May 3, 1902, p. 1246.



into the joint and swelling of the ligaments. Macroscopically the joint is swollen, glazed, tense to the touch, and of a purplish color. In the chronic form the ligaments and fibrocartilages of the joint become infiltrated with chalky deposits (tophi). These consist of sodium urate in the form of crystalline needles or rhombs, which are quickly dissolved by hydrochloric acid, but whetstone-shaped crystals of uric acid make their appearance. Necrosis in the cartilage always precedes the formation of tophi (Ebstein). These deposits may be slight or may lead to enormous distortion of the joint. In some cases the skin may ulcerate and the tophi be extruded. The deposits may be found in the cartilages of the ear, the nose, the eyelids, and occasionally the larynx. In some cases the synovial fluid contains crystals. In chronic cases the joint becomes immovable, due to the exostosis and excess of deposits. The kidneys usually show chronic interstitial inflammation, with deposits of urates in the form of small flakes or stripes, chiefly in the pyramids. Arteriosclerosis, with hypertrophy of the left ventricle, is very common. Cutaneous affections, such as eczema, are not infrequent. The blood contains an excess of uric acid.

*Varieties of Hemorrhage.*—The following terms are applied to hemorrhages from various parts of the body: Epistaxis, hemorrhage from the nose; hæmoptysis, pulmonary hemorrhage; hæmatemesis, or gastrorrhagia, hemorrhage from the stomach; enterorrhagia, hemorrhage from the intestine; metrorrhagia, uterine hemorrhage not occurring during the regular menses; menorrhagia, excessive menstrual flow; post-partum, hemorrhage from uterus after delivery; complementary, hemorrhage occurring in some place other than that in which the original bleeding occurred; consecutive or secondary hemorrhage; extrameningeal, a hemorrhage external to the cerebrospinal meninges; hemorrhage per diapedesis; hemorrhage per rhexin.

*Hemorrhages, Causes of.*—(a) Traumatism. (b) Acute inflammation. (c) Passive congestion. (d) Corrosive poisons. (e) Malignant growths. (f) Diseases of the vessels. (g) Rupture of an aneurism. (h) Cachetic disease. (i) Dyscrasias. (j) Nervous disturbances. (k) Vicarious menstruation. (l) Toxins.

Many coroners' physicians give hemorrhage from the umbilical cord as a cause of death in new-born children. Although this fatality does occur, it is extremely rare, some obstetricians treating without tying the cord hundreds of cases without hemorrhage; nor is the condition seen in the lower animals. It is facilitated by cutting the cord

too close to the abdomen, by forced artificial respiration, and by the presence of hæmophilia. It may come on several days after birth, and at the postmortem the liver appear especially blanched.

**ANGINA PECTORIS.**—A symptomatic affection commonly associated with more or less myocardial degeneration and occlusion of the coronary arteries from atheroma and thrombosis. At the autopsy the heart is often enlarged and the pulmonary artery and the cavities of the heart are filled with post-mortem clots. While aortic and mitral thickenings are usually present, I have examined cases where they were absent. The coronary arteries are "pipe-stem" in character, the anterior one being usually most markedly affected.

**INFILTRATIONS AND DEGENERATIONS.**—In fatty infiltration, or *obesitas cordis*, there is an increase of fat in those places where it is normally deposited, especially along the grooves of the larger blood-vessels. The deposits start from the outside and extend inward along the trabeculae of connective tissue, while in fatty degeneration the changes originate from within. The heart may be embedded within such an enormous deposit of fat as to leave no muscle exposed to view. Fatty infiltration and degeneration occur most markedly in cases of poisoning, as by phosphorus, and the atrophy of the muscle may be very extensive. In such cases the heart is so soft that the finger can readily be pushed through its walls. Hyaline and amyloid degeneration may also occur, as well as calcareous infiltration, fragmentation, and segmentation. (See Myocarditis.) Brown atrophy is common; the degenerated fibres are dark brown in color, contain yellow-brown pigments within the muscle-cells, and the cavities are decreased in size. An atrophy of the left ventricle is sometimes seen in cases of extreme mitral stenosis. Senile atrophy always accompanies fibrosis. Fibrosis occurs most often in the aortic valves; the corpora Arantii are first affected, later the chordae tendineae become thickened, first at the valvular ends. Papillary muscles may also become markedly fibroid. (Osler.)

As a result of degeneration spontaneous rupture may occur, usually in the anterior wall of the left ventricle. This results from fatty infiltration, degeneration, gumma, or tuberculosis. It has been found associated with abscess, aneurism, ulceration, myomalacia, arteriosclerosis, and thrombosis. Fatty degeneration may end in rupture (spontaneous) of heart. Rupture may be due to trauma. Blows upon the chest may rupture the heart and also cause localized myocarditis,

injury to mitral leaflets, or tear holes in valves where the chordæ tendineæ had been attached. This also occurs from extreme action or blows, gunshot wounds, etc.

Aneurism of the heart itself is usually due to myomalacia, with thickening and narrowing of coronary arteries and chronic myocarditis, often associated with valvulitis, syphilis, etc. It occurs usually in the left ventricle near the apex, or may be found in the intraventricular septum or posteriorly. The endocardium is usually opaque, the muscles are sclerotic, and layers of thrombi are found in the sac. The aneurism may or may not be lined with endothelial cells. Two aneurisms may be found existing in one heart. Now and again an aneurism appears on the valves of the heart, and is then spheroidal and projects from the ventricular face of the semilunar valve. Literature is full of reported cases of cardiac and aortic aneurisms, there being several pages devoted to this subject in the *Index Catalogue*.

MYOCARDITIS.—In the myocardium large hemorrhages may be met with, as a result of the rupture of small aneurisms of branches of the coronary arteries or as a hemorrhagic infarct. Anæmic infarct may also be due to a partially obstructing embolus or the formation of a thrombus or to disease of the coronary artery. It usually occurs in the left ventricle, at the apex, or in the septum. It is irregularly shaped, yellow-white in color, and sometimes turbid or parboiled in appearance. This is a common cause of sudden death. Tardieu's spots, or small hemorrhages beneath the endocardium and at times extending into the muscle, are found especially after suffocation and in cases of rapid death from acute infectious fevers. Myocarditis usually is secondary to inflammation of the heart muscle. Parenchymatous myocarditis may be diffuse or limited. When the inflammatory process involves all of the musculature of the heart, as is frequent in the infectious diseases, it is characterized at first by the flabbiness and the turbid grayish-red color of the heart muscle. In the later stages there is much fatty degeneration. Segmentary parenchymatous myocarditis is marked by a cloudy appearance of the heart muscle, which is flabby and friable. (Orth.) Fibres may separate at the cement line. Transverse fragmentation of the fibres is the form which usually occurs during the death agony. Acute circumscribed interstitial myocarditis, or abscess of the heart, is usually a part of a general pyæmic disease, with infection through the coronary circulation. These metastatic abscesses occur in cases of puerperal sepsis, with



osteomyelitis and other intensely septic diseases, but particularly in cases of malignant endocarditis. There may be only a few abscesses or the heart substance may be studded with innumerable suppurating points. In size the abscesses vary from the merest dots to cavities of the size of a cherry; they may perforate or form ulcers in the cardiac wall. Acute diffuse interstitial myocarditis occurs in various forms of infectious fevers. The affected heart muscle is soft and often distinctly friable; there may be spots of hemorrhagic infiltration, but, as a rule, the color is rather lighter than that of the normal organ. The cavities of the heart are frequently dilated, particularly the left ventricle. Chronic interstitial myocarditis or fibrous myocarditis may also be diffuse or localized, though the circumscribed form is the most common. The process is usually secondary, due to a primary disease of the coronary arteries, or to disturbances of the circulation therein, consequent perhaps upon old age, intemperance, gout, syphilis, and the like. This fibroid overgrowth is very commonly met with at the tips of the papillary muscles, on the trabeculæ, or in the substance of the cardiac muscle, and often at the apex of the left ventricle, where it may lead to such a degree of atrophy that a chronic localized aneurism of the heart may be formed by the constant pressure of the blood upon this thinned area. The heart is usually hypertrophied and the cavities dilated. The characteristic change is the formation of dense, grayish sclerotic areas, which appear either as more or less irregular spots or as streaks or lines running in the direction of the cardiac fibres. The entire substance of the heart may be involved and thickening of the walls may result. (Stengel.) The condition of softening of the organ, or myomalacia cordis, has already been referred to. The degenerated tissue may form a scar, but more frequently leads to an aneurismal dilatation, which may subsequently rupture. Aneurisms of the sinus of Valsalva may form and rupture in unexpected places; I have seen, for example, an aneurism of an aortic sinus rupture into the right ventricle.

**ENDOCARDITIS.**—Disturbances of the circulation of the endocardium are rare, as this membrane possesses no blood-vessels of its own. A diffuse redness in this situation may, however, be the result of imbibition, and in the case of long-diseased valves, in which there are newly formed blood-vessels, reddish streaks and spots may be observed, which are due to small hemorrhages. Inflammation of the inner lining of the heart is frequently a secondary affection, dependent



upon inflammatory disorders of other organs, such as suppurating wounds, purulent peritonitis, and pneumonia, or to rheumatism, gonorrhœa, chorea, tuberculosis, cancer, etc. The most common organisms found are the various forms of cocci. Sometimes, however, the endocarditis is the first local manifestation of an infection, the exciting agent of which has left no recognizable traces at the seat of its entrance into the body. In the fœtus endocarditis is usually situated in the right side of the heart, because the blood enters the organ on that side, and may be associated with lesions of the ductus Botalli. After birth the opposite condition prevails, the lesion being most commonly found on the left side. In the great majority of adult patients acute endocarditis affects the endocardium of the valves only,—the mitral, the aortic, and the pulmonary valve in order of frequency; but it is sometimes found in the endocardium of the cavities of the heart,—in the left ventricle, the left auricle, and the right ventricle. Various names have been applied to these conditions, as simple, verrucose, benign, ulcerative, septic, mycotic, rheumatic, syphilitic, diphtheritic, fibrous, or malignant endocarditis. Such cases differ much in their appearance, even when produced by the same organisms. Endocarditis starts on the endocardium as a minute, roughened area, which is red in color and slightly elevated. This can easily be scraped off, but, if the spot where it was found is carefully examined with a hand lens, a small ulcer will be seen. More and more fibrin is now deposited, and the corpuscular elements are caught in its meshes; the organisms multiply and the clot undergoes a liquefaction necrosis, the process not stopping in the newly formed tissue but often penetrating the valves or even the walls of the heart. Embolic occlusion of certain vessels and metastatic inflammations in other organs, especially the kidneys, spleen, brain, lungs, meninges, and skin, are not infrequently associated with endocarditis (Ziegler). Such hemorrhagic areas are to be sought for in the palpebral conjunctiva; their discovery therein during the external examination of the body has more than once led me to suspect ulcerative endocarditis, even when there was no clinical history of its existence. This previous observation is of special value when a bacteriologic examination of the heart is desired. These ulcerative areas of valves on healing are replaced by scar tissue, which, by contraction and by various degenerative changes, such as necrosis, fatty degeneration, and calcification, give rise to the most fantastic shapes and appearances of the parts affected. Old cases are often associated

with aneurism of the leaflets, dilatation, pouching, or perforation of valves. Often small tumor-like masses remain on the leaflets, which become thick, rigid, and calcareous.

**HYPERTROPHY AND DILATATION.**—These conditions are usually associated the one with the other. In concentric hypertrophy the cardiac walls are thickened and the cavities are smaller than normal. As this may be due to post-mortem contraction or to marked systole, it may be well in some cases to soak the heart in tepid water before the measurements are taken. In simple hypertrophy the overgrowth of the walls is associated with normal cavities, while in eccentric hypertrophy dilatation is found along with the thickening of the walls. The highest degrees of enlargement which I have seen have been found in cases of double aortic disease, where, too, moderator bands are sometimes found. It would seem that the long-continued administration of digitalis may produce hypertrophy.

**VALVULAR DISEASES.**—An extreme degree of mitral stenosis is seen in the so-called buttonhole mitral, a condition more frequently observed in England than it is in this country and which causes hypertrophy and dilatation of the left auricle. Cyanotic induration of other viscera, especially of the lungs, and dropsical effusions may follow mitral incompetence. In aortic stenosis the valves are usually thickened, rigid, and cartilaginous; later they become calcified and the division between the different cusps is lost. First there is ventricular hypertrophy, later right-sided enlargement, and finally dilatation with pulmonary congestion. In aortic incompetency arteriosclerotic changes are marked, being seen not only in the valves but also in the aorta, and associated with dilatation and hypertrophy of the left ventricle and of the left auricle, and often followed by fibroid myocarditis. Sudden death is frequently due to aortic stenosis, a condition usually associated with hypertrophy of the left ventricle and a dilated cavity. The aortic ring and segments are atheromatous, puckered, and contracted, often calcareous, and may admit only the tip of the little finger. The aorta above the structure is usually dilated. Tricuspid regurgitation is seen generally associated with cases of cirrhosis of lung and chronic emphysema. Cyanosis is common. Pulmonary-valve disease is rare, except as a congenital lesion. Stenosis is usually associated with patency of the ductus Botalli.

**SYPHILIS, TUBERCULOSIS, ACTINOMYCOSIS, TUMORS, ETC.**—Syphilitic gummata appear in the heart as rather large yellow foci sur-

rounded by fibrous tissue; they may also be found in the arch of the aorta. Miliary tubercles, when present, are usually subendocardial or situated in the large vessels coming off the heart. At a postmortem in Ziegler's mortuary I once saw where caseation of a peribronchial gland had extended through the pulmonary artery and given rise to a most marked local and general miliary tuberculosis. Actinomycosis has been observed. Tumors are rare; myxomata, lipomata, fibromata, fibrous polyps, sarcomata, and rhabdomyomata may be met with as primary tumors of the heart, while, as secondary deposits, carcinomata and especially multiple melanotic sarcomata may be observed. (Plate IV.) In v. Pessl's case<sup>1</sup> of extensive lymphosarcomatosis, the heart at autopsy showed the anterior wall of the left ventricle and a large part of the septum almost converted into a "shell of lime." A carcinoma may thus develop here secondary to one of the penis.<sup>2</sup> Foreign bodies, as needles, hat-pins, pieces of bone, etc., have been found in the cardiac wall and even in the cavities of the heart. Cysticerci, echinococci, and very rarely pentastomata may be discovered in the various parts of the heart.

ARTERIES, MORBID CHANGES IN.—*Arteriosclerosis*.<sup>3</sup>—A chronic thickening and hardening, diffuse or circumscribed, of the arteries, characterized by a diminution of elasticity of the vessels and marked alterations in blood-pressure. The arch of the aorta is the most common seat. In the *first stage* there are a loss of elasticity and some dilatation, due to hyaline or other changes in the subendothelial coat, and thickening of the intima. The *second stage* is characterized by a thickening of the media, atrophy of the muscular and elastic tissue, with proliferative changes in all the coats; the increase of new tissue gives rise to pressure on the vasa vasorum, with interference of nutrition, which leads to the *third stage*. New elastic tissue, derived from the splitting off of the internal elastic layer of the artery, may be found in the intima. This consists of more or less marked macroscopic changes. The vessels are hardened, firm to the touch, do not retract or close when cut, and their lumina may be smaller or larger than normal. On the intima may be seen milk-white or yellowish patches, containing fat, cholesterin, and detritus, intermingled with calcareous


---

<sup>1</sup> *Münch. med. Wchnschr.*, June 10, 1902, p. 956.

<sup>2</sup> PEPPER, *Phila. Path. Soc.*, meeting of November 12, 1903.

<sup>3</sup> WELCH's paper, read at the June, 1904, meeting of the American Medical Association, should be consulted by those interested in this important subject.

# PLATE IV



1, fibroma; 2, chondroma (after Ziegler); 3, cavernous angioma of the liver; 4, myoma; 5, glioma of the brain; 6, giant-celled sarcoma; 7, spindle-celled sarcoma; 8, endothelioma of the pia mater; 9, adenoma; 10, cancer; 11, epithelioma with an epithelial pearl; 12, myxomatous cancer.





plates and areas of ulceration. There may be a marked tendency to dilatation, with the formation of an aneurism, or to contraction with obliteration. Some cases are associated with fibrosis of the aortic valves. As described by Heller, this form when found in the thoracic aorta is usually syphilitic. *Special Forms.*—(a) Senile. To a certain degree this may be regarded as physiologic, the elastica being developed as in scar tissue. The condition affects the larger arteries most; they are dilated, lengthened, tortuous, thin but stiff; often show atheromatous changes in the intima. Even to the naked eye the subendothelial tissue is degenerated. Cyanotic induration and senile atrophy of the heart, liver, and kidneys are common. Moist gangrene of the extremities may follow calcification in the iliac arteries. The changes produced by sclerosis of the coronary artery have already been referred to. (b) Nodular. Knob-like, flat, yellowish-white projections are seen in the aorta and its branches, particularly about the orifices. These in later stages undergo liquefaction and form atheromatous ulcers. Dilatation or aneurism may then ensue. (c) Diffuse. The lesion is wide-spread and more uniform; the intima, as a rule, does not show marked naked-eye changes, though there may be elevated spots of an opaque white color. The aorta and its branches are dilated, the branches sometimes more than the trunk. Cardiac hypertrophy is constant. The kidneys are sclerosed, their capsule is adherent, the cortex irregular and often cystic. (d) Endarteritis obliterans. There is particularly a thickening of the intima, the entire lumen of the vessel being closed. It is not uncommon at the base of the brain. Erythromelalgia is largely due to arteriosclerosis. Lead and alcohol are considered to be frequent causes of arteriosclerosis.

*Amyloid Degeneration.*—Usually microscopic, and best demonstrated by staining. *Atrophy.*—A general diminution in the size of arteries, best seen in stumps after operations. *Calcareous Infiltration.*—In the media of the arteries of the old, particularly involving those of the extremities, calcification of the media occurs. It interferes with the flow of the blood, predisposes to thrombosis, and may be the cause of senile gangrene. Seen as a diffuse or circumscribed process, usually in connection with atheroma, and as a later stage of fatty degeneration. *Fatty Degeneration.*—In fatty degeneration the affected areas of the intima have a white or a citron-yellow appearance. These areas occur in the form of points, stripes, and regular or irregular net-shaped figures. A frequent location is the posterior wall of the aorta around the origin

of the intercostals. For more careful macroscopic study the surface of the fatty area should be removed with a fine forceps and in the centre a shallow incision should be made. Here small and large fat droplets can be seen. When placed in Flemming's solution these droplets become black. The fatty degeneration may involve the media and even be the cause of rupture. Often associated with calcareous degeneration. *Hyaline Degeneration*.—Almost always microscopic; affects mostly the elastic coat and is often the beginning of an arteriosclerosis. It most frequently involves the small arteries. Best seen in the glomeruli of the kidney. *Hypertrophy*.—There is hypertrophy of the muscular layer in some diseases of the kidney, and hypertrophy of this layer in arteries of medium size in aortic insufficiency. A general enlargement, best seen in the collateral circulation after ligation of a large vessel. *Hypoplasia*.—Hypoplasia of the aorta is congenital and is the result of stenosis, most commonly situated near the insertion of the ductus arteriosus Botalli. It usually soon causes death; if not, the aorta is contracted and thinner, but very much more elastic. Virchow attributes chlorosis to it.

*Inflammations*.—I. Acute endarteritis (proliferative or obstructive endarteritis, thrombo-arteritis). This starts with an injury to the endothelium, proliferation occurs, and an obstruction is formed in the vessel wall, on which a thrombus forms, partially or completely obstructing the vessel. The intima is yellow and may be covered with many arcuate limited or confluent ulcers, with a loss of substance. Cholesterin and fat cells are found in the detritus. The lesion may terminate in absorption, suppuration, ulceration, or fibroid change. II. Chronic endarteritis. This usually follows the acute form, but is sometimes primary. It may be local (organization of a thrombus) or general (arthritis deformans). Thrombosis is more common in the veins; it is usually caused by embolus from the right heart or from the left and associated with hemorrhagic infarct. Cases have been reported of thrombosis in the aorta above its bifurcation, in which the aorta and the femoral and popliteal arteries also revealed a marked thrombosis. It may occur in a vein during pregnancy, chronic nephritis, rheumatism, diphtheria, or typhoid fever. Marantic thrombi may form in cases of enfeebled circulation, as in the cachexia and anæmia of severe infections, sometimes associated with arteritis obliterans. In these cases it occurs usually in the large superficial veins of the lower extremities and in the sinuses of the brain. Thrombosis of the superior

mesenteric artery, with a slate-colored to black gangrene of the small intestines for many feet, is sometimes seen. The thrombi are gray-white, red, or mixed; if recent, they are attached, of bright-yellow color and membranous consistency; if old, they are firmly organized and may be calcareous. Thrombus is generally supposed to be due primarily to bacteria. Thrombi are not at all rare in the heart, occurring usually in the auricle of the right side or the apices of the ventricles between the papillary muscles. They often have a root-like process, which is especially marked in the appendix, and are most frequently found in cases of cardiac aneurism and in endocarditis.

*Syphilis* is usually a diffuse process affecting all the coats, especially the intima, and thus giving rise to a local or general sclerosis. Gummata are rare. Orth, quoting from Heller, gives the numerous small foci of cell-infiltration necrosis, and particularly induration, with small thickenings of the inner surface, as the characteristic differences between syphilis and chronic aortitis.

*Tuberculosis*.—Tuberculous lesions are less common in the arteries than in the veins. The small arteries are most frequently affected, as those of the pia, the brain, the kidneys, and particularly the lungs. The disease starts as a local gray tubercle of hæmatogenous origin in the intima or as an extension from a neighboring tuberculous process. *Thrombosis and embolism*, especially in the brain, are of extreme importance. In embolism air, fat, portions of tumors, micro-organisms, etc., are brought to a smaller vessel from a larger one (though the converse may occur), and there set up characteristic changes, as infarcts, softening, abscesses, etc.

*Tumors*.—The new growths found in arteritis are angioma, an erectile tumor; the capillary nævus is usually found on the face, capillaries dilated and tortuous; if made up of dilated and tortuous arteries it forms a racemose or cirroid aneurism and is usually seen in the subcutaneous tissue. The skin over the tumor is, as a rule, very much thinned. An arterial varix consists of a dilatation and lengthening of a single artery. Venous nævus is a tumor made up of communicating spaces lined with endothelium into which arteries empty and from which veins arise.

*Aneurism*.—An aneurism is a circumscribed, tumor-like dilatation of an artery, containing blood in direct connection with the blood-current. A *true aneurism* has a sac composed of one or more of the arterial coats. A *false, spurious, or hernial aneurism* is one in which



some of the walls are formed by the tissues surrounding an opening in the artery; these sometimes attain an enormous size. The aneurism is called *cylindrical* when there is widening in all directions; *saccular* when one side is affected; *cirroid* when a large extent, or even the whole ramification, of an artery becomes dilated and tortuous; this form most often occurs in the frontal, occipital, or iliac arteries; *fusiform* when it is spindle-shaped; sometimes these form near a tear of the intima, the media and adventitia becoming markedly sclerosed (arteriosclerotic). A *traumatic* aneurism results from injury or laceration of the intima by force. A *dissecting* aneurism arises when blood circulates between the coats of an artery. Extensive degeneration must precede this form. I have seen such an aneurism beginning at the transverse arch of the aorta and opening again into the blood-stream just above the aortic bifurcation. *Arteriovenous* aneurism arises where there is a communication existing between an arterial aneurism and a vein; a variety of this is the varicose, where an artery and a vein communicate through a false aneurism lying between them. When bleeding was more frequently practised than it is now, these aneurisms in the arm were of quite common occurrence. *Mycotic* or *infective* aneurisms are multiple and are caused by micro-organisms. This variety is often seen in connection with malignant endocarditis. The mesenteric arteries of the horse sometimes become dilated with considerable numbers of the *Strongylus annatus*. *Miliary* aneurisms are usually multiple and consist of small dilatations; they are found especially in the brain and lungs and often antedate a hemorrhage in these regions. They are best seen in the brain by excising the middle cerebral or basilar artery and floating it out in a white dish partially filled with water. These aneurisms may be due to emboli. *Traction* aneurisms have been reported by Thoma at the concavity of the arch of the aorta. Aneurism by distention, rupture, erosion, anastomosis, also valvular and congenital aneurisms have been described. Aneurism is often associated with arteriosclerosis, embolism, trauma, and infections. The change is now believed to start in the elastic coat.

The walls of the blood-vessels may be present or altogether absent; they may be thickened and opaque or almost transparent. If the aneurism be large, the cavity has a roughened wall, often lined with endothelium, and frequently contains clots which are white, red, organized, or softening. They often show lamination. I have seen a fibrinous clot of an aneurism of the carotid mistaken for a sarcoma of the neck,

a gluteal aneurism opened for an abscess, and a femoral aneurism mistaken for a hernia. Rupture of an aneurism, usually from the aorta into the pericardium, is a most frequent cause of death in cases brought to the notice of the coroner. The rupture often occurs during the act of defecation. Three cases of aneurism of the sinus of Valsalva have come under my notice. The direction of the increase in size of a forming aneurism depends on its location. Constant pressure of the sac may overcome the resistance and cause absorption of the densest tissue, even bone. Hence aneurisms of the arch of the aorta may rupture externally or erode the vertebral column. It is the sacculated aneurism which is now treated by electrolysis. Life may thus be prolonged, though the aneurism usually appears in another spot. In Stewart's case life was prolonged three and a half years, the man finally dying from an alcoholic pneumonia. The clot around the gold wire may become markedly fibrinous. The danger in the operation would seem to be from an embolus. Aneurisms are most common in the thoracic aorta, abdominal aorta, coeliac axis, splenic artery, very rare in the hepatic artery. I have seen a few cases of that rather rare condition, aneurism of the superior mesenteric artery.

Congenital aneurism, periarteritis nodosa, closely resembles at times sarcoma; nodules may be felt in the abdominal wall, in arteries of muscles, and in the viscera. There is marked thickening of the intima and infiltration of other coats.

The question as to the etiology of aneurism is much debated. My own statistics on this subject confirm the opinion that syphilis is a frequent cause, especially in the early stages before marked arteriosclerotic changes have taken place in the arteries. This view is also supported by the fact that animals are rarely affected with aneurism. The experimental production of aneurism in animals by alcohol, trauma, etc., affords an interesting field for future investigators.

DISEASES OF VEINS.—The more common lesions in veins are *thrombosis* and *phlebitis*. Phlebitis occurs in connection with many cases of thrombosis, in gout, and after traumatism; micro-organisms are usually found. It appears commonly in sinuses of the brain, associated with leptomeningitis. The intima is rough, uneven, corroded. Vein walls are infiltrated and discolored. On the surface over an infected vein there is a dusky-red line. Thrombosis always occurs. Suppuration or pyelephlebitis, associated with thrombosis and with purulent softening, occurs in the portal veins. The vein is filled with puriform

fluid, which may cause a branching abscess all through the liver. A varix is a permanent dilatation above a valve,—long fusiform dilatation (phlebectasia) or cirroid aneurism. It is most common in the leg and is often associated with diseases of the heart, liver, lungs, pleura, or may be caused by pressure of a tumor. Phlebosclerosis is seen at times with dilated veins. The vein in these cases is nearly always distorted and thickened. Thrombi often form in dilated parts and are sometimes converted into phleboliths. Associated with this lesion are œdema, chronic catarrh of a mucous surface, chronic ulceration of a cutaneous surface, pigmentation, and productive fibrosis. Certain varices have special names, as varicocele, dilatation of spermatic veins; hemorrhoids, dilatation of veins of lower rectum; caput medusæ, dilatation of superficial abdominal vessels, etc. Primary tumors are rare in veins. Any tumor may be present secondarily. Often there is a fibrous endophlebitis, etc. Syphilitic inflammation is seen in portal and other veins and is a congenital form. Tuberculosis of a vein from a lymph-node is not uncommon in general miliary tuberculosis.

CHIEF LESIONS FOUND IN LYMPH-VESSELS.—*Dilatation* is especially seen in elephantiasis Arabum and is due largely to filaria. Certain parts, as the scrotum, labia, and thigh, are generally the seat of the lymph stasis. This lesion is always associated with hypertrophy and thickening of the tissues. *Lymph tumors (lymphangioma)*—(1) simplex, not much enlarged, (2) cavernosum, much dilated—are usually found in the tongue (macroglossia), cheek, etc. Lymphangioma cysticum occurs most frequently in the neck or the sacral region. Contents of the cyst are often very fatty, at times partially coagulated. Endothelioma and carcinoma may occur. The lesions of tuberculosis and syphilis are found.

## CHAPTER IX

### DISEASES OF THE RESPIRATORY TRACT AND ACCESSORY PARTS<sup>1</sup>

NASAL PASSAGES.—Malformations of the nose are very common. The entire nose, the septum, or the turbinates may be absent. Deviation of the septum occurs in about one out of every ten persons. *Acute Nasal Catarrh (Rhinitis, Coryza)*.—This condition accompanies various infections, as variola, scarlatina, measles, influenza, röteln, and diphtheria. In infants it is often a manifestation of syphilis or gonorrhœal infection. The nasal mucous membrane is red, swollen, and covered with exudate, the accessory sinus, pharynx, and Eustachian tubes being sometimes secondarily inflamed. Empyema of the antrum of Highmore may result, but more frequently arises from carious teeth. Herpetic eruptions are often seen on the lips. *Chronic Nasal Catarrh*.—Varieties: hypertrophic, atrophic, fibrinous, or membranous. There is usually a persistent mucopurulent discharge. In the hypertrophic form there are flattening of the nasal bridge, thickening of the alæ and the mucous membrane, exostoses on the septum, and hypertrophy of the cavernous tissue. In the atrophic variety the mucous membrane is pale, dry, glazed, and covered with scabs, ulcers, and, at times, necrotic tissue, which may lead to perforation of the septum, but this is much more commonly seen in syphilitic rhinitis. Adenoids are, as a rule, confined to the roof of the nasopharynx, but may occur upon the lateral walls. They are smooth, rounded masses, or pedunculated, varying in size from a hemp-seed to an almond and of a pale-pink color. These may replace entirely the normal pharyngeal tonsil. Nasal tuberculosis, especially in the form of lupus, syphilis, glanders, and leprosy, may occur in the nose. Rhinoscleroma gives rise to nodular thickenings and ulcerations of the tissues of the nose, lips, pharynx, and larynx. *Tumors*.—Fibrous, myxomatous, and cystic growths are by far the most common varieties found in the nose, and are most commonly situated in the lower third of the chamber. Cysts, fibroma, fibromyxoma, chondroma, osteoma, epithelioma, and angiosarcoma are seen. Sarcoma usually affects the nares and ethmoid cells. Carcinoma is rare. Rhinoliths are sometimes discovered measuring an inch or more

<sup>1</sup> Based on ALLCHIN'S *Manual of Medicine*, 1902.



THE TRACHEA AND BRONCHI.—*Malformations*.—Fistula is due to imperfect closure of the third and fourth branchial clefts, a small orifice remaining, which may communicate with the trachea or end blindly, on the anterior edge of the sternomastoid muscle three or four centimetres above the inner end of the clavicle. The inner portion is at times dilated, forming a bronchiogenic cyst.

*Asthma*.—This condition being due to a spasmodic contraction of the bronchial tubes and air-vesicles, the lesions found at the post-mortem are not marked, consisting only in hypertrophy and widening of the bronchial tubes with thickened mucous membrane and a dilatation of the air-cells, giving the chest a barrel-shaped appearance and the dorsal spine a curvature. Pressure on the vagus by enlarged bronchial glands is sometimes the only lesion found, or it may be associated with hypertrophic rhinitis. Charcot-Leyden crystals and Curschmann's spirals are often discovered in the sputum.

*Bronchiectasis*.—There is a local or general dilatation of the bronchial tubes, which may be congenital and unilateral or the result of various diseases of the lungs and bronchi. A non-patulous bronchus, closed alveoli (atelectatic bronchiectasis), puckering of the peribronchial or interstitial fibrous tissue, parenchymatous changes of chronic bronchitis, or the circumscribed narrowing of tumors, etc., sometimes produce it. Cylindrical or uniform, saccular, spherical, ovoid, fusiform, and moniliform dilatations are seen. At the postmortem large sacs may be situated immediately beneath the pleura or a number of sacculi varying in size may be found opening one into another. The walls are covered with smooth, glistening, or hypertrophied epithelium, which may be ulcerated in the dependent portion. Putrefaction (putrid bronchitis), fatal gangrene, or a tuberculous lesion may follow the retention of the material thus collected. Calcification sometimes occurs. The lungs usually show some fibroid change. The air-vesicles are emphysematous or condensed by pressure. The liver, spleen, and kidney often show chronic congestion or lardaceous degeneration. Pleuritic abscesses, peritonitis, adhesive pericarditis, dilatation of the right heart, etc., are complications which are sometimes found to be present in bronchiectasis.

*Bronchitis*.—Bronchitis is an acute, subacute, or chronic inflammation of the bronchial tubes, not involving the terminal bronchi, due to infection, exposure, irritants, or extension from neighboring organs. In *acute catarrhal bronchitis* the mucous membrane is thickened,

swollen, at times hemorrhagic, and at first covered with tenacious mucus, which later becomes profuse, thin, and purulent, and may fill the large bronchi. Lobular atelectasis surrounds the affected areas. *Suppurative bronchitis* results from septic embolism, forming small abscesses in the bronchial tubes. *Croupous bronchitis* is characterized by the formation of a diphtheritic membrane, sometimes of complete casts of the smaller bronchi, but is not, as a rule, associated with the Klebs-Löffler bacillus. Bronchopneumonia, extension to the small bronchioles and air-vesicles, œdema, congestion, and local emphysema are frequent complications. In *chronic hypertrophic bronchitis* the whole lung is larger, firmer, and darker than normal. The mucous membrane is thickened, reddish or slate-gray in color, and often shows petechial hemorrhages. The bronchus is sometimes dilated, or is thicker and more fibrous, with its lumen narrowed by small, firm, villous granulations. This obliterating fibrous inflammation usually affects the smaller bronchi and may cause stenosis. The surrounding lung is usually emphysematous, pigmented, and shows an increase of fibrous tissue. The bronchial glands are enlarged, indurated, and pigmented. Dilatation of the right heart and chronic congestion of the liver, spleen, and kidneys are common complications. The lung of *atrophic bronchitis* is smaller and lighter in weight and color. Its elasticity is impaired, it feels "cottony" to the touch, and there may be increase of connective tissue, the longitudinal bands of elastic tissue standing out prominently. The mucous membrane is smooth, atrophied, and the lumen of the tubules may be widened. *Putrid bronchitis* is practically a bronchiectasis. The bronchi are dilated; their walls are usually smooth, but are frequently ulcerated. Fatty plugs and purulent masses such as are found in the sputum during life are seen in the ulcers. Purulent œdema of the lung is more or less general. *Plastic (fibrous) bronchitis* may be regarded as a chronic form of croupous bronchitis, occurs only rarely, and is paroxysmal, usually being limited to a certain number of bronchi. The membrane is a fibrous, fairly consistent pseudomembrane about two millimetres thick with no epithelium under it. The mucous surface is hyperæmic and infiltrated with cells. The thick ducts of the glands push the fibrous tissue off and it is coughed up. The smaller bronchi show catarrhal inflammation, but no membrane. Sometimes coagula are found in the tubes after death. *Tuberculous bronchitis* may be acute, manifesting itself as a part of a diffuse caseous process and involving an entire wall (peribronchitis)

or appear as tuberculous ulcerations like those of the larynx. *Cheesy bronchitis* is a caseation of retained catarrhal secretion. The mucous membrane is infiltrated with cells and these also subsequently caseate. *Gangrenous bronchitis* is associated with bronchiectasis.

*Stenosis of the trachea or bronchi* is due to pressure from tumors, aneurisms, mediastinal abscesses, hemorrhages, swelling of the lining mucous membrane, impacted foreign bodies, etc., to perforating caseous glands or to contraction from syphilitic or tuberculous ulcers.

*Primary tumors of the bronchi* are rare. Carcinoma may develop from the muciparous glands of the bronchial mucous membrane. Secondary tumors are more common. Calcareous, papillomatous excrescences, annular in shape, may be found in the trachea. *Foreign bodies* in the air-passages are discovered usually in the right bronchus. If the blocking is partial, vesicular or interstitial emphysema results. Œdema, local inflammation, and ulceration of the bronchus, lung, and pleura, with rupture of vessels, may occur.

DISEASES OF THE MEDIASTINUM.—*Mediastinitis* is rare, and when present is generally due to infection by *Pneumococci*. The cellular tissue is infiltrated with a puriform lymph. Pericarditis, pleurisy, abscess, gangrene, ascites, œdema of the upper part of the body, and albuminuria may follow mediastinitis. *Mediastinal adenitis* may be simple, suppurative, or tuberculous. The simple form occurs with any inflammation of the neighboring organs. Suppurating glands may rupture into the œsophagus, bronchus, or aorta. Tuberculosis of glands is usually a secondary involvement from the bones, lungs, or pleura. The *spindle-celled sarcoma* is the most common growth in this region. Carcinoma, lymphoma, and lymphosarcoma are also found. Dermoid cysts have been reported as occurring in the anterior mediastinum.

GOITRE.—A local or general hypertrophy of the thyroid gland, characterized pathologically by a variety of morbid changes. In the same gland may be found cystic disease and mucoid, fatty, gelatinous, or colloid degenerations. On section the gland appears as a yellow or brownish mass with scattered areas of colloid matter, varying in size from a pin-head to a millet-seed. In cystic goitre there is a distinct limiting membrane, brownish red if the cyst be due to hemorrhage. If slender masses of tissue project from this membrane, the condition is known as papillary cystadenoma. In some cases the enlargement of the gland may be due to marked vascular dilatation without the



formation of new gland-tissue. If the arteries only are dilated, Orth calls it *struma aneurysmatica*; if the veins only, *struma varicosa*. Fibroid or calcareous changes also occur. In fetal adenoma the structure maintains its fetal characteristics,—i.e., solid masses or rosettes of epithelial cells with little or no colloid material are seen.

*Goitre, Exophthalmic (Basedow's or Graves's Disease).*—A disease common to women of early adult or middle life, which is characterized by functional disturbance of the heart (diffuse or unilateral), hypertrophy of the thyroid gland, rarely as great as in ordinary goitre, a marked increase in the number and size of its blood-vessels, absorption of its colloid material, a replacement of it by a more mucinous fluid, and undue prominence of one or both eyes, due to an increase of the orbital fat. The thymus gland may persist and undergo enlargement, and there is an increased amount of connective tissue in the neck. Marked pigmentation of the skin may simulate Addison's disease. Myxœdema may develop in the later stages or the emaciation may be extreme. Glycosuria and albuminuria are not infrequent. The heart is usually hypertrophied, but may be dilated or even normal in appearance.

*LUNGS.—Abscesses.*—Abscesses of the lung and neighboring parts may arise from pyæmia, embolism, tuberculosis, pneumonia, or the presence of foreign bodies, and are due to many varieties of bacteria, as pneumococci, tubercle bacilli, gonococci, actinomycetes, and various pyogenic micro-organisms. An abscess may discharge itself through a bronchus or otherwise and leave only a dense cicatrix. The solitary abscess is comparatively rare, and usually results from disease of the neighboring parts, as the pleura, liver, or mediastinum. Such an abscess may become encapsulated and contain a greenish-yellow pus of an offensive odor. Multiple abscesses are common, generally superficial, frequently wedge-shaped, rarely encapsulated, and vary in size from that of a pea to an orange. They are at first firm, grayish red in color, and surrounded by a zone of hyperæmia. Later they become distinctly purulent, with an irregular, ragged cavity. The pleura is usually covered with a greenish lymph and may be perforated, causing an empyema, pyæmia, or septic pleuritis. Such abscesses often have their origin in a septic condition following criminal abortion.

*Atelectasis.*—Collapse of the lung, partial or total, may exist in the foetus at birth (fetal atelectasis), or be caused by closure of the bronchi (capillary bronchitis), compression from a tumor, hernia of the



diaphragm, pleuritic transudates and exudates, or marantic conditions. The last is due to weakness and is most marked in the smaller ramifications of the lower and posterior bronchi, often ending by subsequent œdema in pulmonary splenization. The air in the shut-off portion is absorbed, and the portion on section is dark red or bluish red and firm. In old cases the lungs cannot be inflated, the tissue is dense, firm, deeply lobulated, and paler than the rest of the lung. The lung is usually atrophied or may be entirely replaced by a fibrous cicatrix. When the collapse is superficial, as in rickets or pleurisy, the lung is reduced in bulk and wrinkled, fleshy in appearance, smooth, tough and inelastic, and dark red in color. If due to bronchial obstruction, scattered patches of atelectasis occur over the lung, the bronchi leading to these areas being filled with mucopurulent secretion. The surrounding lung may be œdematous or perhaps the seat of chronic pneumonia.

*Circulatory Disturbances.*—Anæmia, hemorrhage, infarcts, fat embolism, or even air embolism, of the lungs may be associated with a similar condition in the right heart. A number of fatal cases of pulmonary embolism have occurred after intramuscular injections of calomel for syphilis, caused by detachment of clots from the femoral, iliac, or uterine arteries, by hydatids, and by phlebolites. Fat embolism of the lung should always be thought of in cases of fractures or of extensive injuries to the subcutaneous fatty tissues or traumatic rupture of the liver. Hæmoptysis may occur from hemorrhagic infarcts, brown induration, tuberculosis, an aneurism rupturing in the trachea or bronchi, acute inflammations, purpura, scurvy, etc. It is interesting to note that a pulmonary hemorrhage in tuberculosis may be the beginning of an attack or precede a fatal termination. Pulmonary thrombosis may arise from embolus, engorgement of the capillaries, or disease of the pulmonary artery. It is not an infrequent complication in pneumonia and tumors, and often occurs in cases of atheroma of the pulmonary artery. Thrombosis of a pulmonary vein has been reported, usually the result of gangrene, pleurisy, or œdema. Hemorrhage into the air-cells and lung-tissue is due to thrombosis or aneurism of the pulmonary artery or to aspiration, as in gangrene and tuberculosis, or to the hemorrhagic diathesis. The extent of lung tissue involved differs very greatly. The lung is large, firm, dark, and heavy. On section there is extravasation of considerable amounts of more or less frothy fluid blood.

*Passive congestion* occurs where there is obstruction of the circulation, in chronic illness requiring the recumbent position, and in dis-

eases of the central nervous system. It is basic or hypostatic. (a) In mechanical congestion, if the condition has lasted some time, the lungs are voluminous, russet-brown in color, œdematous, and cut and tear with difficulty, giving rise to the so-called brown induration. On section they are of a maroon tinge, which on exposure to the air soon gives place to a vivid red. The alveolar capillaries are distended and tortuous, the fibrous tissue is increased, and hæmatoidin deposits are found in the epithelial cells. (b) In hypostatic congestion the bases of the lungs are deeply cyanosed and heavy and the posterior parts engorged with blood and serum. In some instances portions of the tissue will sink in water and on section exude a bloody serum. In prolonged coma the hypostatic congestion may be associated with patches of consolidation due to the aspiration of food into the air-passages. (c) Passive congestion occurring in cerebral apoplexy is most marked in or may even be confined to the paralyzed side.

Notwithstanding its ample collateral circulation, the lung is frequently the seat of small or large *infarcts*, especially of the hemorrhagic variety, usually situated peripherally and associated with brown induration. They may be single or double. Thrombus, rupture from over-distention, and infection of an embolus are the most common causes of this condition. When recent the infarcts are dark red, firm, resistant, and vary in size, sometimes occupying the greater part of a lobe. Sloughing and gangrene may follow. The pleura is congested and covered with exudate, the branches of the artery going to the lesion being filled with clotted blood. In old cases a pigmented scar may alone show the seat of a former infarct.

*Pulmonary edema*, which is a transudation of serum into the alveoli and their walls, may be general or confined to the bases of the lungs. The organ is bulky, heavy, and pale, and pits on pressure. In some cases there is a partial consolidation, the lung appearing gelatinous and containing less air than normal. On section it exudes a clear, frothy serum. The dependent parts may be red in color and boggy.

*Emphysema*.—The dilatation of the air-vesicles is due to some weakness of the lung structure, as a congenital absence of elastic tissue, atrophy of the diaphragm, etc., and a dilating force, usually expiration, as chronic cough, certain occupations, etc. It may follow senile changes or a traumatism. The thorax is barrel-shaped and increased in its anteroposterior diameter. The clavicles, the sternum, and the costal cartilages are prominent. The intercostal spaces are enlarged and the

sternal fossa is deep. The back is rounded and the curve of the spine increased. The neck appears to be shortened. Dilated veins may be seen along the line of the attachment of the diaphragm. On removing the sternum, the anterior mediastinum is found completely occupied by pulmonary tissue, the pericardial sac being entirely covered; the lungs are large, light in color or only slightly pigmented. They are inelastic, do not collapse, but pit readily on pressure. To the touch they are soft like feathers; expulsion of the air causes a crackling sound, and a paper-thin tissue remains. The edges are rounded and obtuse. Beneath the pleura, especially about the anterior margins and the inner surface of the lobe near the centre, enlarged air-vesicles of a delicate bladder-like appearance may be seen, varying in size from that of a pea to a hen's egg (bullous emphysema). Amyloid bodies are sometimes found loose or embedded in the walls. Local emphysema is common around old fibroid or tuberculous lesions, the dilatation affecting also the bronchi. In the atrophic form, really a senile atrophy, the lungs are small, pale, dry, and pigmented, pit on pressure, and collapse when the thorax is opened. The chief seats are at the edges and the apices. The mucous membrane of the large bronchi may be rough and thickened; bronchiectasis may be present, and the lungs irregular in shape. The right heart is dilated and hypertrophied; the pulmonary artery is enlarged and atheromatous. Emphysema may be vesicular, being confined within the dilated alveolar spaces, or it may be interstitial, the alveolar walls being broken. This is seen especially beneath the visceral pleura and may be produced *post mortem* by decomposition.

*Gangrene (Pneumomalacia).*—Gangrene may be circumscribed or diffuse, and affects usually the peripheral portions of the lower lobe rather than the central. It is by some supposed to be due to a specific bacillus. The gangrenous part is large, firm and solid or of a pulpy consistence, heavy, and of an ash-gray to greenish-black color. The outer tissues are intensely oedematous, next is an area of deep congestion, and then a cavity with shreddy, irregular walls containing a greenish fluid of a most offensive odor. The pleura may be inflamed and contain an abnormal amount of exudate, or it may be perforated, causing a pyopneumothorax. The gangrenous material gives rise to an intense bronchitis, the bronchial tubes being obstructed by a thin, highly offensive pus or by mucus containing fatty acids, tyrosin, and leucin. The elastic threads disintegrate later



than the remaining tissue, a fact of considerable diagnostic value. Embolic processes are common, abscesses of the various organs, especially the brain, being the result.

*Parasites.*—Certain parasites may infest the lungs. The *Aspergillus niger* and *fumigatus* are sometimes found in these organs, always associated with a pneumomycosis, and the *Mucor mucedo*, a yeast fungus, in cases of cancer. *Cysticercus cellulosæ*, strongylus, and *Distoma hæmatobium* are now and again found. Mackenzie<sup>1</sup> reports a case in a Japanese from Portland, Oregon, of parasitic hæmoptysis or infection with the *Distoma Westermanii*.

*Pneumonia.*—The chief forms of pneumonia are catarrhal (bronchopneumonia), chronic interstitial, and lobar (croupous and fibrinous). In cattle there is also found a very infectious variety known as pleuropneumonia. *Catarrhal pneumonia* is an acute or chronic inflammation of the lungs, involving both the bronchial tubes and air-vesicles, and due to extension of inflammation from neighboring parts, aspiration or inhalation of irritants, or micro-organisms,—e.g., *Diplococcus pneumoniae*, staphylococci, streptococci, the diphtheria bacillus, and the bacillus of pneumonia, or it may follow as a sequel to the infectious fevers. The lung is larger, heavier, and firmer than normal, and in my experience the lower lobe of the right lung is most frequently affected. On section the surface is somewhat dark red in color, distinctly mottled, and may drip blood. On palpation irregular nodular areas of gray hepatization can be felt, surrounded by crepitant tissue. The nodules, seldom larger than a hazel-nut, contain a central bronchiole surrounded by a grayish-red elevated area of consolidation and filled with tenacious purulent mucus which can be pressed out. Recent patches are red-brown in color, firm, smooth or finely granular, but later they are gray and soft. Minute hemorrhages are common near the affected areas and on the pleural surfaces. The pleura is bluish in color and rough. Emphysema is seen on the anterior and upper portions of the lung, especially within the inflamed areas. Fibroid changes seldom follow bronchopneumonia. Associated with this variety of pneumonia may be found enlarged bronchial glands, a dilated right heart, gastritis, enteritis, congestion of the liver and kidneys, and rarely pericarditis or pulmonary thrombosis. Catarrhal pneumonia in itself, except in the very young or the very old, is rarely fatal.

---

<sup>1</sup> *Jr. Amer. Med. Assoc.*, April 30, 1904, p. 1133.



*Chronic interstitial pneumonia* may be due to acute inflammations (rare), tuberculosis, chronic pleurisy, chronic poisoning, or syphilis, and is usually unilateral. The chest on the affected side is sunken, deformed, and the shoulder depressed, the heart being drawn over to the affected side. The opposite lung is usually emphysematous. On opening the chest the affected part, more or less deeply pigmented, is seen to be almost airless, quite firm, and very resistant to the knife, lying back against the spine, and usually held by dense adhesions. On section grayish fibroid tissue of variable amount is found, which may be more or less dilated. The unaffected lung is much enlarged, occupying the greater portion of the mediastinum. The heart is hypertrophied and the blood-vessels may be atheromatous. Associated lesions are tuberculosis or syphilis, a cavity of the apex, pulmonary aneurism, and amyloid disease of the viscera.

*Croupous (lobar) pneumonia*, an infectious and contagious disease, occurs especially in adult males, the *Diplococcus pneumoniae* of Fränkel being present in a large proportion of cases, though other organisms, such as the *Aspergillus bronchialis*, may be found. The organism, readily demonstrated in cover-glass preparations stained by Gram's method, is found in the bronchial secretions and in sections of the affected lung, in pairs, surrounded by a lanceolate capsule. Osler considers the mortality to be about one in four persons affected. The disease is divided into three distinct stages,—hyperæmia, red hepatization, and gray hepatization. (a) In the stage of engorgement, which lasts about twenty-four hours, the lung is heavier, firmer, more solid, and redder than normal. It still crepitates, though not so distinctly as the healthy tissue. The cut section exudes a red, frothy serum and will partially float. (b) In red hepatization, which lasts from one to four days, the affected lobe or lobes are larger, heavier, and firmer than normal, and are of a deep-red color. They are airless, do not collapse on exposure to the atmosphere, and excised portions sink in water. The pleural surface of the lung is covered with a more or less extensive layer of fibrin, which forms a false membrane that contrasts markedly with the smooth shiny appearance of the unaffected portions of the lung. The surface may retain the impressions of the ribs. On section the lung is dry, reddish brown, and exceedingly friable. Careful inspection shows that the surface is distinctly granular, due to fibrinous plugs in the smaller bronchi and blood-vessels, which are lighter in color than the intensely red

tissue, and which can be scraped off with a knife together with a reddish-viscid serum. Such fibrinous masses may extend into the larger tubes and thus form perfect casts. The bronchi may contain a mucous secretion tinged with blood, or more rarely the tenacious mucus so characteristic of pneumonic sputum. The microscope reveals in the alveolar meshes fibrinous threads, epithelial cells which have undergone hyaline and necrotic changes, leucocytes, red blood-cells, micro-organisms, etc. Sections taken from the central portion of the lung show more cellular elements, while those from the surface are richer in fibrin, showing that infection probably takes place from the bronchi. In this connection it is well to remember that there is a pneumonic form of plague and of several of the other infectious fevers. (c) In gray hepatization the color varies from a reddish brown to a grayish white. The surface is more moist, the exudate more turbid, the color grayish yellow or green. The granules are less distinct and the pulmonary tissue is still more friable. The exudate is softened and the pneumococcus is usually no longer to be demonstrated. The cell-elements are disintegrated and prepared for absorption. Gray and red hepatization may coexist in the same lobe. *Lesions in Other Organs.*—(d) The bronchial glands are swollen, soft, and hemorrhagic. The overlying pleura is inflamed, with more or less extensive exudate, which may be serous, fibrinous, or, more rarely, purulent. The cavities of the right heart are often distended with firm tenacious coagula. Pericarditis is not infrequent with pneumonia of the left side or double pneumonia, and is most common in children. Endocarditis is more common, and may be malignant and associated with meningitis, usually of the cortical variety. Myocarditis is rare. In many cases the spleen is enlarged. The kidneys and liver show cloudy swelling or acute parenchymatous changes. The hepatic veins are often engorged.

*Complications.*—Otitis media, conjunctivitis, and arthritis are not unusual in children. Severe and often fatal toxæmia may develop with a comparatively slight lesion in the lung. Jaundice, croupous gastritis, croupous colitis, and peritonitis also occur.

*Terminations.*—(a) Liquefaction, absorption, and resolution. (b) Suppuration. The lung is then an airless, firm, regular, gray or red mass. Abscesses should always be examined for tubercle bacilli. (c) Gangrene. (d) Fibroid changes or carnification. (e) Lymphangitis and perilymphangitis may occur.

TABLE SHOWING DIFFERENCES BETWEEN CROUPOUS AND CATARRHAL PNEUMONIA.

CROUPOUS PNEUMONIA.	CATARRHAL PNEUMONIA.
1. Whole lobe usually affected; hence the name lobar pneumonia.	1. Lobules affected; hence the name lobular pneumonia.
2. No areas of healthy lung tissue in affected lobe; other lobes healthy, but may be congested, especially those near the affected lobe.	2. Irregular areas of lung tissue in various stages of degeneration intermingled with normal lobules.
3. Lung weighs much more than normal. An entire lobe may sink in water.	3. Lung weighs but slightly more than normal. An entire lobe will float on water, though small portions may sink.
4. Microscopic appearance varies according to stage. Much fibrin; hence the name fibrinous pneumonia for this condition.	4. Microscope reveals three zones: central, a small bronchus; middle, a desquamative area containing many cells, but little or no fibrin; outer, a zone of congestion. Hence, the synonym, bronchopneumonia.
5. An extensive fibrinous exudate on the pleura covering the affected area ("bread-and-butter" pleurisy); hence the name pleuropneumonia for this affection.	5. Exudate slight, if present.
6. Pneumococcus usually found.	6. Pneumococcus rarely found.
7. Usually at base and posteriorly.	7. Usually at the termination of the smaller bronchioles and anywhere in the lung.
8. Usually one-sided.	8. Usually on both sides and associated with other diseases.
9. On section the lung varies according to stage, the marbled appearance being especially striking in the third stage. Notice the fibrinous plugs.	9. On section the lung is congested. Small angular irregular patches, the central portion being the oldest, are seen.
10. Sputum, so-called rusty sputum.	10. Sputum more purulent.
11. Lung lesions of same age.	11. Diseased portion of the lung varies; some spots are old, some are new, the oldest being around the bronchioles; healthy tissue between affected areas. Caseous pneumonia, really a form of catarrhal pneumonia, is due to the action of a toxin, as from the tubercle bacilli. In phthisis there may be small areas of croupous pneumonia.
	12. Capillary bronchitis and catarrhal pneumonia are, pathologically, practically the same.



*Pneumoconiosis.*—This fibroid condition of the lung, often associated with tuberculosis, and produced by the inhalation of particles of mineral or metallic substances, occurs in persons employed in such occupations as coal-mining, the manufacture of pottery, steel-grinding, stone-cutting, tobacco-sorting, etc. Various names are thus applied to it, depending upon the nature of the inspired dust,—*e.g.*, anthracosis, siderosis, calcicosis, lithosis, silicosis, etc. Unless, as is frequently the case, emphysema coexists, the affected lungs are harder, firmer, often smaller than normal, and usually of a blue-black, yellowish, or buff color, affording a striking contrast to the lung of a child. Even when the inspired dust is white, the lungs are apt to be of a dark color, due to the carbon and the altered blood pigment. In advanced stages of anthracosis an ink-like juice may exude from the cut surface. In siderosis, caused by oxide of iron, the lung is of a reddish color.\* On section condensed portions of highly fibroid tissue are seen, with numerous raised points, which give it a coarse granular appearance. These raised points are small, thickened, fibroid bronchial tubes protruding above the surface. The deposits are found microscopically everywhere along the course of the lymphatics. The pleura is usually adherent, thickened, and pigmented. The signs of chronic bronchitis are present, though the mucous membrane of the bronchi remains unpigmented. The bronchial and peribronchial glands as well as the peribronchial lymph-nodules are frequently intensely pigmented, and may be either soft or indurated. The liver and spleen may also be pigmented. True osseous formations, coral-like in shape, may be found in the lungs.

*Traumatism.*—In accidents foreign bodies may enter the lungs. In one case reported<sup>1</sup> a woman fell from a ladder and a broom-handle passed through her chest from one axilla to the other; she recovered.

*Tumors.*—The benign tumors of the lungs are fibroma, adenoma, osteoma, and chondroma. Hydatids are common in countries infested by that disease, and may attain considerable size. Dermoid cysts are found, but very rarely. Primary malignant growths are rare, involving one lung only, while secondary tumors are comparatively common, affecting both lungs. Carcinoma may originate in the epithelium of the alveoli, the bronchi, or the mucous glands. Secondary cancer is more frequent in women than in men and may

<sup>1</sup> FRANKE, *Arch. f. klin. Chirurgie*, 1903, vol. lxxi, no. 2, p. 543.



be scirrhus, encephaloid, epitheliomatous, or colloid. Endothelioma starts from the lymphatic apparatus. Primary spindle-celled sarcoma and melanosarcoma are found, which often extend to the liver. The tracheal or bronchial glands are sometimes the seat of metastatic growths. In malignant diseases of the lungs, pleurisy, generally of a hemorrhagic type, is commonly present.

**PLEURA.**—*Empyema (Pyothorax).*—Suppuration in the pleural cavity is usually accompanied by the presence of air, and is due to pleurisy, extension of inflammation from neighboring organs, trauma, and micro-organisms, especially the tubercle bacillus, the diplococcus of pneumonia, and the streptococcus and staphylococcus pyogenes. The pleuræ are much thickened; their surfaces are irregular and covered with a yellowish-green exudate of varying thickness. There may be evidences of more or less extensive hemorrhage, also erosions, fistulæ, or perforations. In severe cases there may even be gangrene. The pus separates into two layers,—a clear greenish-yellow serum above, a thick cream-like pus below. It has a heavy sweet odor, and is rarely fetid unless gangrene supervenes. A sterile culture on ordinary media suggests tuberculosis.

*New Growths.*—The benign tumors of the pleura are fibroma, osteoma, chondroma, and lipoma. Endothelioma originating from the lymphatics may cause a diffuse thickening of the pleura. Teratoma has been reported, in one case attached to the tenth rib, aorta, and vena cava. Carcinoma and sarcoma occur, usually as secondary deposits. Hydatid cysts are very rare in this country. Tuberculosis of the pleura is described elsewhere.

*Pneumothorax.*—This is the presence of air in the pleural cavity, and may be due to traumatism, tuberculosis of the lung rupturing into the pleura, other infectious granulomata, and malignant growths. The thorax is usually distended and the intercostal spaces may be obliterated. The introduction of a trocar allows the escape of the air. Unless pneumothorax kills suddenly, it is always accompanied by a pleurisy, generally of a purulent variety. The pericardium and heart are pushed or drawn to the opposite side. The lung is usually compressed and carnified and may be adherent to the chest wall at the apex, this site being frequently the seat of caseous nodules or cavities. Localized pneumothorax is probably often overlooked at the postmortem.

*Hydrothorax.*—This is part of a general dropsy, and is usually

due to chronic valvular disease, chronic Bright's disease, cirrhosis of the liver, cachexia, or pressure on the azygos veins. As a rule, it is bilateral, although not equal in extent in the two sides. The lungs are compressed and the pericardium and heart pushed upward.

*Hæmothorax*.—This is generally due to trauma, sometimes to cancer of the lung or pleura, also found with tuberculosis, purpura, scurvy, leukæmia, cirrhosis of the liver, and granular kidneys.

*Chylothorax*.—An effusion of chylous fluid is rare, and is due to traumatic rupture or obstruction to the thoracic duct.

*Pleurisy*.—Inflammation of the pleura may be acute or chronic. It is due to exposure to cold and wet, traumatism, extension of inflammation from neighboring organs, pyogenic micro-organisms, many infectious fevers, infectious granulomata, or malignant tumors. *Acute pleurisy* is classified by the character of its exudate into serous, serofibrinous, fibrinous, purulent, and hemorrhagic. In all, the serous membrane is at first red, sticky, and lustreless, and the vessels are dilated; later it becomes pale, thick, and rough. The pleural cavity may contain an inflammatory exudate, varying in amount from a few cubic centimetres to one or more litres, resembling that seen in other serous cavities. The serofibrinous exudate contains more fibrin but less fluid. Coagula may be found *in situ*. The characteristic of the fibrinous exudate is the so-called "bread-and-butter" appearance of the pleura. The deposit varies in thickness from a millimetre to a centimetre or more. *Purulent pleurisy* may follow the acute form or may be primary. It is frequently associated with tuberculosis. The serous membranes are covered with a creamy exudate and the cavity contains from a few cubic centimetres to a litre or more of greenish-yellow, offensive pus. In the tuberculous exudate Ravaut and Vidal have found a predominance of lymphocytes, while other effusions contain polymorphonuclear neutrophiles. *Hemorrhagic pleurisy* may be due to asthenic conditions, as tuberculosis and cancer, or may occur in perfectly healthy individuals, from wounds to the lungs during aspiration by the mixing of any fluid present with blood. The pleural cavity contains blood, usually fluid and varying considerably in density. The serous membranes are generally inflamed and stained with blood-coloring matter. *Chronic pleurisy* with effusion may persist for months without undergoing any alteration in its character. The post-mortem appearances are very similar to those of an acute pleurisy. *Chronic dry pleurisy*, resulting from the partial absorption of

a pleuritic exudate and the organization of the remainder, occurs usually at the base, causing marked flattening of the chest. Small pockets of fluid are often found and it is frequently impossible to separate the layers of pleura. The lung is compressed, airless, and fibroid. *Primitive dry pleurisy* may be limited or universal, unilateral or bilateral, and may be accompanied by a similar condition of the pericardium and peritoneum. The layers of the pleura are firmly adherent to one another and, especially about the lower lobe, are much thickened. In tuberculous cases reddish-gray fibroid masses and small tubercles are present between the layers, sometimes infiltrated with serum. The bronchi may present marked dilatations and the pulmonary tissue be more or less sclerosed. In diaphragmatic, encysted, and interlobar pleurisy the morbid anatomy is similar.

## CHAPTER X

### CRITICAL EXAMINATION OF THE ORGANS OF THE ABDOMINAL CAVITY

THE OMENTUM, MESENTERIES, AND PERITONEUM.—The superficial examination of the peritoneum having been made during the general inspection of the abdominal cavity, any thick regions are now felt with the index-finger and thumb and, should anything abnormal be found, such areas are at once incised and critically studied. A regular order should be chosen for the study of the peritoneum, say from above downward, so that nothing of importance shall escape the attention of the one making the autopsy.

The peritoneum covering the diaphragm may become inflamed as part of a general peritonitis, or show the presence of aberrant pulmonary tissue, cysts, filaria, actinomycosis, lipomata, fibromata, endothelioma, and secondary tumors. The *Distoma hepaticum* has been found embedded in the peritoneum of this region. Subdiaphragmatic (subphrenic) abscesses are not uncommon, especially on the right side, and may rupture into the pleural cavity or remain localized. In Körte's<sup>1</sup> sixty cases, which were operated on by himself, infection was found to originate most frequently from the vermiform appendix. Among the other causes giving rise to this condition may be mentioned: (a) perforation of a gastric or duodenal ulcer; (b) abscess of the liver and pancreas; and (c) diseases of the lower ribs, pleura, and mediastinum. Mandl has tabulated 179 cases from the literature on this subject.

A volvulus or hernial opening of the omentum may require its removal; although, as a rule, it is best to make examination of any abnormalities or pathologic lesions that appear and afterwards to remove the omentum along with the transverse colon. In thin subjects the separation of the omentum into its four layers, one anterior and three posterior, forms a striking picture, especially if studded with recent miliary tubercles or the wild strawberry-like nodules of a sarcoma. The hæmolymph-nodes in man and in sheep have recently been studied with special care by Warthin. He thinks that they have to do with the destruction of the red blood-corpuscles.

<sup>1</sup> GRÜNEISEN, *Archiv f. klin. Chirurgie*, 1903, vol. lxx, p. 1.



The mesentery may be shortened by contraction, as by granulation of the tissue, or lengthened, as by traction upon the bowel in a strangulated hernia. Congenital redundancy of the sigmoid flexure may later cause chronic obstipation and hypertrophy of the colon. Hemorrhage may take place into the mesentery in phosphorus poisoning and acute yellow atrophy of the liver. The glands are red and swollen in enteritis, especially in typhoid fever, where they may be very numerous and break down. They afford a favorable spot from which to secure cultures for the different varieties of colon and typhoid bacilli. When the glands become tuberculous, they often caseate and may reach a large size. In the *tabes mesenterica* of children they are usually enlarged, even in non-tuberculous cases. All statistics bearing upon tuberculous infection of these glands are extremely useful at the present time, in order to place upon a sound scientific basis the relation of tuberculous milk to infant mortality. One also finds here enlarged glands in leukæmia and Hodgkin's disease. By the stopping up of the blood-vessels, the mesentery may become dark in color and cause many feet of the small intestine to become gangrenous. It may be wholly converted into a mass of fat. Search should be made for calcified tubercles, tumors, parasites, chylocysts, etc. Hemorrhagic infarcts are sometimes seen. A recent postmortem of infiltrated blood into the mesentery from a ruptured aneurism of the superior mesenteric artery showed the distribution of the blood to the mesenteries of the jejunum, ileum, cæcum, vermiform appendix, colons, and rectum. It is well to remember that the duodenum is not supplied with a mesentery.

Cancer of the peritoneum is found especially in the female sex after the change of life, and is most often secondary to cancer of the stomach or ovaries. It is spoken of as "miliary carcinosis" because the nodules are small, spherical, and diffuse. The serous membranes are pale, thickened, with marked fibrinous deposits, which form adhesions to neighboring viscera; the omentum is indurated, and forms a mass transversely across the abdomen; the bowels are often firmly matted together. Ascites is usually found; the amount of fluid present may be several pints or only a few ounces. In some cases of colloid cancer the masses are of large size.

Among the micro-organisms capable of demonstration in acute general peritonitis may be mentioned the *Streptococcus pyogenes*, *Bacillus coli com* aureus, *Streptococcus lanceolatus*,

*Bacillus proteus*, *Bacillus pyocyaneus*, and, more rarely, the gonococcus (in the female) and the anthrax and typhoid bacilli. The cause of the peritonitis is usually from a perforation of the bowel. It may be: (a) Serous. (b) Serofibrinous. (c) Fibrinous. (d) Purulent. (e) Putrid. (f) Hemorrhagic. (g) Ulcerative. In acute general peritonitis the peritoneum has lost its lustre, is opaque, and is covered with an exudate varying with the type of the disease. The intestinal coils are distended and glued together with lymph. They are more or less displaced and compressed, and their walls are easily torn. The serous membrane may easily be separated from the muscular coat. In peritonitis due to perforation, the peritoneum and its contents are discolored by the faeces, while the peritoneal cavity contains gas, which escapes with a hissing noise when an opening is first made in the abdominal cavity.

Causes of chronic peritonitis: (a) Follows acute. (b) Tuberculosis. (c) Extension of inflammation from the abdominal organs. (d) Cancer. *Classification*.—(a) Local adhesive. (b) Diffuse adhesive. (c) Proliferative. (d) Hemorrhagic. (1) Localized peritonitis occurs about the spleen, diaphragm, liver, intestines, mesentery, and pelvic organs. Bands of connective tissue more or less firmly organized bind the various organs together, producing marked alterations in the appearance and position of the parts. The peritoneum is thickened and puckered. (2) Diffuse adhesive peritonitis follows acute inflammation, either of a simple or tuberculous nature. The abdominal cavity is often obliterated; the intestinal coils are firmly matted together by the plastic exudate, which eventually becomes converted into bands of fibrous tissue. The spleen and liver are usually involved in the adhesions. In this variety I have seen a central cavity produced which contained the entrance and exit of several coils of the small intestines, the functions of life having apparently been carried on for a long while. (3) In the proliferative form there is great thickening of the peritoneum, which is opaque and white in color. The omentum is usually rolled into a thick mass between the stomach and the colon. The liver and spleen are the subjects of a chronic capsular inflammation; both are usually smaller in size, with thickened, wrinkled capsules. There are seldom many adhesions, and serous effusion may be present in the abdominal cavity. The intestinal wall is greatly thickened and the mucous membrane of the ileum is thrown into folds. Nodular thickenings may be present

and be mistaken for tubercles. (4) The hemorrhagic form occurs particularly in cancerous and tuberculous conditions. Layers of new connective tissue form on the surface of the peritoneum; they contain large blood-vessels, from which the bleeding occurs. It is commonly a circumscribed process. Orth compares it to chronic internal hemorrhagic pachymeningitis.

In order to obtain more room for the examination of the abdominal cavity and a more favorable opportunity for the subsequent inspection of the gall-bladder, biliary ducts, and portal vessels, the attachments of the diaphragm to the ribs on the right side may now be severed with the knife and the liver rolled over into the thoracic cavity of this side.

**THE SPLEEN.**—The spleen varies greatly in size and weight, even during health and in the same individual at different times. Its normal weight is about five ounces and the measurements are five by one and one-fourth inches. I have removed a spleen which weighed only one hundred and eighty-six and one-half grains (senile atrophy) and another weighing over seven pounds (malarial enlargement.) Enlargement of the spleen is also seen in sepsis, typhus, syphilis, etc. Remember that the spleen affords a favorable opportunity for the study of micro-organisms, especially of the typhoid and colon groups. The spleen may now be removed from the abdominal cavity, although some pathologists recommend its excision later in connection with the pancreas. It is easily found by passing the hand along the left under surface of the diaphragm from the eighth to the eleventh rib, well towards the side and beneath the cardiac end of the stomach. Usually but little force is necessary to bring it into view, with the gastrosplenic omentum and splenic artery and vein still intact. These parts are then cut or torn with a sort of twisting movement. In some cases the spleen is so soft that lacerations may be made in its substance by the fingers. These should not be mistaken for traumatic rupture of the organ, as from a kick, or for the rupture that sometimes, although rarely, results from disease. Occasionally the spleen is absent, its place being taken by a large number of supernumerary spleens or by an increase in the lymph-nodes of the peritoneum. The spleen may be found attached to the surrounding parts, or a wandering spleen may even be found in the left inguinal region. Before detaching it, examine the course of the splenic artery for aneurisms, supernumerary spleens, enlarged glands, etc. When this has been done,



the artery may be divided and the organ removed from the body. Notice whether or not the capsule is normal or thickened; it should be thin, smooth, and transparent. At times the capsule, from which trabeculae extend into its pulp, appears as if melted tallow had been poured over the surface and allowed to dry.

Now lay the spleen, resting upon the hilum (posterior surface), on the table, fix it with the left fingers, and with one stroke incise it in its longest diameter. The spleen being turned, transverse incisions to those made upon the anterior surface may be made for further investigation. The color of the normal spleen is dark red, somewhat darker and of a bluish tinge in children; it may be brownish, from the presence of hæmosiderin; or yellow, as in jaundice (or in the new-born, due to bilirubin crystals); or streaked with blue, owing to the presence of melanin. Coal-dust may be found in the spleen, having probably entered the circulation through the peribronchial glands. Hyperplasia of the fibrous stroma in cases of chronic enlargement of the organ, as in malaria and leukæmia, may give to the spleen a grayish tinge.

The structure of the splenic tissue may then be examined, and the changes in the splenic pulp, the Malpighian bodies, and the connective-tissue trabeculae noted. The elastic tissue of the spleen may be destroyed, as in tuberculosis, or hypertrophied around the capillaries, as in leukæmia.<sup>1</sup> A disturbance of the local circulation may lead to various changes. Oligæmia is marked by the light-red or grayish-red color of the spleen, with wrinkling of the capsule and prominence of the trabeculae. Obstruction to the portal circulation causes congestion. Hyperæmia due to congestion is characterized by an enlarged, hard, dark-red splenic pulp, with smooth surface on section and thickening of the capsule, trabeculae, and vessel-walls. Infarcts of the spleen are common, and are usually wedge-shaped, with the apex towards the hilum. They vary in size from that of a pea to that of a cherry, and may at times include half of the spleen; they may become infected. Anæmic infarcts are of a cloudy-yellow color, while the less common hemorrhagic infarcts are very dark red, and later become yellowish red, and even whitish yellow as the coloring-matter of the blood disappears. Acute splenitis, resulting in the formation of pus, is not frequent. An acute proliferative splenitis, the

---

<sup>1</sup> FISCHER, *Virchow's Archiv*, 1904, vol. clxxv, no. 1, p. 69.



cause of the so-called splenic tumor, is characterized by enlargement of the spleen, with the capsule markedly on the stretch, and the pulp, on section, being of a vivid red, at first darkish and later somewhat lighter. The pulp is soft and exudes on section, so as to conceal the Malpighian bodies. Fibrous, productive, or chronic inflammation of the spleen causes the chronic splenic tumor, recognized by the large size of the organ, which is hard, of a light or dark brownish hue, with thickened trabeculae, that may appear as streaks through the splenic substance. A leukæmic spleen with its umbilicated nodules is hard and of a reddish-gray color, sometimes weighing twenty pounds. Miliary tubercles, with caseation, and other tumors of the spleen occur. In the colored race miliary tubercles at times do not undergo caseation and may attain the largest size of any developing in the body. I have not infrequently seen them as large as wild cherries. The arteries in the splenic pulp rarely show macroscopic atheroma, although the tissue of the spleen may be infiltrated with the salts of lime. The most important of all the forms of retrocessive disturbances of nutrition of the spleen is amyloid degeneration. In this disease the spleen is firm and inelastic, so that the pressure of the finger leaves a decided mark. Amyloid degeneration of the pulp is characterized by the smooth, shining, almost transparent appearance of the cut surface, while the so-called sago spleen—the amyloid degeneration of the Malpighian bodies—is recognized by the enlargement of the lymph-nodules, which on section appear somewhat transparent and scattered over the cut surface. The amyloid reaction would be more frequently demonstrated if Lugol's solution were applied as a routine practice. A small piece of the spleen should also be tested for iron with ammonium sulphohydrate. The *Pentastomum denticulatum* and echinococcus cysts of the spleen are sometimes found, as well as multiple angiomas, cancer, and sarcoma, the latter occurring, in rare instances, primarily in this organ. On healing, gummata leave behind stellate scars of varying size.

THE INTESTINES, EXCEPT THE DUODENUM.—When the exudation in the peritoneal cavity is fibrinopurulent and has a fetid odor, its source should be sought in a perforation of the intestine, although it may have originated elsewhere, as from the uterus or adnexa. If the peritoneal fluid suggest perforation, the gut may be examined under water, as pressure on the intestine will then cause bubbles of gas to appear. The site of perforation is usually marked by an area of

fibrinous exudation, which may be so dense as to occlude the opening; or there may be several perforations, as in a case of typhoid fever. In duodenal ulcer the contents will be stained with bile. Erysipelas and poisoning by arsenic should be remembered as occasional, though rare, causes of intestinal ulcer. The exterior of the entire intestinal tract should be critically inspected, starting from below and going upward, and any adhesions should be very gently broken down, care being taken not to make an artificial opening in the bowel,—an accident quite apt to occur in certain diseased conditions. When, however, the intestines are extensively agglutinated, as in appendicitis, tuberculous peritonitis, etc., the parts may often be better studied by first carefully noting their relations and then removing them *en masse*. Observe whether there be distention or contraction of the bowels. Distention is marked in cases of stenosis or strangulated hernia, and when a large amount of faeces is contained within the intestines. Contraction is noted in enteritis and after starvation. Localized constrictions may be due to bands of peritoneal adhesions. A Meckel's diverticulum should not be overlooked, and its omphalomesenteric attachment going to the umbilicus should be searched for. The duct sometimes remains patulous until puberty, or even later. The lymph-follicles may be injected, and are noticeable as irregular whitish lines which, when pricked, exude a drop of milky fluid,—chyle.

A proper examination of the intestines can be made only after they have been removed from the body. For this purpose the intestine is doubly ligated in three places,—viz., (1) at the end of the duodenum and the commencement of the jejunum; (2) in the ileum, several feet above the ileocaecal valve; and (3) at the end of the sigmoid flexure and the commencement of the rectum. The method of doing this is as follows. A loop of string is carried by the nail of the index-finger (Fig. 90) through an opening in the mesentery made with an instrument or the fingers and the intestine is ligated. A second ligature, far enough from the first to allow of the gut being divided between them later, is then applied (Fig. 91, facing p. 111). Care should be taken that the ligatures be tightly held so as to prevent slipping, thus affording an opportunity for the escape of



FIG. 90.—Method of passing the string through an opening in the mesentery previous to tying it.

fecal matter. The large intestine is then cut between the ligatures, and its proximal extremity grasped and the mesentery severed by a sawing or fiddle-bow movement close to its intestinal attachment along the whole extent of the colon, until the ligatured spot in the ileum is reached, when the small intestine is incised between the ligatures. This portion is then removed to the sink or bucket preparatory to being cleansed (Fig. 92), and the remainder of the ileum and the jejunum are removed, as seen in Fig. 93, and also placed in the sink or bucket.

The intestines are opened either by pushing them into the open blades of the scissors or, better, by thrusting the enterotome or scissors through the bowels along the line of the mesenteric attachment (Fig. 94). Meckel's diverticula, like Peyer's patches, usually occur opposite the mesenteric attachment and, therefore, on the free anterior border of the ileum; hence the reason for opening the bowel at its mesenteric attachment. The ileocaecal opening is guarded by the two valves of Bauhin, the superior one normally covering the inferior. Each valve is to be examined on its iliac and caecal surface. The ileocaecal cut is to be made between the two valves; this is readily done by dissecting the ileum down to the valves, and then, as is usual, pushing the enterotome through the opening in the direction in which the chyle passes,—*i.e.*, towards the caecum and between the two valves. The appendix is opened opposite its mesenteric attachment, contrary to the rule in the case of the intestines. The situation of the valve of Gerlach is very variable. If a competent dead-house assistant is at hand, the opening of the bowel may be intrusted to him, as it saves the operator's time and prevents his hands from becoming impregnated with the disagreeable fetid odor of the gut. The assistant is instructed to call attention at once to any abnormalities observed, and opens all the intestinal tract except the ileocaecal valve and the vermiform appendix. These are to be incised and examined by the pathologist himself. The small and large intestines, after being freed of their contents, should be measured and weighed apart, the ileocaecal valve counting with the large intestine. After washing, the entire bowel is arranged, mucous surface upward, upon the post-mortem table, so that the pathologist may at a glance examine the intestines throughout their entire extent. In cases in which haste is a matter of importance, the intestines need not be removed from the body, but at the end of the autopsy the region of the ileocaecal valve is opened as well as the sigmoid and rectum, and, if no lesions are discovered here, the remain-





FIG. 93.—Method of removing the small intestines. They are elevated and traction is made on the mesenteric, which is cut as close as possible to its attachment to the bowel.



FIG. 94.—Opening of the small intestine after its removal from the body, the incision being made with the enterotome along the mesenteric attachment.



FIG. 95.—Method of incising the kidney with its ureter still attached.






FIG. 98.—Method of removing the capsule of the kidney with the bare hands.



FIG. 99.—Method of removing the capsule of the kidney with the knife when gloves are employed.



FIG. 101.—Position in which the body is advantageously placed for examination of the rectovaginal region and for the performance of a postmortem *per vaginam* or *per rectum*. The body is drawn down towards the end of the table and the hips are elevated with a block. The thighs are then strongly flexed and each one held securely in place by a separate bandage fastened beneath the table.



FIGS. 101-108. Method of performing a post-mortem extirpation of bladder, uterus, and adnexa through the vagina and the restoration of the parts afterwards. An avoidal incision, Fig. 102, is made through the skin about an inch from the external genitalia; this is then enlarged and deepened until the pelvic cavity is reached, when with the hand the bladder is loosened and the parts desired brought through the opening. Figs. 103 and 104 show the parts *in situ*; Fig. 105, after their removal. In sewing up the parts afterwards to prevent leakage double thread is used. It is introduced at the upper or lower portion of the wound and the needle passed through the two threads so that when it is pulled tight there will be no chance of the thread slipping. (Fig. 106.) Figs. 107, 108, and 109 show method of sewing. (In this case there was a prolapse of the uterus, which is well seen in Figs. 107 and 108. The figures are numbered, starting at the top, from left to right.)



ing portion remains unopened, unless palpation or inspection in the preliminary examination of the abdominal cavity has led one to suspect a lesion in other situations. If a spigot is at hand, the open end of the intestine may be drawn over it, and the water allowed to run through until clean. Passage of water from a stop-cock to cleanse the intestines is not, however, always a proper procedure, as it may injure the mucous membrane, rupture a Peyer's patch in typhoid fever, loosen some of the intestinal contents, change their consistence by admixture with water, etc. The bucket method of opening and cleansing the intestines—a very useful one in private cases—is illustrated in Fig. 92. In warm weather these viscera are particularly liable to undergo rapid decomposition after exposure to the air.

The color of the normal mucous membrane of the intestine is light gray, varying according to the contents of the bowel and the amount of blood present. Congestion of the small capillaries causes a general redness, while injection of the larger vessels produces red streaks; the two conditions may occur together. The greater the distention of the bowel the more pale is the grayish shade of the mucous surface, and if the contents of the gut are bloody, the walls are dark red. This diffused color is to be distinguished from the redness due to hyperæmia, occurring in inflammations, congestions, etc., by the marked injection of the capillary blood-vessels which is seen in the latter case. Even the vessels of the submucosa are observed to be overfilled. Thickening of the walls as well as partial overgrowth of the mucous membrane, often in the form of small polyps, may be observed in many of the chronic inflammations of the intestines. Enlarged villi, individually made out with the naked eye, may be noted in some cases of inflammation. (Orth.)

The following tabulation will be found of use in distinguishing typhoid and tuberculous ulcers.

DIFFERENCES BETWEEN TYPHOID AND TUBERCULOUS ULCERS.<sup>1</sup>

TYPHOID ULCERS.	TUBERCULOUS ULCERS.
1. Direction often longitudinal, involving the Peyer's patches, which are larger in size; actual amount of surface involved greater.	1. Direction transverse (frequently). This distinction is not so characteristic as is sometimes held. The ulcers are smaller and may be very numerous.

---

<sup>1</sup> After WOODHEAD, *Practical Pathology*, 3d edition, p. 455.



## DIFFERENCES BETWEEN TYPHOID AND TUBERCULOUS ULCERS.

## TYPHOID ULCERS.

2. Edges undermined, ragged, and can be floated out on water; thin, vascular, and composed of mucosa and submucosa; red.
3. Floor smooth and vascular.
4. Peritoneal surface unaltered, except that it may be inflamed. No thickening and no gray or yellow patches.
5. Mesentery unaltered; glands enlarged, vascular, pink, and softened.
6. Perforation more common both by separation of slough and by direct extension of the ulcerative process. Small opening by which the fæces may escape. Peritonitis. Hemorrhage may occur during either of these processes.
7. Microscopically: A specific inflammation affecting the adenoid tissue; blood-vessels distended, and increased vascularity of the mucosa and the submucosa. Dense masses of small round cells—lymphoid cells and leucocytes—with some large multinucleated cells, the latter of which are derived directly from endothelioid cells. A line of demarcation is formed and abscess results, beginning in the solitary glands and other lymphoid tissue of the mucosa and submucosa. Widal test positive.
8. Extension takes place laterally or in depth.
9. Heals by granulation, the thin edges falling on to and uniting with the granulating floor of the ulcer.

## TUBERCULOUS ULCERS.

2. Edges not undermined; thick, prominent, nodulated, terraced, or sloping; pale or red; composed of tissue infiltrated with tuberculous nodules.
3. Floor nodular, irregular, thickened, vascular, with pale or yellow points or areas.
4. Peritoneum thickened; small yellow or gray points in the floor of the ulcer running along the lines of the lymphatics.
5. Mesentery thickened at its attachment to the bowel; glands enlarged, firm and gelatinous on section, or caseous.
6. Perforation, peritonitis, and hemorrhage are all rare.
7. Microscopically: A specific inflammatory affection of the adenoid tissue and the mucous membrane, ending in caseation and connective-tissue formation; vascularity of the mucosa and submucosa; increase of the connective-tissue cells and lymphoid cells; tubular nodules, typical or caseating. It begins in the mucous membrane, and, like the typhoid lesion, is due to direct contagion or infection. Widal test negative.
8. Extension usually takes places laterally.
9. Very rarely heals.

## DIFFERENCES BETWEEN TYPHOID AND TUBERCULOUS ULCERS.

TYPHOID ULCERS.	TUBERCULOUS ULCERS.
10. Leaves a smooth, often depressed, pale, anæmic, or pigmented cicatrix, covered by a layer of epithelium, but no gland tissue. Seldom breaks out afresh, relapses being due to the affection of adenoid patches previously little damaged.	10. Leaves a puckered cicatrix in which are gray or white nodules; often breaks out afresh.
11. Presence of typhoid bacilli, which are also found in the enlarged mesenteric glands and in the spleen.	11. Presence of tubercle bacilli easily demonstrated.
12. Spleen enlarged and soft.	12. Evidence of tuberculosis elsewhere, especially in the lungs.

Cases of paratyphoid fever explain the occasional failure of the Widal test. A most careful study of all typhoid cases should, therefore, be made where the Widal reaction was not obtainable during life. The anatomic findings in the cases of paratyphoid fever<sup>1</sup> which have come to autopsy are those of septicæmia with splenic swelling and at times ulcers which resemble those of dysentery and do not affect Peyer's patches.

Whether during life a rectal enema may, by reversed peristalsis, be carried to the stomach and then vomited is an interesting but debated question which I believe should be answered in the affirmative. It is very difficult by pressure to force liquid past the ileocæcal valve, but in relaxed conditions, as in cholera, this is perfectly possible. The problem is interesting as bearing on the possibility of a gastrocolic fistula and reversed agonal invaginations.

Mayo Robson<sup>2</sup> has reported a case of peptic ulcer which developed in the jejunum forty months after the performance of a gastro-enterostomy.

An abundance of fæces in the large intestine indicates constipation, which occurs in an extreme form in partakers of opium, where one may find scybalous masses lying in pouches in the transverse colon as hard and dry as if they had been retained there for many weeks or

<sup>1</sup> WELLS and SCOTT, *Journal of Infectious Diseases*, vol. i, January, 1904, p. 72.

<sup>2</sup> *Annals of Surgery*, August, 1904, p. 186.

even months. A similar condition is sometimes found in old persons subject to chronic constipation; the masses even become encrusted with salts of lime. Distention of the small intestine shows that considerable food was recently taken. When the lacteals are well dilated, some three and a half hours have elapsed since the taking of the food which has reached this portion of the intestine. Pavlof finds that psychical secretion of the intestinal juices varies markedly according to the character of the food ingested. When the fæces are light in color, an absence of bile is shown; when dark or light red, blood is probably present, although it must be remembered that medicines, such as hæmatoxylin, may give a similar appearance. When dark or black, the presence of iron or bismuth may be suspected; if yellow, the possible administration of rhubarb should be considered.

Gall-stones and worms may be found anywhere in the intestinal tract, but most frequently above the ileocæcal valve and in the lower rectum. In one of my cases I found, not far apart, two *Tania medio-canellata*, their heads being firmly attached to the mucous membrane beneath folds of the valvulæ conniventes at the end of the duodenum. A specimen of ascarides in the Wistar and Horner museum of Philadelphia shows where one of them had penetrated the bile-ducts. As these worms try to escape from the body after death, this may be an instance of post-mortem penetration. Seat-worms are found in the lower rectum. Packard removed *post mortem*, at the Pennsylvania Hospital, a specimen of *Tania nana*. The *Ankylostoma americana* has been seen several times in Philadelphia. Loeb and Smith have recently pointed out the presence of a substance inhibiting the coagulation of the blood in the *Ankylostoma caninum*. Of course any of the varieties of intestinal worms seen in man may be found here, but it is surprising how few cases are described in post-mortem notes of our hospitals. The foulest odors arise in icterus and dysentery, while in cholera the odor may be hardly perceptible. True intestinal sand may be found and is largely composed of the phosphate and carbonate of calcium. It is most often caused by a pure milk diet or one of milk and lime-water. False sand, composed of biliary and fecal concretions, is more common, being seen especially in the vermiform appendix. Salol, when taken medicinally, may cause the formation of crystalline enteroliths. In a case of Brossard,<sup>1</sup> one of a number of calculi weighed

---

<sup>1</sup> *Bull. gén. de thérap.*, 1897, vol. cxxvii, p. 363.

two grammes. The tumors of the intestines are myoma, fibroma, polyps, lipoma, adenoma, carcinoma, and sarcoma, the latter variety being of rare occurrence. (Plate IV.)

In hemorrhage of the bowel the bleeding may be localized or diffused. In the former variety petechial spots or ecchymoses are found on the mucous membrane. The mucous membrane surrounding the hemorrhages may be normal in appearance or show the results of active or passive congestion. In diffuse hemorrhages the blood is free in the bowel or may be extravasated into the mucous membrane. In the former case it is brownish black or black in color and usually semi-liquid or tarry. In the latter case the extravasated blood is in slate-colored or black patches.

The average length of the appendix is about three inches, although it may measure as much as six. There is a mesentery often reaching to the tip and containing fatty deposits. The appendix may be absent. Its usual direction is towards the brim of the pelvis, but it may point in any direction. Appendicitis is most common in males and in early adult life, and is favored by fecal concretions, but rarely by foreign bodies. Among the articles found in the appendix have been pins, fecal masses, calculi, worms, gall-stones, fish-bones, tip of a thermometer bulb, seeds and fruit-stones, as of grapes, cherries, prunes, etc. The theory has recently been advanced that influenza and syphilis are common causes of many cases of appendicitis. Metschnikoff thinks that the condition is often associated with worms of various sorts. The principal micro-organisms are the *Bacillus coli* (most common), *Streptococcus pyogenes*, *Staphylococcus pyogenes*, *B. tuberculosis*, *B. typhosus*, *B. influenzae*, *Proteus vulgaris*, *B. pyocyaneus*, *Actinomyces*, *B. pseudotetanus*, *Micrococcus tetragenus*, and *B. adematous maligni*. A mixed infection is usually present, upon an average three species being found in each case. The normal appendix is never sterile, while the diseased organ is sterile in ten per cent. of cases. (Lanz and Tavel.) Acute forms: catarrhal, follicular, suppurative, and gangrenous. Of the chronic: catarrhal, obliterative, and chronic infective. In acute forms the appendix is reddish brown, black, or greenish yellow in color. The mucous membrane is swollen, reddened, and presents hypertrophied follicles, ulcerations, or a false membrane. The whole appendix is thickened, the serous membrane red and lustreless. In the suppurative form the abscess may be small and limited to the appendix; when large the pus frequently invades the peritoneal cavity, the sac being



formed by peritoneum, fibrinous exudate, and fibrous adhesions. It should be remembered that in cases of appendicitis abscess formation may start outside of the appendix and there be no perforation. In severe cases following ulcerative or obliterative conditions the abscess-cavity may contain the whole or a portion of the appendix which has been sloughed off. The abscess-cavity may become limited and remain so and be subsequently absorbed, or it may later open into the general peritoneal cavity. Rarely it breaks through the skin. It may rupture into surrounding organs or structures, as the vagina, bladder, and rectum. The appendix may become invaginated into the cæcum and, by obstructing the blood-supply, become gangrenous, slough off, and be passed by the bowel.

Ulceration following typhoid is often seen, and perforation is not unknown. In obliterative appendicitis the entire tube is thickened, firm, and stiff; the peritoneal surface is smooth or injected, and may be adherent or free. It may become cystic, the contents being clear fluid or pus. The situation of the appendix varies greatly; rarely it may be found on the left side, as in transposition of the viscera, or it may be entirely absent. I have seen the tip of the appendix resting beneath a distended gall-bladder, entering into the formation of a *left* femoral hernia, or lying in the sigmoid flexure in a case of ileocæcal intussusception. On microscopic examination the lymph-follicles are numerous and close together, but as age advances they become separated and smaller. Late in life the appendix undergoes marked fibrous change, which must be distinguished from obliterative appendicitis. Primary cancer and sarcoma of the appendix have been found in a number of cases. Lafforgue<sup>1</sup> reports a double hydatid cyst of the appendix.

More people die from dysentery than from plague, cholera, and yellow fever. It occurs especially in warm climates and after eating improper food. I. *Acute*.—(a) Catarrhal. (b) Amœbic. (c) Gangrenous. II. *Chronic*. In the early stages the bacillus of Chantemesse<sup>2</sup> is found, and in the later stages, especially where abscess develops, the amœba coli is seen. The blood of patients affected with tropical dysentery has an agglutinative reaction with the bacillus of

---

<sup>1</sup> *Gaz. des Hôpit.*, January 12, 1904, p. 33.

<sup>2</sup> Commonly spoken of as the bacillus of Shiga, although described by CHANTEMESSE and VIDAL in 1888. *Presse méd.*, July 23, 1902. For the latest information on this subject, see *Diarrheal Diseases of Infancy*, vol. i of studies from the Rockefeller Institute for Medical Research.

dysentery. Summer diarrhoea of children has also recently been shown to be due to the same organism. All the lesions of dysentery have certain points of election for the starting of the inflammatory process,—viz., the large bowel, the flexures of the large bowel, and the course of the valvulæ conniventes. (1) *Acute Catarrhal Dysentery*.—The mucous membrane is enlarged, swollen, and covered with tenacious blood-stained mucus. The solitary follicles stand out prominently and in protracted cases often show necrotic or suppurative change. In some cases numerous ulcers appear throughout the large bowel. In children the picture is that of an acute follicular colitis. At first glance the mucous membrane seems to be universally congested; on closer examination it is found to be more or less streaky, with bright-red pin-point areas of intense congestion. The peritoneal surface is enlarged, lustreless, and sticky. (2) *Amœbic Dysentery*.—In this form the amœbæ, of which there are several kinds, both pathogenic and non-pathogenic, are almost always present. These are unicellular protoplasmic motile organisms, five or six times the size of a white blood-corpuscle. They contain a nucleus and one or more vacuoles. The characteristic lesion is an ulcer, which has a small external opening, with extensive undermined infiltrated edges. Sometimes these ulcers run together, forming deep sinuous tracts bridged over by apparently healthy mucous membrane. There is a progressive infiltration of the connective-tissue layers of the intestine, causing pressure upon the blood-vessels and subsequent necrotic changes in the overlying structures, so that the mucosa or the muscularis may be sloughed off *en masse* in certain parts of the bowel. In severe cases the whole of the intestine may be much thickened and riddled with ulcers, with only here and there islands of intact mucous membrane. More rarely these ulcers have but slightly undermined edges, the borders being more or less cleanly cut. In some cases there is a tendency to purulent formations. (3) *Gangrenous Dysentery*.—This form is characterized by the formation of a diphtheritic membrane, which is more or less irregularly distributed; it is at first yellowish-brown, in later stages becoming black or ashen-gray; in the latter case it appears as sloughs more or less easily detachable. There is thickening of all the coats of the intestine, with great interference with the blood-supply, so that in severe cases whole portions of the bowel may become gangrenous. (4) *Chronic Dysentery*.—In this form the anatomic changes are variable. Deeply pigmented ulcers are often present or there may be cicatrizations; again, no trace of ulceration

may appear, but the entire mucous membrane presents a rough, irregular, figured appearance, in places slate-gray or blackish in color. Certain parts of the mucosa are greatly thickened and the muscular coat is hypertrophied. In some cases the solitary follicles are enlarged and pigmented. At times the outlets of tubules of the glands are closed, thus forming "slime cysts" (Orth), varying in size from a pin-head to a pea. The condition is called chronic cystic enteritis. The calibre of the bowel may be reduced, but stricture is very rare. *Complications*.—(a) In all cases dysentery may be complicated by peritonitis, pleurisy, pericarditis, or pyæmic manifestations. (b) In amœbic dysentery the characteristic complication is the abscess of the liver, which is usually single and occupies the right lobe. It may be multiple, when it is apt to be distributed superficially in any or all of the lobes. It is a large solitary abscess, the wall of which is made up of broken-down, rough, shaggy liver-tissue, without any of the ordinary pyogenic membrane. The contents of this abscess vary. The outer portions are gelatinous and composed of broken-down liver-tissue, blood-pigments, pus-cells, amœbæ coli, etc. The interior is usually of an almost watery consistency, and of a brownish or reddish color. In some cases cultures made from these abscesses are sterile. In hot climates the amœbæ coli are almost always found on microscopic examination of old cases.

In colitis, or inflammation of the large bowel, consider: (a) Early life. (b) Hot weather. (c) Improper foods. (d) Certain micro-organisms of the colon group. (e) Poisons. (f) Some infectious diseases. *Classification*.—(a) Simple. (b) Membranous. (c) Ulcerative. (d) Chronic. (1) In *simple colitis* the mucous membrane is much thickened and reddened, the rugæ are prominent, and petechial hemorrhages are common. In ordinary inflammation the follicles are inflamed and œdematous and on section they appear like pearls. When there is a marked cell increase, they are white or gray and more prominent. These follicles may become confluent. (2) *Membranous colitis* is characterized by the formation of a more or less complete cast of the intestine, usually from one to six inches in length, but it may extend a distance of several feet. The membrane usually appears homogeneous, but may be distinctly laminated and show deposits of fecal matter between the layers. The end of the cast may be well defined, but often shades off into a transparent gelatinous material. Associated are swelling and œdema of the submucosa. The mucous membrane not involved is very much inflamed and there may be hemorrhagic infiltration. The intes-



tine may show that perforation has occurred and gangrene may sometimes supervene. (3) In *ulcerative colitis* the appearances vary greatly: the ulcers may be small and numerous or they may be large in size and few in number. They may be perfectly regular in outline, but are usually irregular, with slightly undermined edges. The floor of the ulcer generally shows a somewhat sloughing bowel. The ulcers may communicate by the separation of layers of the intestines. In long-standing cases they are often intensely congested and tend to become transverse. Sometimes the floor of the ulcer becomes so thin as to be pushed out and form pouches. In very acute cases the mucous membrane is much reddened, highly vascular, and the surface is soft. The peritoneal coat of the bowel may be normal in appearance, but is usually red, somewhat sticky, and shows many dilated blood-vessels. Small hemorrhages are common. (4) In *chronic colitis* the bowel is often much thickened in all its coats. It may be larger in diameter. It is firm, even leathery, to the touch. The mucous membrane is hypertrophied, often much pigmented, and shows many small hemorrhages. The follicles are swollen and have a slaty appearance. There may or may not be ulceration.

There are four forms of dilatation of the colon: (a) Distention from gas. (b) Distention due to some solid substance within the bowel. (c) Distention caused by an organic obstruction in front of the dilated bowel. (d) The so-called idiopathic dilatation.

Malignant disease of the colon is generally a cylindrical-celled epithelioma, usually confined at the start to a small area, where its contraction sets up an annular stricture.

THE KIDNEYS AND ADRENALS.—The spleen and intestines having been removed and the liver turned over into the thorax, the kidneys and adrenals yet remain behind the peritoneum, often deeply embedded in the perinephrial fat. Of course, in anomalous cases, in certain diseases and deformities (notably Pott's disease), and in floating kidney they may be considerably displaced. In any event it is best and simplest first to find the ureters as they descend on the psoas muscles and enter the pelvis. The exact situation of the ureters is as follows: Each ureter at first passes obliquely downward and inward to enter the cavity of the true pelvis and then curves forward and inward to reach the base of the bladder. In its whole course it lies close behind the peritoneum and is connected to neighboring parts by loose areolar tissue. Superiorly it rests upon the psoas muscle and is crossed very obliquely



from within outward by the spermatic vessels, which descend in front of it. The right ureter is close to the inferior vena cava. Lower down the ureter passes either over the common or the external iliac vessels, behind the termination of the ileum on the right side and the sigmoid flexure of the colon on the left. Descending into the pelvis, it enters the fold of the peritoneum forming the corresponding posterior false ligament of the bladder, and, reaching the side of the bladder near its base, runs downward and forward in contact with it, below the obliterated hypogastric artery, and in the male is crossed upon its inner side by the vas deferens, which passes down between the ureter and the bladder. In the female the ureters run along the sides of the cervix uteri and the upper part of the vagina before reaching the bladder. (Quain's Anatomy.)

Incise the peritoneum on the left side first, then on the right over and in the direction of the brim of the pelvis, and follow up each ureter, gently tearing away the loose connective tissue, but being careful not to disturb seriously the relationship of the kidney and adrenal and their vessels until they have been noted. If this method be adopted, there is no need of making an incision in the peritoneum directly over the kidney, as is recommended by most pathologists. A careful examination of the vessels entering and leaving the kidney is next made,—vein, artery, ureter, etc.,—all of them being subject to many anomalies. The left spermatic or utero-ovarian vein enters at right angles into the renal vein, which I have known to be followed out in mistake for the ureter. The organs may next be “shelled out” of their bed of cellular tissue and fat and the vessels severed, thus permitting their removal from the body. The adrenal<sup>1</sup> is then separated from the kidney, weighed, measured, and incised in its greatest plane. Should disease of the bladder or ureters be present, the kidneys may be removed from the body with the ureters attached. This is always better in those very common cases in which double ureters are found. One nick is then put in the left kidney at its upper or lower border, and the kidney and adrenal are removed, or the kidney may first be dissected. Another method of distinguishing the right kidney from its fellow is to make a uniform rule as to which ureter shall be left the longer, by several inches, on

---

<sup>1</sup> The right adrenal is more difficult to find than the left, and may be permitted to remain in the body until *after* the removal of the stomach, duodenum, and pancreas, but should be sought for *before* the removal of the liver. Testut, quoted by Gerrish, gives admirable illustrations of the situation of the adrenals.

the separation of the kidneys from the body. The kidney is then cleaned and weighed, and any peculiarities are noted.

To remove the kidney while the intestines are still in the body, first hold aside the left sigmoid flexure and pull away the fundus of the stomach and the tail of the pancreas. Then make an incision over the convex border of the kidney. Next separate it from the surrounding tissue and cut the kidney out along with the adrenal. The right kidney lies under the liver, and in removing this adrenal be careful not to cut the inferior vena cava. If you remove the ureters with it (Fig. 95), on the right side a long incision must be made through the peritoneum that goes from the abdominal wall to the cæcum and colon. (Orth.) Nauwerck recommends a more complicated method. He cuts the descending colon from the mesocolon first. His primary incision is vertical and between the hilum and the spinal column, a second one being made in the convex border of the kidney.



FIG. 96.—Method of opening the kidney. The organ is held in the left hand with its hilum downward, and an incision is made with a brain-knife along its upper convex border and more than half through the renal substance. It is then reversed (Fig. 97) and the incision continued until the gland is nearly, but not quite, divided. In this manner there is no danger of cutting the hand.

Holding the kidney longitudinally in the hand, the hilum towards the palm and the convexity upward, a clean brain-knife or large cartilage-knife is used to divide it through its middle parallel to its greatest surface. The knife must be so sharp that it will cut without tearing, and care should be taken not to extend the incision through to the hand (Fig. 96). The wisest precaution for this purpose is first to bisect the kidney only to its centre, then reverse the organ in the hand and com-

plete the incision by cutting outward (Fig. 97). The pyramids and the calices with their papillæ will now be completely exposed and the two halves held together by the tissues composing the pelvis. If it be desired to lay open the hilum or a hydronephrosis, scissors should be employed. Precipitates of urinary salts in the pelvis are often mistaken for pus. A microscopic examination, especially if acetic acid be added, will at once reveal the true nature of the fluid. Now examine the surface for cysts, stellate veins (veins of Verhagen), aberrant adrenals, miliary tubercles, tumors, etc. Large cysts can readily be



FIG. 97.—Method of opening the kidney in such a manner as not to injure the hands of the operator.

seen. When incising a cystic kidney, it should be remembered that the liquid therein is often under considerable pressure, and may squirt several feet when the cavity is opened, and thus injure the eyes or soil the clothing of the operator or of those present at the autopsy.

The capsule, which when normal is transparent, is next stripped off from one side (Figs. 98 and 99), and its condition noted as to whether or not it is thickened, adherent, or non-adherent. If adherent, see if any of the cortical substance is removed with it,—*i.e.*, whether the inner surface is smooth or rough. In those cases where the capsule is adherent, this portion of the kidney should be saved for microscopic study along with the renal tissue lying directly beneath. The normal color of the surface of the kidney after removal of the capsule is brownish red.



The relation existing between the lighter cortex and the darker medulla is determined by drawing a straight line from the apex of one of the largest central cones of a pyramid to the surface of the kidney. Normally this relation is as one (cortex) to three (medulla); it is, however, frequently altered and should always be noted. The cortical substance is increased in parenchymatous nephritis and decreased in chronic interstitial nephritis. Also study the color of the external and cut surfaces, the quantity of blood or fluid exuding and its character, and the consistence of the organ. Thus, in parenchymatous nephritis the color of the cortex is a grayish white or light yellow. In poisoning by hydrocyanic acid much blood exudes, and in chronic interstitial nephritis the nephritic tissue is dense and hard. Both anæmic and hemorrhagic infarcts occur. Scars are often found, and may be due to many different causes, as gummata, thromboses, infarcts, stones, former operations, etc. Tumors of the kidney, especially fibroids, are quite common. With arteriosclerosis and granular kidneys, suspect apoplexy, especially if there has been a clinical history of flushing of the face. As a routine practice in the examination of the kidney, the amyloid reaction should be tried. A thin slice about one inch square, including both cortex and medulla, is removed from the organ and placed in Lugol's solution (which is preferably diluted four or five times) for several minutes and then examined with a hand glass in a good light. In weighing the kidney the fat which accumulates—as found in old renal cases in the renal hilum—is weighed along with the organ and unless its presence is mentioned may give a false idea as to the real weight.

Where decapsulation as an operative therapeutic measure has been practised, also after the scraping of the hepatic peritoneum for ascites, the post-mortem examination should be very thorough, as any information concerning such cases is most important at the present time.

The adrenals are covered by the under surface of the diaphragm, although not usually attached to it, and above and lateral to the tips of the kidneys. It is embedded in the same kind of tissue as surrounds the kidney, which is of a fatty cellular nature, the difficulty of finding the gland being in proportion to the amount of this tissue, which differs, however, in its color unless stained with bile. After the gland is found, its dissection is best accomplished by a pair of scissors. It is sometimes intimately connected with the kidney or even with the liver by bands of fibrous tissue. Accessory adrenals (mostly found by

microscopic study of other parts) are found at times in the neighborhood of the main gland or even several inches away. They may be found in the kidney (hydronephroma) or even in the liver. The adrenal of one side may in rare cases be absent.

The adrenals are best removed attached to the kidneys, though, as already stated, the ablation of the right adrenal with the kidney is more difficult than that of its fellow, and for this reason it is often left in the body and examined at the time of the removal of the pancreas. The adrenals are very delicate, and care must be exercised lest they be injured in their excision. Normally the adrenals consist of three layers, which differ more or less in the young and the old. The outer or cortical layer is light yellow in adults and grayish red in children. This tissue somewhat resembles that found in the thyroid gland. It is composed of radiating follicles whose cells are undergoing fatty degeneration. It will be seen in the new-born that the adrenals are relatively of large size in comparison with the kidneys and when examined microscopically no fatty metamorphosis is discovered. The inner or medullary substance is composed of neuroglia and ganglionic cells connected with a rich vascular supply. The middle zone, or intermediary substance, is brown, owing to pigmentation of the follicles. The amount of intermediary substance is subject to considerable variation. (Langerhans.) Later in life there is a tendency for the central part to become separated from the intermediary portion,<sup>1</sup> and in atrophy of this organ, when it takes place unevenly (as it frequently does), nodes are left on the surface which are not infrequently mistaken for tubercles. These organs are subject to numerous pathologic changes and are hyperplastic in many varieties of congenital deformities in which other nerve-tissue is affected. There may be here hæmatoma, melanoma, cysts, hypertrophy, glioma, primary cancer, echinococcic cysts, hæmatoid degeneration, tuberculosis, purulent infiltration, infarcts, and internal proliferations. The recent discovery of the marked action of adrenalin would seem to show the presence of an internal secretion acting directly upon the vascular apparatus. It by no means follows that the adrenals will be found affected, either macroscopically or microscopically, in all cases of Addison's disease. Exquisite miliary tubercles are seen in the adrenals, and in advanced tuberculosis the caseating mass may reach the size of a walnut.

---

<sup>1</sup> Letulle considers the formation of a central cavity as usually due to traumatism in its removal or to post-mortem changes.

Addison's disease is most frequently seen in laborers between the ages of twenty and forty years. It may be due to: (a) Tuberculosis, simple atrophy, cirrhosis, hemorrhage, or tumors of the adrenals. (b) Inflammation or pressure of structures bordering the adrenals. (c) Changes in the semilunar ganglia and the sympathetic system. The adrenals are not infrequently tuberculous, and there is then a deficiency of the internal secretion of these organs. The brownish pigmentation (bronze disease) is most marked on the chest. The spleen may be enlarged, as may also the thymus, if the latter organ persists. The stomach and intestines may show hypertrophied lymphoid follicles. No specific blood-changes have been observed. One of the most marked cases of pigmentation of the abdomen which I ever saw was that of a girl who had undergone an operation for the removal of a large dermoid cyst of the ovary. It is possible that in this case the semilunar ganglia or the adrenals were affected by pressure or otherwise. In two cases of primary sarcoma of the adrenal, and in one of general tuberculosis with marked involvement by caseous tubercles of both adrenals, I observed no pigmentation of the skin at the time of the autopsy.

**THE SEMILUNAR GANGLIA.**—The semilunar ganglion or celiac plexus, which receives the great splanchnic nerve and the pneumogastric, is situated behind the stomach and in front of the crura of the diaphragm, by the side of the celiac axis and the root of the superior mesenteric artery, and close to the suprarenal body (Fig. 100). It may also be found by tracing the nerves from the adrenals to their entrance into the ganglion. The ganglia should be carefully studied microscopically in all cases in which lesions are suspected in the adrenals or in the sympathetic system. The color and vascularity as well as the condition of the surrounding connective tissue should be noted. In cholera and typhus fever the ganglia are hyperæmic and may show evidence of the occurrence of hemorrhage (Rokitansky).

**THE URETERS AND BLADDER.**—The ureters may be distended with urine, as from an impacted stone, from cancer of the uterus, or from overfilling of the bladder. They are often double, most frequently uniting in their middle third, more rarely in the structure of the bladder, but may enter this viscus by separate papillæ. The ureters being slit open throughout their entire extent, the appearance of the mucosa is described, taking into account the color and character of any catarrhal exudate, should it be present. Many microscopists teach methods of diagnosing the situation of a lesion in the urinary tract



from the shape of the epithelial cells. A most interesting experiment is to take at a postmortem scrapings from the pelvis of the kidney, the ureter, bladder, and urethra, examine them under the microscope, and determine whether or not such a diagnosis is possible. Hemorrhages, abscesses, papillary fibromata, the *Distoma hæmatobium*, calci-




FIG. 100.—The relations of the pancreas, kidney, ureter, adrenal, and solar plexus are shown, the liver having been turned upward and the intestines shoved over to the right.

fied bodies, etc., are found in the ureter. Miliary tubercles of the mucous membrane are seen, often of typical shape and large size.

In some three hundred consecutive autopsies performed in one year, I met with three cases in which the ureter had been tied during abdominal operations on the uterus and its adnexa. The right ureter seems to be ligatured oftener than the left. In pregnancy there may be considerable pressure hydronephrosis.

If it be desired to collect the urine for microscopic, chemic, or medicolegal examination, it should be drawn off into a sterilized vessel

with a new catheter. Should strychnine poisoning be suspected, place a live frog in the urine, and if strychnine is present in any amount the frog will show the typical strychnine convulsions. Unfortunately, however, in strychnine poisoning the quantity of urine secreted is often very small, and the alkaloid is not always present in the urine of those dying from its effects.

PELVIC ORGANS.—*Removal of the Female Genitalia.*—The parietal peritoneum is freed around the entire brim of the true pelvis by a cir-




FIG. 110.—Method of opening the uterus; the lines show the places for the incisions, one of which has already been started at the cervix. Letulle prefers to open the uterus posteriorly.

cular incision, starting and ending at the symphysis pubis and including the anterior portion of the sacrum. Orth begins the incision between the rectum and the sacrum, while Schottelius recommends the ending of the incision at the posterior superior spine of the ilium. The body is then placed in the position seen in Fig. 101, and the thighs are separated. An oval incision is next made, starting above the external genitalia, below the symphysis pubis, and ending behind the anus near the



coccyx (which may be examined at this time), passing to the outside of the labia on each side. Traction is then made upon the soft parts towards the median line and the incision deepened, keeping as close as possible to the pelvic bones and taking care that the knife or scissors cutting in the direction of the long axis of the body does not injure the rectum, bladder, or external genitalia. It is now possible to remove the external genitals, bladder, and rectum through the abdominal cavity, or the internal parts through the oval incision exteriorly (Figs. 102 to 109 inclusive). Whichever method is adopted, the muscles, fatty tissue,




FIG. 111.—The uterus has been incised in the manner called for in Fig. 110. The ovary and the tube are opened. The fimbriated extremity, the hydatid of Morgagni, and a corpus luteum are well shown in the illustration.

and fascia holding the parts in place are to be severed without injury to the tissues desired to be preserved; or an internal or external hysterectomy may be performed, if for any reason the external incisions should be avoided. If the ureters and kidneys have been left connected, they may be removed at the same time. The pelvic organs having been excised, they are placed on a board upon the table in the same relative position that they occupied while they were in the body. The bladder is then incised anteriorly with the scissors on the median line from the fundus to the urethra, which should be opened. In the male, the prostate should be carefully observed.<sup>1</sup> The rectum is slit up along its

<sup>1</sup> Many obducenters partially open the bladder while it is still attached to the body; indeed, the entire examination of the pelvic organs can be made with the parts *in situ*.



posterior wall, while an anterior incision is chosen through which to examine the uterus. When it is desirable to preserve the exterior of the bladder intact, the rectum may be dissected away and the womb incised posteriorly, or the bladder may be removed so as to permit of the uterus being opened up anteriorly. A transverse incision in the uterus from the entrance of one oviduct to that of the other will give an opportunity for a study of their uterine termini, which are sometimes rather difficult to find. Each ovary is completely bisected through its free surface, with the exception of enough tissue at the bottom to hold the two halves together (Figs. 110 and 111). The oviducts are now opened. Ch. Robin has pointed out that the normal mucous membrane of the oviducts secretes a creamy material which, without a microscopic examination, may be mistaken for pus. For the method of closing the external opening, see directions under Figs. 102 to 109 inclusive. The older the ovary the more it is cut up, irregular, and covered with cicatrices.

*Removal of the Male Organs of Generation.*—In the male the bladder is pressed downward well towards the rectum, and the tissues thus put on a stretch are incised close to the under portion of the symphysis pubis. A circular incision is then made anterior to the rectum and as close as possible to the parts to be removed (seminal vesicles, prostate, Cowper's gland, bulbus, etc.) without injuring them or buttonholing the skin. The soft tissues of the penis (cavernous and membranous portions of the urethra) are dissected away from the skin from *within* the pelvis, traction being made to bring these parts into the pelvic cavity as fast as those above are loosened. The corpora cavernosa and corpus spongiosum being now fully exposed, they are incised transversely near the attachment of the prepuce, just below the corona glandis and frænum. By pulling on the spermatic cords from above and pushing up the testicles from below, these organs are then removed together. To facilitate removal, a few cuts may first be made into the deep inguinal ring. The skin of the penis and scrotum is well stuffed with cotton, so that they may conform as nearly as possible to their original shape. A preliminary symphysiotomy may even be performed, or a V-shaped portion of bone taken from the symphysis pubis, or, if desired, after dissection of the testes and their appendages, they may be returned to their normal situations. Unless by an accidental perforation of the skin,—as the knife is working in the dark,—there need be no visible deformity, if this method be properly carried out.

The rectum and the bladder and its component parts may be left attached or they can be separated, as preferred. The testicles may also be removed and examined by dissecting beneath the skin *in front of* the symphysis pubis until their situation in the scrotum is reached. They are then pushed up with the hand from below. The tunica vaginalis and the spermatic duct and its vessels are then dissected out.

The technic of my external method of examining the testicles, urethra, spermatic cord, etc., without mutilating or disfiguring the external genitals, is as follows. The penis is grasped with the left hand and drawn upward and backward over the symphysis pubis in such a manner as to expose its under surface and the scrotum. With the thumb and forefinger of the same hand a fold of skin is taken up at the point where the integument of the penis merges into that of the scrotum. This fold, which should be in the line of the long axis of the penis, having been drawn taut, incision is made across it at right angles to the line of the penis. If this transverse incision be not carried too far, it will leave an oval gap about an inch and a half in diameter when the fold of skin is allowed to fall back. This will be quite large enough to permit the proper execution of the subsequent steps of the operation, and the wound, after being sewed up, is so small that it is entirely concealed by the penis when replaced in its normal pendent position. The finger is next introduced into the scrotum and swept around so as to break up the delicate areolar connective tissue that forms the septum scroti and unites the dartos with the testes; then by slightly dilating the external wound the testicles can be removed from the scrotal sac. Next the root of the penis is grasped from within, and the extremely loose bands of connective tissue that unite the body of the organ to the integument are broken up, still using only the finger. These connections having been severed, the body of the penis can be drawn from its cutaneous sheath as far as the point of union of the prepuce with the tissues at the cervix, so that now the testes and the penis, as far as the glans, are exposed without their cutaneous investment. In severing the body of the penis from the glans and the tissue included in the inverted sheath of skin, great care must be exercised not to "bottonhole" the delicate structure of the prepuce. This accident can be avoided by amputating the glans at a point one-fourth of an inch from the corona (which can be plainly seen and felt through the delicate skin covering it) and carrying the incision parallel to its plane. The direction of the



FIG. 112.—Author's method of examining testicles, epididymis, spermatic cord, etc., without disfigurement. The primary incision is made in the median raphe in such a manner as to be covered when the penis is returned to its normal situation.



FIG. 113.—Testicles shelled out of the scrotum through the opening made in Fig. 112.



FIG. 114.—Appearance of the male external genitalia preparatory to minute examination in the author's method of exposing them without disfigurement.






FIG. 115.—Method of examining the seminal vesicles, which are exposed by incisions at the places indicated by the lines above the seminal ducts. *a, a*, edge of severed portion of peritoneum; *b*, urinary bladder; *c, c*, seminal vesicles; *d, d*, spermatic ducts. (After Nauwerck.)




FIG. 116.—Relations of the gall-ducts and duodenum. The gall-bladder in this case was packed with stones and one large one was found in the common bile-duct; the pancreatic duct communicated with the duodenum by a separate outlet, and a probe is seen emerging from the opening through which the bile normally finds its way into the duodenum.




FIG. 117.—Examining the bile-ducts. The left index-finger is introduced into the foramen of Winslow and supports the hepatic artery, the portal vein, and the ductus choledochus, into the latter of which a sound has been introduced and is seen coming out of the opening in the duodenum. (After Nauwerck.)



FIG. 118.—Method of examining the stomach, which in this case was markedly hypertrophied. Rubber gloves are very useful for this purpose.



FIG. 119.—Removal of the liver from the body. It is held in the left hand and an incision is made towards the operator. This stretches the diaphragmatic attachments of the large blood-vessels, so that they may readily be incised.



incision will be downward and forward, for in the position in which the integument attached to the cervix now holds the penis, the frænum is in front. The amputation of the glans is most conveniently performed with scissors, the body of the penis being supported by the thumb and first finger of the left hand (Figs. 112, 113, and 114).

Nauwerck describes the following method of finding the seminal vesicles. They lie as long, flattened organs on the lateral side of the spermatic duct immediately above the prostate and the posterior wall of the bladder. The fundus of the rectovesical excavation is held up, and the index-finger is placed in the incision in the prostate, the middle finger in the posterior wall of the bladder, and the thumb on the rectum, which on being pulled downward exposes the back part of the neck of the bladder, upon which rest the seminal vesicles. Or, cut through the peritoneum in the depth of the excavatio rectovesicalis, and dissect up the spermatic cord until the vesicles are reached. They are then to be incised and the duct opened up with a fine pair of scissors (Fig. 115). The mucous membrane of the seminal vesicles is of a brownish color, like that of the testicle.

THE DUODENUM AND ITS DUCTS.—The duodenum may be slit while still *in situ*, or it can be excised together with the stomach, liver, and pancreas, and the whole dissected after removal from the body. If a careful dissection of the pancreas be desired, it is well to leave the duodenum attached to the stomach and not to dissect these parts away from the pancreas, which in the Russian language is so aptly called the "under-the-stomach gland." The length of the duodenum is determined by laying a string along the centre of its anterior surface and measuring the same. The gut is best opened with a knife, starting at the tied end about the centre of its anterior surface and with the enterotome cutting more and more to the right until at the pylorus the incision almost reaches the posterior surface of the duodenum (Figs. 116 and 117). Notwithstanding the presence of the glands of Brunner in the lower third of the duodenum, the appearance of the mucous membrane closely resembles that of the jejunum. The papilla, the outlet of the ductus choledochus communis, can usually be discovered if it be remembered that it appears as an elevation of the mucosa near the junction of the second (descending) portion and the third (transverse or oblique) portion of the duodenum, about three and one-half inches from the pylorus, just below the head of the pancreas, and towards the inner and back part of the duodenum. The

duct runs for three-quarters of an inch in the muscular coat of the bowel, where it is usually joined by the pancreatic duct. A small magnifying-glass will often enable one to distinguish the papilla from the valvulæ conniventes. Pressure upon the gall-bladder, as suggested by Virchow, will cause bile to flow out (but care must be taken not to dislodge a gall-stone, either here or in the cystic duct) and thus reveal the opening of the duct. Another way is to follow down the cystic duct, make a transverse incision in it, introduce downward a small probe or splint of broom until this emerges through the opening in the papilla, and then slit it with a knife or scissors. Orth says that if, after finding the head of the pancreas, the intestines are stretched transversely, the outlet will readily be discovered a little below the middle of the head. Congenital diverticula of the duodenum are sometimes found, as well as those of the stomach and œsophagus. Accessory pancreatic tissue may be found hid in the walls of the duodenum. The canal of Wirsung and its accessory canal should be opened. For this purpose the transverse incisions stop at the centre of the gland, and the canal is hunted for. It is usually situated about the centre, is small in size, and is recognized by its pearl-like color. When found, it is dissected out until within about one and one-half centimetres of the ampulla, when a flexible probe or small grooved director is passed through the ampulla. The opening of the caruncula minor is often closed and the pancreatic fluid finds its way out from the portion of the pancreas drained by it by means of the ampulla of Vater, or the converse may be true. Branches from the canal of Wirsung, when dissected out, may lead to accessory pancreases.

THE STOMACH AND ŒSOPHAGUS.—Unless poisoning is suspected (see pages 242 and 343), the stomach is incised along the greater curvature, a *little below* the cardiac orifice and a *little above* the pyloric, the contents are removed, and the openings examined, after which the incision is extended in both directions until the entire viscus is laid open. The mucous membrane may be washed by allowing a gentle stream of water, as from a sponge, to flow over it, but it should not be rubbed with the sponge. The organ may be opened and examined without removal from the body (Fig. 118). Should it be desired to find the artery from which a hemorrhage has occurred in a gastric ulcer, water is injected into the gastric artery supplying this area, and it will be seen to exude from the open part. The usual situation of a gastric ulcer is upon the posterior wall near the pylorus. Examples



of a carcinoma developing from the edges of a gastric ulcer are sometimes found. Guit  ras has pointed out the frequency of small abrasions of the mucous membrane near the pylorus and the collection of small round cells in this vicinity. I know of no extended series of examinations of the gastric contents made after death. In nine cases of pernicious anaemia, Arneill<sup>1</sup> found free hydrochloric acid in none, whereas it was present in some of the gastric carcinomata examined.

The contents of the stomach should be examined as to their quantity, consistency, reaction, odor, gas formation, foreign bodies, color, inflammation, and infectious granulomata. Blood coming from the lungs is apt to be mixed with air, frothy in character, and redder than blood issuing from the   sophagus or the stomach itself, where, if the vessel be of good size, large, compact, blackish-red lumps appear. The blood from cancer is blackish brown (the so-called coffee-grounds appearance); that from diapedesis, cirrhosis of the liver, and inflammations is a brownish homogeneous mixture combined with mucus. The biliary pigments often impart to it a yellowish or greenish hue. In peritonitis and in obstruction of the bowel the gastric contents may be fecal in character.

The most unexpected articles may be found in the stomach,—gail-stones, hair-balls, scarf-pins, glass, rupees (in one case weighing seventeen and three-fourths ounces), hundreds of pins and needles, etc. Thus, eight teaspoons and seventeen other articles were removed by operation from a would-be suicide.<sup>2</sup> Thieves often swallow articles stolen. The larv   of the *Diptera*, maggots of cheese, earth-worms, ascarides, t  ni  , and *Oxyuris vermicularis* have been found. (Ewald.)

The   sophagus is opened up along its anterior surface throughout its entire median extent, either while *in situ*, in case it has not been removed in the manner suggested on page 110 (Fig. 86), or after its removal from the body (Fig. 87). Its caliber may be directly determined by graduated cones, or may be calculated by dividing its circumference by 3.14. Its linear measurements can be made after it has been laid open. The longitudinal folds can plainly be seen, and in their normal state *post mortem* may be discolored. Note carefully the change of color and elevation of the   sophageal epithelium as it passes into that of the stomach. Sometimes it is wise to differentiate by placing in

<sup>1</sup> *Amer. Med.*, January 16, 1904, p. 93.

<sup>2</sup> MONNIER, *Bull. de l'acad. de m  d.*, 1903, vol. lxvii, no. 34, p. 210.



Müller's fluid or in alcohol for several hours, thus coloring or bleaching the part. Diverticula are not uncommon, and an aneurism with a very small opening, usually slit-like, may rupture into the œsophagus. The collateral circulation is often established by means of the veins in the lower third of the œsophagus. Peptic, typhoid, syphilitic, and tuberculous ulcers occur here, as well as abscesses, congenital diverticula, and stricture due to a cicatrix, neoplasm, spastic contractures, etc.

In cancer consider heredity, sex (more common in the male), age (average about fifty years), previous history of a gastric ulcer, and place of origin, inquiring particularly whether or not other cases have occurred in the same house. Cylindrical-celled cancers are found especially at the pylorus, while squamous epitheliomata occur mostly at the cardiac end of the stomach. The tumor may be hard (scirrhus), soft (medullary), or colloid. (a) *Scirrhus*.—The growth starts as a small nodule, usually at the pylorus, often sharply defined, and very hard. It is whitish on section and no cancer-juice exudes from the cut surface. Stricture of the pylorus with hypertrophy and dilatation of the stomach is common. Connective tissue is very abundant and cancer-cells are few. Ulceration occurs late in the disease. (b) *Medullary*.—This tumor tends to become larger than the previous one. It contains much less connective tissue and is therefore softer. It involves all the coats and is not circumscribed. It ulcerates very early and hemorrhages are frequent. As in the previous instance, metastasis is very common. (c) *Colloid*.—This variety usually consists of gelatinous cancer-cells in a condition of colloid degeneration. It extends over the entire stomach and metastasis is very rapid. Metastasis in all the forms affects the various tissues and organs in the following order: lymphatic glands, liver, peritoneum, omentum and intestine, pancreas, pleura, lung, and spleen. The squamous variety is a somewhat flat tubular swelling involving the superficial layers. It may constrict the œsophageal orifice and cause atrophy of the stomach. Cases in which a cancerous stomach has been removed entire during life demand special attention at the postmortem.

Gastrectasis, or dilatation of the stomach, is due to: I. *Pyloric Stenosis*.—(a) Carcinoma. (b) Congenital conditions. (c) Hypertrophy of the pyloric sphincter. (d) Cicatrix of an ulcer. (e) Peritoneal adhesions. (f) Cancer of the head of the pancreas or other structure pressing on the duodenum. (g) Spasm of the sphincter.

II. *Atony of the Gastric Walls*.—(a) From chronic gastritis. (b) Excessive ingestion of solids and liquids. (c) Traumatism. (d) Surgical intervention. (e) Serious infectious diseases. (f) Neurasthenia. (Hemmeter.) At first there is hypertrophy of the muscular walls. Soon, however, interstitial sclerosis comes on, the stomach may become either pyriform or hour-glass in shape, and the mucous coat is thrown into exaggerated folds. As atrophy advances all the layers of the stomach become thinner; the bundles of muscles are separated by connective tissue; the surface may show evidences of pigmentation and petechial hemorrhage; and while the serous surface sometimes remains unaltered, it is usually thick, pale, and opaque.

Gastritis, or inflammation of the stomach: I. *Acute*.—(a) Errors in diet both as regards quantity and quality. (b) Irritant poisons. (c) Mechanical: external (severe injury to the epigastrium); local (fish-bone, etc.). (d) Thermal (hot or cold ingesta). (e) Infectious diseases. (f) Psychic shock (grief, sorrow, etc.). (g) Extension of inflammation. II. *Chronic*.—(a) Follows repeated acute attacks. (b) Slow poisons (alcohol, tobacco, gout, rheumatism). (c) Diet. (d) Anæmia and chlorosis.

I. (a) In simple gastritis the mucous membrane is hyperæmic, swollen, and covered with profuse thick mucus. There are localized areas of ecchymosis and often small erosions. In severe cases there is considerable denudation of epithelium, with perhaps an exudate of grumous blood. (b) Phlegmonous or suppurative gastritis may exist in two forms: the abscesses may be small, multiple, and miliary, or they may be diffuse. The pyloric end is most commonly involved. The submucous and muscular layers are much altered, being swollen, œdematous, purulent, and sometimes even bloody. The mucous membrane overlying the abscess may be normal in appearance, it may slough off, or, again, it may be swollen and hemorrhagic. Abscesses generally grow towards serous and not mucous surfaces. On the other hand, the surface is sometimes studded with numerous areas of focal necrosis of a yellowish appearance, and, on section, may discharge pus. (c) Diphtheritic gastritis sometimes follows laryngeal or pharyngeal diphtheria, and frequently accompanies pyæmia, scarlet fever, variola, and malignant endocarditis. In this form of gastritis we find a variable number of circumscribed areas of false membrane firmly adherent to the underlying structures and leaving a raw surface when removed. It is apt to attack particularly the crests of the rugæ. The

epithelial patches are usually surrounded by areas of more or less pronounced congestion. (d) In toxic gastritis the appearance of the viscus depends upon the amount of contained food at the time of ingestion and the concentration and kind of poison. If the latter is diluted, the mucous membrane alone suffers; if concentrated, all the coats may be involved. Alkalies appear to be more destructive than acids, the lesions produced resembling those of an intense congestion, more or less localized. Around an area of necrosis is a brown-black eschar. In very severe cases perforation may follow. Sloughs or ulcers are almost invariably found where the poison has been concentrated. Mycotic gastritis may be due to: (a) Anthrax. (b) Favus. (c) Thrush. II. *Chronic*.—(a) *Hypertrophic*.—Virchow calls a condition of the mucosa when there are swelling, cloudiness, and a yellow color, gastritis parenchymatosa or glandularis; it is due to poisons, as arsenic and phosphorus, to acute infectious diseases, to acute atrophy of the liver, etc. This may be localized or diffuse. In the former case numerous mucous polyps can be seen over the affected area. This variety occurs in drunkards. These warty elevations show considerable cystic degeneration. In the diffuse variety the stomach is almost invariably enlarged and the walls are thickened, particularly the mucous coat, which is decidedly velvety both to sight and touch, slate gray in color, with nodular, deeply injected areas of scarlet and brown-red thickened patches. Besides being swollen, rugæ are often present in exaggerated folds. Petechial hemorrhages and areas of pigmentation are common. There are often evidences of previous ulcerations (cicatrices). The stomach frequently contains a variable quantity of thick, tenacious, sour smelling, greenish mucus. (b) *Atrophic*.—When this variety of the disease exists the walls of the stomach become thinner. There is connective tissue overgrowth, which by its contraction causes the epithelial cells to undergo degeneration and disappear. The mucous membrane is thin, smooth, and pigmented.

In hemorrhage from the stomach, if the blood come from without, as from a rupture of an aneurism, the stomach presents but few changes. The blood may be fluid or clotted; it may be bright red or dark in color. When the hemorrhage is due to actual disease of the stomach, this blood is apt to be coffee-brown. Petechial hemorrhages in the mucous membrane are common. Extensive hemorrhage from the wall of the stomach is most usually associated with gastric ulcer. Behrend reviews the symptoms of those cases in which death resulted



from the diagnostic and therapeutic inflation of the stomach with carbon dioxid gas.<sup>1</sup>

**THE LIVER AND GALL-BLADDER.**—The clinician, having felt the lower border of the liver during life, often wants to know its exact situation at the postmortem, and is disappointed, on reading the report of the autopsy, if he does not there find what he desires. The attachment and the presence of any lesions near the suspensory ligament are carefully noted. In the round ligament are sometimes found small collections of blood in places where this vessel has not become entirely obliterated. The bile may now be collected in a sterilized tube such as is described on page 347. The so-called "corset-line" produced by tight lacing may be due to other causes, as a pleuro-pneumonia or subphrenic abscess.

The liver is removed from the body by severing its attachments to the diaphragm, falciform ligament, blood-vessels, and ducts, and breaking up existing adhesions. For this purpose traction is made by introducing the left hand behind the right lobe and raising the liver so that it hangs over the ribs of the right side (Fig. 119). Nauwerck removes the organ by finding the hepatoduodenal ligament and then, introducing the index-finger into the foramen of Winslow, pulling it somewhat towards the duodenum and cutting, from right to left over the finger, the ductus choledochus to the right, the hepatic artery on the left, and, lastly, the portal vein with its four main branches lying between the two posteriorly (Fig. 117) and quite constantly distended with blood. The liver is then weighed and measured, and the color, normally of a chestnut brown, and the condition of the surface are noted. The true color of the surface of the liver is best determined from an examination of its anterior aspect, as its lower part is apt to be bile-stained and, being in contact with the intestine, is more apt to show post-mortem changes. The right and left lobes may sometimes be measured separately with advantage. I have sometimes made a tracing of the outline of the liver by cleansing it from blood, placing it on paper, and then drawing with a pencil its outlines, indicating in their proper places any lesions which may be noted or the areas from which pieces are cut for microscopic study. As blood, when fresh, is quite adhesive, the paper must not be folded until any blood which may be on it has dried. After examination of the serous surface of

---

<sup>1</sup> *Med. News*, December 19, 1903.

the gall-bladder and duct, the sac should be laid open by a longitudinal incision carried through the duct. To find the ductus choledochus, first note the situation of the gall-bladder and then follow down the cystic duct either with the eye or by dissection to where the hepatic duct joins it. The bile-duct running to the right of the portal vein may then be dissected out to its outlet at the papilla in the duodenum. In case the liver is to be removed at once, the dissection should be



FIG. 120.—Method of incising the liver. Long parallel incisions are made from the right lobe to the left, care being taken not to cut entirely through the organ, which would prevent reconstructing it in its normal state, nor to extend the incisions so deeply as to injure the gall-bladder. If desired, the liver may now be turned and a second set of incisions at a right angle to the first may be made upon its posterior aspect. The structures of the under surface of the liver have been previously dissected out in a manner similar to that described in the case of the lungs. The history of the case will usually give information as to which of the vessels may have to be sacrificed in the dissection. On p. 344 will be found methods for injecting the vessels with different colored materials.

continued beyond the place where it is to be cut. Should resistance be met with in the passing of a probe during the process of dissection, such a part should be at once investigated. The hepatic duct may be opened with scissors until it has branched several times in the substance of the liver. A bacteriologic examination of the bile-ducts or gall-bladder may now be made. It will be recalled that the portal vein is formed by the union of the splenic and superior mesenteric veins and after running three or four inches divides in the liver substance



into two main branches. The inferior mesenteric vein may empty into the splenic or superior mesenteric or take part in the formation of the portal vein.

The liver is laid on its posterior surface and a series of parallel incisions about half an inch apart, which do not completely pass through the organ, are made, either longitudinally or, still better, transversely (Fig. 120).

In pernicious anæmia the presence of free iron may be shown by placing a thin strip of hepatic tissue in a ten per cent. solution of potassium ferrocyanide for several minutes and then washing it thoroughly with a two per cent. solution of hydrochloric acid. The production of a blue color (Prussian blue) indicates the presence of iron. When Lugol's solution is applied to test the presence of the amyloid reaction, it is well to remember that the glycogenic reaction is produced by the iodine. A weak solution of the violet of Paris is recommended by Letulle for securing the amyloid reaction. Observe: Bile-ducts: (*a*) caliber,—normally that of a thin goose-quill, may be closed or may be of the size of a finger; (*b*) gall-stones; (*c*) ulcers. Portal vein: (*a*) color of blood; (*b*) thrombosis; (*c*) caliber,—may be thin, as result of old inflammation; (*d*) periphlebitis. I recall a case operated upon for cirrhosis of the liver where the postmortem showed an infected thrombus of the portal vein. Gall-bladder: (*a*) size; (*b*) adhesions; (*c*) tumors; (*d*) contents,—1, bile (note its color,—light or dark yellow, reddish yellow, greenish yellow,—quantity, quality, etc.); 2, foreign bodies,—gall-stones; 3, mucous membrane,—thickening, change in color, and inflammation. Liver: (*a*) position; (*b*) size,—increased in parenchymatous inflammation, decreased in atrophy; (*c*) form,—fissures or granular distortion of surface; (*d*) color,—brown or brownish red normally, yellow in fatty infiltration, dark brown in atrophy, gray in amyloid and interstitial overgrowth, ochre-yellow in acute yellow atrophy, green in icterus, or dirty green when decomposition sets in; (*e*) consistence,—normally rather hard (pitting soon disappears), increased in amyloid disease, the pitting remaining for some time, softer in parenchymatous affections and early stages of acute yellow atrophy, fluctuates in echinococcus cysts and abscesses; (*f*) capsules,—normally transparent, but thickened in chronic inflammation, syphilis, etc.; (*g*) section,—smooth, uneven, rough, or granular; (*h*) lobules,—notice that they are separated by connective tissue, more distinct in cirrhosis, less so in acute yellow



atrophy. It is well to remember that in man the separation of the lobules by the connective tissue of Glisson's capsule is not at all well marked. Observe whether the lobules are larger or smaller than normal. Notice that the color is darker in the centre of the lobule than at its periphery (cyanotic induration). See if the periphery is yellow (fatty infiltration). On section note whether the tissues retract.

PANCREAS.—The position of the pancreas having been determined in the preliminary examination of the abdominal cavity, its isolation and detachment are attended by no difficulty unless there be disease of neighboring parts, in which case its removal may necessitate taking an additional viscus with it. Many students are singularly unfamiliar with the normal anatomy of the pancreas, the splenic artery often being mistaken for the pancreatic duct and the sensation of hardness which this gland normally imparts to the touch being regarded as an evidence of sclerosis; the head and tail of the pancreas, too, are not infrequently left in the body and thus escape examination. In warm weather the pancreas is early affected with signs of decomposition, consisting in a brownish-red color, softening of its tissue, and the escape of a greasy brownish-red serum. Disease may extend from the pancreas to the portal vein, bile-ducts, pylorus, or duodenum, or from these organs to the pancreas. Hemorrhages, tumors, degenerations, calculi,<sup>1</sup> atrophic changes, cysts, etc., may be found in this organ. The possible presence of fat necrosis—a not infrequent cause of sudden death—should be borne in mind. The submucous, interstitial, or subperitoneal patches of pancreatic tissue in the wall of the intestine should not be forgotten. Thorel<sup>2</sup> describes seven cases of accessory pancreas, in the stomach, intestines, and mesentery, the opening, except in the latter case, going to the intestinal tract. Langerhans's islands were absent in nearly all the cases examined.

EXAMINATION OF THE RETROPERITONEAL LYMPH-GLANDS, DIAPHRAGM, VENA CAVA, CHYLE-DUCT, ETC. — The retroperitoneal lymph-glands, best exposed by dividing the vertebral attachments of the mesentery at its roots, may be thickened from inflammation (as in syphilis) or be the seat of primary tumors (especially sarcoma and lipoma), secondary cancer, amyloid degeneration, and tuberculosis,

---

<sup>1</sup> MÜLLER, *Proceedings Path. Soc.*, December 10, 1903. These stones may be found in the intestines.

<sup>2</sup> *Virchow's Archiv*, vol. clxxiii, no. 2.

or may have undergone changes due to various other inflammatory, cystic, and systemic affections.

Examination of the diaphragm may reveal the existence of hernia, abscess on the under surface, perforation (as in echinococcus cysts or amœbic abscess of the liver), trichina spiralis, inflammation of its serous investment, fatty degeneration and brown atrophy, hypertrophy (as in obstruction to normal respiration), atrophy (as in pseudo-hypertrophic muscular atrophy), etc. The muscular fibres of the diaphragm may undergo granular, cloudy, or fatty degeneration. Atrophy may arise from lesions of the phrenic nerve or as a part of a general muscular atrophy. Traumatic rupture and congenital deficiency are occasionally met with.

The vena cava and the aorta should be inspected for signs of inflammation, thrombosis, etc. To remove the aorta it should be grasped as high up as possible, drawn forcibly forward, and cut obliquely from within and above outward and downward. In order to secure a firmer hold one finger may be inserted in its lumen. (Orth.) Its elasticity should always be tested by pulling both longitudinally and laterally. The color should also be noted and the presence of atheromatous patches and plates described, especially when found around or near the point of exit of its various branches, a frequent cause of arteriosclerosis.

The receptaculum chyli rests behind the aorta, mainly on the body of the second lumbar vertebra, and between the pillars of the diaphragm and the insertion of the psoas muscles. It arises from three roots which spread out over the third lumbar vertebra. As the thoracic duct ascends, it crosses above the left azygos vein, and lies between the aorta and the right azygos vein, and has a caliber of from three to eight millimetres. At the fourth dorsal vertebra it passes behind the œsophagus and opens into the left subclavian vein, at or near the entrance of the left common jugular vein. It is readily found by dissecting away, with an up-and-down movement of the tip of a grooved director, the cellular tissue situated at the top of the arch of the aorta and the œsophagus. This is near the left subclavian artery and *before* the duct which bends around this artery, like a shepherd's crook, to terminate in the vein. It has a rosy-white tint with longitudinal striæ, and at this point gives off few collateral branches. The azygos vein is much larger and not nearly so elastic. It can be opened by splitting with a pair of fine scissors. The *right* thoracic duct emp-

ties into the vena anonyma and collects the lymph from the upper part of the right thorax, neck, heart, and upper extremity. Failure to find the duct may be due to its previous removal while still attached to the descending thoracic aorta. Tuberculosis of the thoracic duct is quite common in abdominal tuberculosis.

Abscesses in the psoas muscles may be secondary to Pott's disease, coxitis, perforation of the intestine, tumors, etc. Examine the spinal column for kyphosis, lordosis, and scoliosis.

In death after fright and from chloroform narcosis, a large amount of blood is collected in the abdominal veins, as the result of vasomotor paralysis.



## CHAPTER XI

### DISEASES OF THE GENITO-URINARY TRACT

**KIDNEY.**—No little confusion exists in the description of the pathologic lesions of the kidney, owing to the multiplicity of terms employed. A classification of renal diseases depending upon the structure affected is: 1, epithelial (parenchymatous) nephritis; 2, fibrous (interstitial) nephritis; 3, vascular nephritis. It should be borne in mind that there is no such thing as a perfectly pure form of nephritis and that the condition which predominates gives the name to the lesion. For example, when we speak of parenchymatous nephritis, we do not mean that the epithelial cells alone are affected without involvement of the connective tissue, for it is entirely proper to describe a case as chronic parenchymatous nephritis in which the interstitial changes are beginning to predominate. The epithelial cells of any portion of the kidney may be affected primarily, hence the name glomerulonephritis, etc.

*Amyloid Changes.*—These may be due to (a) prolonged suppuration (tuberculous or syphilitic), (b) chronic disease of the kidney, or (c) lack of cardiac compensation. The amyloid kidney is usually enlarged (the condition occasionally occurs in a contracted kidney), pale in color, and firm in consistency. The capsule is adherent in places and shows petechial hemorrhages beneath it. The cortex is increased in size. The glomeruli are first affected and usually prominent, although the cortex is pale in contrast to the somewhat reddish color of the pyramids. The organ has a bacony or waxy appearance. The urine contains albumin. The tube-casts are hyaline, waxy, or finely granular. Edema of the extremities is common.

*Congenital Defects.*—(1) Total absence. (2) Absence of one, with hypertrophy of the other. (3) Rudimentary, cystic. (4) Duplication. (5) Partial coalition, usually lower end (horseshoe). (6) Remnants of fetal lobulation.

*Congestion.*—(a) In traumatism the kidney is large; the capsule is tense; the color is dark red. On opening the capsule the contents are found to be soft and bulge out and blood drips freely from the surface of the section. The dependent portions are more congested than

the cortex. In passive congestion the organ is enlarged and firm; the capsule strips off readily; the cortex is wider than normal; the surface on section looks coarse and connective tissue is plainly visible; the cortex is of a deep-red color and the pyramids are of a purple-red. Congestion may be due to (*b*) drugs, as cantharides or turpentine, (*c*) infectious fevers, (*d*) alterations of the circulation in the kidney itself or in the vena cava (rare), (*e*) valvular lesions of the heart, (*f*) diseases of the liver, or (*g*) diseases of the lungs. Hæmatoma of the kidney occurs, sometimes reaching a large size and holding over a quart of blood and clots.

*Cystic Disease.*—(*a*) Congenital cystic kidneys are greatly enlarged, so much so at times as to impede labor. There may be a conglomeration of cysts varying in size from that of a pea to a small apple. In some cases no renal tissue can be seen without the aid of a microscope. The cysts are lined with flattened epithelium and contain a fluid in which are found albumin, blood-crystals, cholesterin, triple phosphates, and fat-drops. (*b*) Chronic nephritis (which see). (*c*) Adenocystomata, of similar origin as the corresponding cysts in the ovary. (*d*) Concretions block up the uriniferous tubules and press upon the still intact epithelial cells, which later become flattened and disappear. The stroma and vascular supply are next affected and a cystic condition is produced, or the disease may go on to the formation of large concretions.

*Hydronephrosis.*—The outflow of liquid from the pelvis of a kidney may be obstructed by (*a*) congenital deformities, as when the pelvis comes off too high up on the kidney, (*b*) twists of the ureter, (*c*) calculi, (*d*) morbid growths, or (*e*) cicatricial bands. There is an accumulation of non-purulent fluid, which by steady pressure produces an atrophy of the organ and a gradual distention of its pelvis. The papillæ become more flattened and disappear, and their place is taken by concave recesses in the medulla, which becomes narrower. In extreme cases the kidney may be converted into a large cyst with some imperfect septa. There may be an enormous quantity of the contained fluid or only a few ounces. It is yellowish in color and contains urea, uric acid, and sometimes albumin and sugar. There is usually compensatory hypertrophy of the opposite kidney.

*Infarcts.*—(*a*) Calcareous infarcts extend through the tips of the papillæ as stripes through one-half or more of the medulla, mainly along the canals, but also in interstitial tissue. There is effervescence

on the application of hydrochloric acid. (b) Uric acid—found as acid ammonium urates in very young children and as acid sodium urates in mature years in cases of gout—may be deposited within the kidneys in the form of flakes (uric acid nephritis or gouty kidneys). In babes they appear as yellow radiations from papillæ into medulla, and show that the child was born alive, as they occur only after breathing has taken place. Sodium hydrate dissolves the acid ammonium urates. (c) Hæmoglobin occurs in hæmoglobinuria. It exists in the canals first as lumpy brown, later as granular, and seldom as crystalline masses. Hæmatoidin crystals are seen where old hemorrhages were (Virchow). (d) Bilirubin infarct gives the bile reactions. It occurs in the icterous new-born, in acute atrophy of the liver, and in progressive pernicious anæmia. (e) Infarcts caused by salts of silver are very rare. (f) Hemorrhagic infarcts. (g) Anæmic infarcts.

*Interstitial Nephritis.*—In acute interstitial nephritis the whole kidney is increased in size; the color is uniform, making it hard to distinguish the border line between the cortex (which is swollen) and the medulla. The process is essentially a productive one. There is a marked migration of the leucocytes and the connective tissue undergoes proliferation. The cells increase in number and the intercellular substance disappears. The pus-cells get between the epithelial cells and the lumina of the canals can no longer be followed. Such areas may be found anywhere in the kidney substance. The process is essentially due to pyogenic bacteria brought from the heart, as in malignant endocarditis, or the uterus, as in puerperal sepsis. The process ends in abscess formation, often affecting the perinephric tissues. (Langerhans.) A similar condition may start from without the kidney or extend up from the pelvis or further down the urinary tract.

Chronic interstitial nephritis may start as an acute form, but most frequently affects alone the connective tissue of its stroma, the blood-vessels not being involved. The process naturally ends in contraction. The canals are freed from their epithelial cells and the glomeruli may be brought so close together as to touch each other. The capsule is adherent and the surface lumpy or granular and grayish red in color. The cortex is much smaller and may measure only a few millimetres in thickness, but its consistence is markedly increased. Compensatory hypertrophy may occur. If the canals are fatty they appear as yellow stripes or points. Cysts are common and are most marked at the junction of the cortex and medulla. The vessel walls are thickened. Local-



ized interstitial nephritis is usually syphilitic, while the diffuse form is due to gout, lithæmia, lead, over-indulgence, etc. In the latter form we have granular atrophy, the so-called red granular kidney, in which, as contraction takes place, cysts are found.

*Movable Kidney.*—(a) Especially in females. (b) Due to absorption of perinephric fat. (c) Repeated pregnancies. (d) Traumatism. (e) Displacement by tumors. As a rule, the displacement is not great. The kidney usually moves downward or upward and inward, generally rotating so that the outer border and upper end move forward and the hilum is directed inward and backward. Nearly all cases are associated with a medial displacement of the colon. The right kidney is the one most frequently affected.

*Parasites.*—Of the parasites the following are found: (a) *Distoma hæmatobium* (*Bilharzia hæmatobia*). (b) *Filaria sanguinis hominis*. (c) Echinococci. (d) Cysticerci. (e) Pentastoma. (f) *Strongylus gigas*. All are rare in this country.

*Parenchymatous Nephritis.*—Acute diffuse inflammation of the kidney is due to: (a) Acute infectious fevers. (b) Poisons,—e.g., turpentine, arsenic, etc. (c) Traumatism. (d) Exposure to cold and wet. Macroscopically the organ is swollen, tense to the touch as the capsule is stretched, but the substance of the kidney is softer than normal, the color is gray to yellowish, and the stellate veins on the surface are prominent. The capsule strips off easily and is somewhat thinner than when normal. On slitting the capsule the renal substance bulges out. The cortex, which is increased in amount, is somewhat pale, swollen, and soft; the glomeruli appear as minute red dots. The pyramids are distinct and striated. The radiations in the medulla may be gray or transparent, gelatinous or watery. The larger blood-vessels are overfilled and prominent.

*Parenchymatous Nephritis, Subacute.*—The large white kidney is more swollen than in the acute form and the tissue itself is of greater consistency. The cortex may be increased, therefore, before contraction commences. Yellow spots where the degenerative changes are most marked are found in the gray glossy substance. Cysts are absent, unless interstitial changes are associated. The kidney is dry on section, and the pyramids of the medulla show reddened stripes pointing towards a papilla. This condition may be associated with amyloid degeneration, most marked in the glomeruli. The mucous membrane of the pelvis is frequently swollen and of a pinkish color.

Microscopically the changes are those of an acute diffuse inflammation, including cloudy swelling, proliferation, desquamation, and a granular change in the cells lining the tubules. The straight connecting tubules may entirely escape, though there is a form of catarrhal nephritis, usually of an ascending variety, in which this part of the kidney is alone affected. In the surgical kidney there is an acute parenchymatous nephritis with abscess formations. Each individual cell is larger, the transverse diameter of the tubule increased, and the lumen diminished or even obliterated. Death most frequently takes place before the degenerative changes are complete; otherwise resorption and contraction follow, and on the surface there are slight indentations, often associated with a hemorrhagic condition, hence bloody casts, as in poisoning by cantharides and potassium chlorate, where even pigmentary infarcts may be found. The urine is scanty, high colored, albuminous, and contains casts and free blood. There may be an extensive œdema, with effusions into the serous cavities.

*Parenchymatous Nephritis, Chronic.*—This process is latent and runs a slow course, often of years; not all of the kidney is affected at once, some portions showing normal parenchyma while at other places degenerative changes are going on and at still others degeneration is complete and the parts are already in an atrophic condition. The cortex contracts irregularly, and has not the regular granular appearance seen in the kidney affected with interstitial nephritis (Langerhans), nor is there much increase in the stroma except at those places where contraction has taken place.

*Perinephritic Abscess.*—(a) Traumatism. (b) Extension of inflammation from the kidney or from neighboring organs. (c) Perforation of the bowel. (d) Infectious fevers, particularly in children. The kidney is surrounded by pus, especially posteriorly. The abscess-cavity is usually extensive. The pus is often offensive and may have a distinctly fecal odor. It may burrow and discharge into the lung, bowel, peritoneum, or bladder, or it may follow the psoas muscle and appear in the groin.

*Pyelitis and Pyelonephritis.*—(a) Tuberculosis. (b) Infectious fevers. (c) Calculi. (d) Cystitis. (e) Tumors. (f) Drugs. (g) Cold and wet. *Classification.*—(a) Simple catarrhal. (b) Purulent. (c) Hemorrhagic. (d) Calculous. In simple acute pyelitis the mucous membrane of the pelvis is swollen, hemorrhagic, and turbid. In the purulent form the mucous membrane is swollen and covered with

a cream-like exudate of a yellowish or yellowish-green color. Ecchymoses are common. The kidney itself is enlarged, softened, œdematous, grayish in color, and shows little distinction between cortex and medulla. Areas of necrosis or miliary abscesses are distributed through the kidney substance. The kidney may attain the size of a human head. It is usually firmly adherent to the adjacent organs, tissues, and vessels. A quart of pus may be contained in the cavity; in these extreme cases all appearance of the gland substance may be lost. The hemorrhagic variety occurs in anthrax, sepsis, and leukæmia. In calculous pyelitis the mucosa is roughened, grayish in color, and thickened. There are also more or less dilatation of the calyces and flattening of the papillæ. These may be covered by a gray membrane. After the renal substance has been destroyed, if the pelvic orifice is still obstructed, the pus may become inspissated and ultimately impregnated with the salts of lime.

*Stones.*—The following varieties of stone may be found in the kidney or its pelvis: (*a*) Oxalate. This is very hard, dark, brownish yellow or gray in color, with rough surface and mulberry shape. (*b*) Uric acid. This is usually smooth or a little rough, light brownish yellow in color and often striped, and of medium consistence. (*c*) Phosphate stones are white, crumbling, and chalky. (*d*) Cystin and xanthin stones are rare.

*Tumors.*—(*a*) Fibromata are the most common of benign tumors. (*b*) Lipomata. (*c*) Myxomata. (*d*) Myomata. (*e*) Angiomata. (*f*) Lymphadenomata (or lymphomata). (*g*) Rhabdomyomata. (*h*) Carcinoma may be primary or secondary; it is comparatively rare. The cancer may infiltrate the whole cortex or may be knotty and separated sharply from the surrounding tissue. (*i*) Sarcoma may be primary or secondary. It is more common than cancer, usually occurs in children, and may attain to an enormous size. Here it exists as a myosarcoma. (*j*) Carcinoma sarcomatodes. (*k*) Patches of adrenal tissue may start growing and give rise to large tumors, the so-called hypernephroma heterotopes. Such growths are by no means rare. (*l*) Deposits of liver-tissue may be found in the kidney, especially in the cortical layer.

*URETERS.*—In some malformed fœtuses both ureters are absent. In other cases double bilateral ureters are found. They may open into the vagina or the uterus. Stenosis often occurs as a congenital or acquired condition. Cysts and polyps are found not infrequently.



Parasites, as the *Distoma hæmatobium*, round worms, and echinococcus, are found. A calculus may fill the pelvis of the kidney and extend down in the ureter.

BLADDER.—The color of the vesical mucous membrane is normally a pale gray, but is red in recent inflammation and blackish red if the inflammation be very severe. The mucosa affords a favorite location for the multiplication of various organisms, which usually reach the bladder either from the kidney or from the urethra. Typhoid bacilli may frequently be detected in the urine of patients suffering from typhoid fever. Cystitis is due to irritants in the urine, extension of inflammation from adjacent parts, traumatism, septic infection through the blood of the urethra, infectious diseases, stricture of the urethra, enlarged prostate, or diseases of the cord (myelitis). It may be œdematous and especially hyperæmic after the ingestion of certain poisons, as phosphorus and cantharides. When caused by the colon bacillus, it may give the agglutinative reaction in a dilution of one to fifty of the blood of a patient suffering therefrom. In acute cases the mucous membrane is swollen, reddened, and covered with a thin film of mucus or pus. The veins may be distended, especially when hemorrhoids exist and venous thrombosis occurs. When hemorrhage has occurred, the surface of the membrane is of a universal gray tint or mottled with gray, black, or reddish-brown patches. In severe cases necrosis, abscess, or even perforation may occur. In the diphtheritic form of the disease necrotic patches are seen and also small hemorrhages in the region of the trigone and the surrounding fundus. These tend to increase in size. There is submucous swelling, which subsequently becomes infiltrated with pus. The whole mucous membrane over it degenerates and can easily be removed from the muscular coat. In chronic cystitis the bladder may be enlarged, but it is often smaller than when normal. The various coats are much thickened and there may be true hypertrophy of the muscular coat. This condition is best seen in cases of long-standing chronic cystitis, where the inner surface may even be thrown into folds and roughened so that the picture resembles that of the interior of a heart, and shows how difficult it is for injections into the bladder to cleanse thoroughly the walls when there is inflammation. In severe cases the inner coats often feel rough and sandy to the touch, on account of encrusted salts. Gangrene and tuberculosis may occur. There is a considerable variety of tumors in the bladder: adenoma, carcinoma, fibroma, myoma, sarcoma, and

mixed tumors. They frequently assume the form of polyps and villi. Cavernous angiomas, dermoids, and echinococcal cysts are met with. Pockets (diverticula) may develop in the walls of the bladder, sometimes being covered only by the peritoneum. Their openings may be very small, though the size of each diverticulum may reach that of a hen's egg. These pockets are at times produced by and may contain stones. In exstrophy the inner surface of the bladder is exposed externally above the pubes through a hiatus in the median abdominal wall. The intestines may protrude into or open through the bladder. Professor Guit  ras once related to me an interesting case of primary diphtheria which developed upon this exposed mucous membrane. The organ may be completely or partially divided by an anteroposterior septum. The bladder may be entirely absent or may be double. Cases have been reported in which the two bladders were of the same size and located the one behind the other. The remnant of the urachus may undergo cystic change. In hypertrophy dilatation of the cavity exists along with increase in the thickness of the wall, which may extend an inch or more. The female bladder may become inverted and appear through the urethra. It may also take part in herni  e of various forms. The bladder may be ruptured by external violence, of which there may be no external visible sign. In over-distention from hemorrhage the bladder may reach to the umbilicus or may open into the rectum or vagina (vesicorectal or vesicovaginal fistula). In the interesting condition called trichosis vesic  e the hair is usually referable to the breaking of a dermoid cyst into the bladder or it may be a product of growth from the mucosa itself. In one case—an autopsy on a female—I could not see where the dermoid had arisen if not in the walls of the bladder itself. In a body examined at the Pennsylvania Hospital a bundle of hair was found which had become encrusted with salts, thus forming a calculus.

Garrod<sup>1</sup> gives the following causes of black urine: (1) jaundice, especially when of long standing; (2) h  maturia; (3) h  moglobinuria; (4) h  matoporphyrinuria; (5) melanotic sarcoma; (6) alkaptonuria; (7) ochronosis; (8) abundance of indican; (9) long-standing pulmonary tuberculosis; (10) the taking of certain drugs and articles of diet; (11) certain rare cases of undetermined nature.

Vesical calculi, usually associated with some form of cystitis, may

---

<sup>1</sup> *The Practitioner*, March, 1904.

contain any of the normal or abnormal constituents of the urine. If this liquid be allowed to stand, precipitation occurs, the character of which depends upon the acidity or alkalinity of the urine. Bacteria in the body may cause an alkaline decomposition, with the formation from urea of carbon dioxide and ammonia, which uniting with the uric acid forms ammonium urates and triple phosphates. The most important sediments are uric acid, sodium urate, ammonium urate,—all of which give the murexide test,—calcium oxalate, calcium carbonate, calcium diphosphate, calcium triphosphate, and triple phosphate. Concretions may be found in the form of sand or as calculi. They are held together by an albuminous or cement-like substance, to which may be added cast-off epithelial cells, shreds of tissue, blood, mucus, etc. Primary stone formation may take place in urine which has not undergone decomposition; such calculi are usually composed of uric acid and urates. Secondary stone formation occurs in an alkaline urine, the starting-point being a foreign body introduced through the urethra from without or a small calculus which has found its way down from the kidney; these stones are apt to be composed of ammonium urates and phosphates. They often consist of different substances concentrically arranged. Metamorphosed calculi are produced where, for example, a primary stone has been partially dissolved by the action of an alkaline menstruum and the remainder covered by secondary deposits. (Schmaus.) Calculi assume a large number of shapes and differ much in size. In addition to those named above, cystin and xanthin stones exist.

Parasites in the bladder are rare. The following have been found: (a) *Distoma hæmatobium* (*Bilharzia hæmatobia*). (b) *Filaria sanguinis hominis*. (c) Echinococci. (d) Cysticerci. (e) *Pentastoma*. (f) *Eustrongylus gigas*.

FEMALE GENITAL TRACT.—*Fallopian Tubes*.—One Fallopian tube may be absent or rudimentary, and, on the contrary, I have seen an oviduct lengthened to over ten inches by traction from a growing uterine fibroid, and have observed in a tube extra openings supplied with fimbriæ, the presence of which might at times have an important bearing upon the question of ectopic pregnancy. This dangerous condition may occur anywhere within the tube, or the fecundated ovum may escape into the abdominal cavity or become caught in a corpus luteum of either ovary. It is doubtful whether ovarian extra-uterine pregnancy ever existed; in two cases so diagnosed and brought to me



for examination, careful study showed that fecundation occurred near the ostium, and the fimbriated extremities became attached to the ovary just as in a case of ovarian abscess, making it to appear as if the pregnancy had started in the ovary. Zinke tabulates a series of 88 cases of simultaneous intra- and extra-uterine pregnancies. An interesting abnormality is lithopædion, where a foetus may stay in the abdominal cavity for thirty or forty years with certain of its tissues remaining recognizable. The convoluted interior of the oviduct offers a favorable place for the growth of various organisms, especially the gonococcus, the streptococcus, the colon bacillus, and the organism of tuberculosis. The tube itself may be affected with cysts and with many kinds of benign and malignant tumors, the latter being primary or metastatic. It is subject to hemorrhages and different forms of inflammation.

In acute salpingitis the Fallopian tubes are swollen. The neighboring blood-vessels are dilated, tortuous, and overfilled with blood. There is often a considerable exudate on the serous surfaces, causing adhesions of the tubes to the surrounding structures. On section the lumen of the tubule is found to contain serum (hydrosalpinx), mucopurulent matter (pyosalpinx), or hemorrhagic fluid (hæmatosalpinx). The tube may rupture and give rise to a general peritonitis. The mucous membrane is thickened, swollen, and often intensely congested. To show the ciliated cells, though these may have been shed by the inflammatory process, care must be taken to harden the tissue at once after removal and according to the methods for showing karyokinesis. In chronic (proliferous) salpingitis the tubes may become enormously thickened, hard, and resistant to the touch. The adhesions to surrounding tissues are very marked and more or less completely organized. The new connective tissue contracts, throwing the organs out of their proper relation and often obliterating their normal appearance.

*Ovaries.*—These show perhaps a greater variety of pathologic changes than any other part of the body. The ovary may be divided into lobes by bands of connective tissue, or actual duplications of the parts may occur. Supernumerary ovaries are found. An ovary may form part of a hernia, and in a child I found one in the canal of Nuck. The organ may be absent, hypoplastic, or prolapsed in abnormal position. As the opportunity arises, study the differences between a true and a false corpus luteum and a corpus hæmorrhagicum. These glands are subject to various forms of inflammation, an entire ovary at times

being converted into a pus-sac. They are often bound down by adhesions, and in later life undergo senile atrophy and may even become calcareous or calcareous concretions may be found in them. They undergo hypertrophy. Among the tumors here found may be mentioned adenomata, dermoid cysts, which are of an almost endless variety, enchondromata, endotheliomata, fibromata, fibromyomata, myomata, cancers, cystomata, sarcomata, psammocarcinomata, angioma, etc. Dropsy of follicles, fungous excrescences, and tuberculosis occur. Ovarian cysts may grow to an enormous size and contain over two hundred pounds of fluid. The ovaries may be enlarged in mumps.

*Uterus.*—In examining the womb notice any abnormalities on the exterior and be sure to search every portion of the interior for any lesions which may exist. The situation of the organ may be markedly altered, both as to its entirety and its individual parts. Thus, we may discover anteflexion, anteversion, retroflexion, retroversion, prolapse, inversion, dilatations, elongations, bendings, or even find it forming part of an inguinal or a crural hernia. The chief congenital malformations are uterus bicornis, bicornis duplex, bilocularis, subseptus, and bipartitus, unicornis, didelphys, cordiformis, septus duplex, and double uterus. The uterus is subject to atrophy, hypoplasia, rudimentary (infantile) atresia, stenosis, and hypertrophy. Uterine tumors are of great variety,—adenoma, adenocystoma, cancer, deciduoma malignum (syncytioma malignum), hæmatoma polyposum, fibroma, myoma, myofibroma, myosarcoma, lipoma, leiomyoma, etc. A placental polyp may assume destructive characteristics. Hydatid moles occur, and at times number several thousand. Fleishy moles are the result of hemorrhage into the decidua. Dermoid cysts are found. Hemorrhages are common, and, besides those due to menstrual disturbances, are often associated with polyps, cancer, etc. After parturition and after the menopause marked changes take place in the blood-vessels, which may undergo amyloid degeneration. Infarcts are seen. Many varieties of endometritis exist, such as gonorrhœal, tuberculous, diphtheritic, syphilitic, decidual, fibrous, gangrenous, glandular, interstitial, catarrhal, purulent, mycotic, villous, etc. Langerhans describes an interesting case of an old woman in which the womb was so enlarged by a solid mass of thrush fungi and other bacteria that it measured some two inches in diameter. The uterus may rupture, as from childbirth, trauma, etc.

In acute forms of endometritis the mucous membrane is red,

swollen, and sodden; the discharge is profuse, stringy, and often purulent; in severe cases blood is present. If infection follow contusion during labor, there may develop a suppurative process which transforms the parts into a soft, stinking, grayish-green or brown mass that tends to become gangrenous. The cervix is the most often involved. In diphtheritic endometritis there is formed a thick grayish-yellow or white membrane, the decidua lying loosely on the surface. The process may begin and remain at the placental insertion or may involve the cervical portion of the uterus. The infection may spread through the lymph stream or blood-vessels. Acute (ulcerative) endocarditis is a frequent complication of puerperal infection. In hemorrhagic endometritis the mucous membrane is red from engorgement of blood-vessels and numerous punctiform hemorrhages. It is distinguished by the condition of the ovaries from a similar appearance in menstruation. Tuberculous endometritis sometimes resembles carcinoma in gross appearance, but on microscopic and bacteriologic examination shows the presence of tubercle bacilli.

In chronic hypertrophic endometritis there is a hyperplasia of the mucosa, with softening and congestion, forming polypoid excrescences; the glandular structures also hypertrophy, become occluded, and form cysts of various sizes. In the cervix enlarged Nabothian cysts should be looked for. In atrophic endometritis the mucous membrane becomes thin and pigmented and the glandular structures disappear. Follicular erosion of the cervix occurs after lacerations.

Foreign bodies may be found in the uterus. These are introduced to prevent conception or to produce abortion, or find place through surgical manipulation or expulsion from some adjacent organ. Twenty-four hours after the birth of a full-term child the uterus weighs from seven hundred to twelve hundred grammes. Friable, elevated, mushroom-like fibrous masses formed after the removal of the placenta are found in the area where it was attached.

*Vagina.*—The vagina may be absent or appear as a mere connective-tissue cord. It may be wholly or partially divided by a longitudinal or transverse septum. It may be entirely closed or so small that coitus, if attempted, takes place through the urethra, which thus becomes markedly dilated. The normal flora is considerable, and of pathogenic organisms the *Gonococcus*, *Bacillus diphtheriæ*, and *Oidium albicans* (thrush) are of importance. It is well to remember that diplococci other than the *Gonococci* are frequently found in the vagina.



After rape the condition of any secretion present should be noted,—whether dried, fluid, watery, or purulent; also its color and odor. Examine microscopically for spermatozoa and gonococci. Observe all discharges in cases of abortion. Severe inflammation may follow, and even gangrene may supervene. Erosions, fissures, fistulæ, and lacerations are seen after labor. Syphilitic ulcers are common; those of a tuberculous nature are rare. Tumors of the vagina include cysts, carcinoma, fibroma, fibromyoma, myxoma, rhabdomyoma, and sarcoma. Malignant neoplasms of the cervix uteri may by extension of the growth involve the vaginal walls, which should, therefore, always be examined in such cases. The epithelioma is almost always verrucose or nodose. Prolapse often accompanies tumors or is seen in multiparæ accompanied by prolapse of uterus, rectocele, and cystocele. The exposed surface is eroded, covered with ulcers and patches of necrosis. Vaginal hæmatocele, hernia, abscesses, and hypertrophic vegetations are not uncommon. Poisons, as mercury, arsenic, have been found in the vagina. In cases of abortion various materials have been discovered.

MALE GENITAL TRACT.—*Testicles*.—The testes are subject to many lesions, but the exposed situation and the extremely specialized character of these organs are sufficient to account therefor. The undescended testicles are peculiarly liable to injury and the subsequent development of tumors. Adenoma, sarcoma, enchondroma, fibroma, osteoma, myxoma, and rhabdomyoma occur. Dermoids and mixed neoplasms containing cartilage are not rare. Inflammation is common. In typhoid fever the condition of the testicles should always be noted, as they may become infected with the typhoid bacillus. They may be affected by syphilis, tuberculosis, leprosy, etc. In guinea-pigs infected with glanders the testes are especially apt to become diseased. Sarcoma or a cancer of these organs is often indistinguishable. Cysts also occur, often combined with tumors. Hemorrhage may take place in the tunica vaginalis and the testicle may atrophy owing to pressure from the fluid in a hydrocele. True abscesses are found in them, and they may undergo brown atrophy, glycogenic infiltration, pigmentary or amyloid changes. In elephantiasis they may show hypertrophy. The cords sometimes rupture and varix is common. Albers Schoenberg experimenting on guinea-pigs found that Röntgen rays caused necrostermia (in 197 minutes) and azoospermia (377 minutes). The action of radium upon these organs should also be studied.

*Spermatic Cord.*—The cord may become twisted; this condition is, as a rule, associated with undescended testicle or swollen epididymis. The cord may be thickened or lobulated. Its arteries sometimes show marked atheroma and its veins varices. I have also several times seen the duct itself converted into a rigid thickened tube by salt-like deposits.

*Prostate.*—The most common lesion of the prostate is enlargement due to interstitial hyperplasia. This is almost always accompanied by atrophy of the gland cells. Acute inflammation is very common. Abscesses occur; they are often not recognized until their sequelæ are prominent, and are usually seen in the area around the posterior urethra. Cystic formation (cystic adenoma) is seen in many enlarged prostates, but true neoplasms are much less common. Carcinoma and fibroma are the most frequent. Moderate atheroma and endarteritis are seen almost always in the arteries of the part. In men over sixty years of age small prostatic calculi or sand, often dark in color, are quite commonly found on careful sectioning of the prostate.

## CHAPTER XII

### DISEASES OF THE LIVER AND PANCREAS AND THEIR DUCTS<sup>1</sup>

**LIVER.**—*Abscess.*—Hepatic abscesses may be multiple, often originating from the appendix, or single, as in the amoebic abscess of the tropics. Perihepatitis is usually present, and rupture into the pleural cavity may occur.

*Acute Yellow Atrophy.*—This is an acute disease of the liver, presumably of infectious origin, characterized by a rapid fatty degeneration of the organ, with invariably fatal termination. Due to: (a) A specific micro-organism(?). (b) The ordinary micro-organisms of suppuration and infectious diseases have been found in this condition. (c) Certain poisons,—e.g., phosphorus. (d) Female sex. (e) Pregnancy or the puerperium. The liver is greatly reduced in size,—one-half to one-third; in one of my cases, however, the condition had been preceded by hypertrophic cirrhosis and the organ weighed over five pounds. The liver is thin, flattened, and flabby, the capsule is wrinkled, and the gland is of a pale-yellow color. Both on the surface and on section may be seen a number of orange-yellow patches, in the centre of which are usually marked hemorrhagic areas. The remainder of the liver is of a yellowish-brown or mottled color. The outlines of the lobules are very indistinct. The bile-ducts and gall-bladder are empty. Bilirubin crystals may be seen under the microscope. If a section of the liver be allowed to remain in the air for some time, a thin, white coating appears on its surface, which on examination is found to consist of crystals of leucin and tyrosin. The adjacent organs are usually stained with bile and present numerous hemorrhages, especially on the surface. The spleen is enlarged and the heart and kidney show marked granular

<sup>1</sup> Those wishing to go more deeply into this subject will find WARING's *Diseases of the Liver, Gall-Bladder, and Biliary System* (1897) and OPIE's *Disease of the Pancreas* (1903) most instructive reading. VIRCHOW's remark in his "*Post-Mortem Examinations*" (1876) of "the slight importance of the pancreas" is interesting as showing the small consequence attached even until a short time ago to the functional activities of this gland. An account of the "Zuckergussleber" and fibrous polyserosities will be found in ROSE, *Würzburger Abhandlungen aus dem Gesamtgebiet der prakt. Medizin*, 1904, vol. iv, no. 5, and KELLY, *Trans. Coll. of Phys. of Phila.*, 1902, p. 62. CAMMIDGE's article on the chemistry of the urine in diseases of the pancreas is contained in the *Lancet* of March 19, 1904, p. 782.



change. The color of the liver in acute yellow atrophy depends on the time at which death took place: in the earlier stages the organ is ochreyellow, in the later stages it is mottled, and if much blood be present it is grayish red.

*Amyloid Degeneration.*—Found in cases of: (a) Prolonged suppuration, tuberculous or syphilitic. (b) Infectious fevers. (c) Chronic visceral diseases with cachexia. The liver is large in size, smooth in outline, and pale in color. The edges are distinctly rounded; small hemorrhages are common on the surface. On section the surface is anæmic, semi-transparent, and infiltrated. It presents the characteristic lardaceous or waxy appearance. The process may be a localized or a generalized one; in either case staining by Lugol's solution is never uniform, as the diseased brown spots appear only in certain areas. The characteristic coloration may be seen upon the lining of both hepatic and portal vessels. Early in the disease this reaction is hard to detect, except by special stains under the microscope. Very thin pieces of the liver should be sectioned with a scalpel and put in a small glass dish. Add a solution of iodine and then wash out with water. Put something white under the dish and the characteristic coloration can be more readily seen.

*Bile.*—T. Kimura<sup>1</sup> has investigated the bile taken from the human gall-bladder shortly after death. He finds its pigments to be variable in quantity, being low in tuberculosis and high in conditions of stagnation, such as heart disease. The specific gravity varies from 1012 to 1040, and the dry residuum from 2.68 per cent. to 20.63 per cent. The relative viscosity varies widely,—from 1.46 to 58.24. These factors are all greatly increased in cases of obstruction of the common bile-duct. Urobilinogen is found regularly, urobilin very frequently; but both are wanting in cases of complete biliary obstruction, marked diarrhoea, and in the new-born. This fact supports the enterogenous theory as to the formation of urobilin. Normal faeces contain urobilin regularly, but it is wanting in cases of biliary obstruction. Meconium does not contain any. In a case of obstruction of the cystic duct, a hitherto undescribed brown pigment was found.

*Cancer.*—I. *Secondary Cancer.*—Most common. Histologically shows same structure as primary growth, which is usually in the stomach, bowel, or pancreas. The liver is enormously enlarged, irregu-

---

<sup>1</sup> *Deut. Arch. f. klin. Med.*, 1904, vol. lxxix, p. 274.

lar, and nodular. The nodules are usually symmetrical, often superficial, flattened, discrete, and umbilicated; they may be more or less evenly distributed throughout the liver. On section whitish masses of varying size are seen, contrasting with the red color of hepatic tissue, the yellow staining of bile, pigmentation due to blood, and the light-yellow areas of fatty degeneration. The cancerous masses may undergo fatty degeneration, suppuration, or fibroid change. II. *Primary Cancer*.—Rare. (a) Massive. Causes great enlargement. On section the mass is uniform grayish white in color, somewhat firm, and distinctly outlined from the liver substance. (b) Nodular. Large and small nodules are scattered throughout the organ. These usually consist of a primary growth and numerous secondary nodules. (c) Cancer with cirrhosis is rare. Liver not much enlarged. Surface of section is grayish yellow, studded with nodular yellowish masses. In one of my cases of primary cancer of the gall-bladder the cancerous portions and the liver had become infected by the *Bacillus pyocyaneus*.

*Cholecystitis, Acute Infectious*.—There exists an acute inflammation of the gall-bladder due to: (a) The introduction of pyogenic micro-organisms,—for example, the *Bacillus coli communis* and the typhoid bacillus, pneumococcus, staphylococcus, and streptococcus. (b) Gall-stones. (c) Extension of inflammation from the bile-ducts. The gall-bladder is distended; its walls are thickened and tense. The mucous membrane is swollen, hyperæmic, and may be covered with a purulent exudate. The submucosa may also be involved. The contents of this sac are cloudy and dark in color, and may be mucopurulent or hemorrhagic. Orth states that the inflammation is usually of a necrotic character. The tissue is of a dirty yellow-brown color and sometimes is rotten and easily torn. Gall-stones are frequently present. The cystic duct is often obliterated. There may be adhesions with the bile-duct or omentum. The common bile-duct may be congenitally absent and yet the child may live five months.<sup>1</sup> The gall-bladder may be absent without serious impairment of the hepatic function.

*Cholelithiasis*.—Gall-stones may be formed within the gall-bladder or in the ducts leading to or from it. Consider: (a) Most frequent in females. (b) Age, fifty per cent. over forty years old. (c) Sedentary habits. (d) Overeating. (e) Carcinoma(?). (1) The calculi are usually multiple, rarely single. They vary in size as well as

<sup>1</sup> MENZIES, *Australasia Medical Gazette*, January 20, 1904, p. 20.

in number. When multiple they are faceted, sometimes mulberry-shaped. They are of a dark bluish or greenish color. On section there is a nucleus consisting of epithelium, rarely a foreign body, then comes a layer of inspissated bile-salts, the outer covering being cholesterin. There may also be bile-acids, fatty acids, salts of calcium and magnesium, with a trace of iron and copper. When the stones consist of pigment exclusively, they are very easily broken and vary from yellowish brown to black in color. When composed of cholesterin entirely, they are softer, easily indented with the finger-nail, but not brittle, and are crystalline, the crystals forming layers. They are colorless and more or less transparent, but turn blue when iodine and sulphuric acid are added. They generally consist of both pigment and cholesterin, which may be combined or may be separated by layers. These stones are usually firm in consistence, rarely friable. (2) The gall-stones may lead to impaction of the gall-bladder or to obstruction of the cystic and common ducts or even of the bile-duct alone. There may be formation of a fistula, external or internal, with escape of bile. The bladder itself is much thickened, sometimes dilated, sometimes smaller than normal through chronic inflammation.

*Cirrhosis.*—Under this heading are classified various forms of disease of the liver characterized by a marked increase of its connective tissue, which may be capsular, interlobular, or intralobular, with or without increase or decrease in the size of the organ. Causes: (a) Alcohol. (b) Certain infectious diseases,—e.g., syphilis, tuberculosis, malaria, scarlet fever. (c) Micro-organismal infection. (d) Mechanical obstruction to the onward flow of the blood. (e) Rickets. (f) Anthracosis. (g) Poisons, as phosphorus and cantharides. *Classification.*—(a) Alcoholic. (b) Fatty. (c) Hypertrophic. (d) Capsular. (e) Syphilitic. (f) Cyanotic. (g) Malarial. (h) Scarlatinal. (i) Tuberculous. (j) Rhachitic. (k) Anthracotic. (1) In the atrophic cirrhosis of Laennec the organ is greatly reduced in size, although in the beginning it may be slightly enlarged, and later is altered in shape. The surface is irregular and nodular and the capsule thickened. The nodules are usually small, but in some cases they may be greatly increased in size. The tissue is firm, hard, and resistant to the knife. The surface of section presents a mottled appearance, the lobules being divided by bands of connective tissue. The liver substance itself is of a yellowish or greenish-yellow color. The areas of connective tissue which are periportal are gray. (2) In fatty cir-



rhosis, found usually in drunkards, the organ is enlarged, somewhat smooth, although often slightly granular. It is paler than normal and of a yellowish-white color. It is firm and resistant to the knife. The capsule is opaque and often much thickened. The peritoneal cavity usually contains ascitic fluid. The membrane is opaque and thickened. Chronic involvement of the stomach and small intestine is always present. The spleen is enlarged; the kidneys are often cirrhotic. Owing to interference with the portal circulation by the cirrhotic liver, extensive compensatory circulation is formed. The abdominal vessels above and below the umbilicus are markedly enlarged. Around the umbilicus is found the caput of Medusa. Acute tuberculosis of the peritoneal cavity may be associated with it. (3) Hypertrophic cirrhosis is most common in young men. Ackerman compares it to elephantiasis. The organ is enlarged, but the outline is normal. The surface is usually smooth and its color an olive-green; the consistency of the organ is increased and the capsule is thickened. The surface of section is uniformly greenish yellow and the lobules may be separated by distinct bands of connective tissue. The spleen is greatly enlarged. Jaundice is a marked symptom of this disease. Ascites is usually absent. (4) In capsular cirrhosis there is enormous thickening of the capsule, which is irregular and somewhat wrinkled, producing great contraction of the liver. The organ itself is rarely markedly cirrhotic, its tissue being usually soft. Chronic capsulitis of the spleen, chronic perisplenitis, and ascites are often present. The kidneys usually show granular change. (5) In syphilitic cirrhosis the liver is markedly irregular in shape, being divided into peculiarly shaped lobes by extensive bands of fibrous tissue traversing the organ in indefinite directions. In one of my cases over forty distinct lobulations were present. The cut surface is mottled, often fatty in appearance, and shows the presence of gummata or of syphilitic scars. The connective-tissue bands are of a gray or reddish-gray color. (6) For cyanotic cirrhosis see Passive Congestion of the Liver. (7) In malarial cirrhosis the liver is markedly enlarged, commonly extending to the level of the umbilicus. It is firm in consistence, of a dark-red color, smooth in outline, and bleeds freely on section. (8) Klein has pointed out that chronic interstitial hepatitis may follow an attack of scarlet fever, which may account for some cases of cirrhosis of the liver in children. (9) Rhachitic cirrhosis is a form of the disease in which there is a marked increase of connective tissue around the individual lobules. (10) Anthracotic cirrhosis occurs in coal-miners,

in whom the coal-dust may occasionally reach the liver in sufficient quantities to cause a marked connective-tissue formation about the portal canal. (Welch.) Sears and Lord from a study of seventy-eight autopsies of hepatic cirrhosis consider the condition to be part of a systemic disease.

*Congestion.*—(a) Acute infectious diseases. (b) Traumatism. (c) Extension of inflammation,—*e.g.*, from the intestines. (d) Valvular heart-disease. (e) Pressure of tumors. (f) Other mechanical obstructions to the circulation. The condition is most marked when the veins of the liver are closed, as in periphlebitis or Chiari's endophlebitis. (1) The post-mortem appearances of active congestion are not characteristic. The liver is swollen, dark in color, and full of blood; the hyperæmia is not limited to any one portion of the liver substance. (2) In passive congestion the liver is large in size, smooth or slightly granular in outline, and of a distinctly mottled hue. The surface of section presents the characteristic nutmeg appearance (the centre of the lobule being deeper), due to a marked congestion occurring in the central veins, the congested tissues being of a reddish-brown color. This is surrounded by a large area of a pale-yellowish color (fatty infiltration), with a third zone of cellular infiltration and new connective tissue. In rare cases this order is reversed, the congested area occurring at the periphery of the lobe and the lighter or fatty parts towards the centre. In chronic and well-marked cases there may be considerable induration and shrinkage of the liver substance, with irregular surface, so that the hypertrophy gives place to an atrophy, called cyanotic atrophy or Virchow's red atrophy.

*Emphysema.*—Portions of the liver when squeezed under water show the escape of bubbles. This condition may be due to putrefaction or to the growth of gas-forming organisms during life.

*Fatty Changes.*—(a) Middle life. (b) Alcohol. (c) Sedentary habits. (d) Infectious fevers. (e) Certain poisons. (f) Cachexias. (g) Interference with local or general circulation. *Classification.*—(1) Fatty degeneration. The liver may be increased or diminished in size. The capsule may be smooth or wrinkled. The consistence is usually somewhat decreased; the organ is paler than normal and somewhat mottled in appearance. Periphery of lobule is first involved. The surface of section is smooth, usually bloodless, and

---

<sup>1</sup> *Bost. Med. and Surg. Jr.*, September 11, 1902, p. 285.



imparts a greasy stain to the knife. The general color is a dull gray or grayish yellow. (2) In fatty infiltration the liver is often markedly enlarged, normal in outline, smooth to the touch, and of a somewhat pale, excessively fatty color. Globules of fat may be readily expressed with a knife. Hyperæmia may obscure the characteristic appearance.

*Hepatitis, Suppurative.*—Abscess of the liver may be due to: (a) Traumatism. (b) Extension from neighboring organs,—e.g., the bowel and the pleura. (c) Pyæmia. (d) Amœbic dysentery. (e) Malignant emboli. (f) Diseases of veins, as periphlebitis, thrombophlebitis, and thrombo-umbilicalis. (g) Stoppage of bile, as from gall-stone or dead ascarides. (h) Idiopathic tropical disorders. *Classification.*—(a) Pyæmic hepatitis. (b) Portal pyæmia. (c) Pyo-septicæmia or multiple abscess. (d) Tropical or endemic hepatitis. (e) Suppurative cholangitis. (1) In multiple abscess the change in the liver depends upon the number of the abscesses. If these be few, the liver walls may be comparatively little altered; if they are very many, the liver is apt to be enlarged, softened, and friable. The abscesses themselves appear as minute foci which are non-encapsulated, the centre containing a thick white, yellow, or greenish pus surrounded by a zone of congestion. The abscesses may number from five to ten, or many hundreds. These multiple abscesses frequently arise from pyæmic embolism of the portal vein or hepatic artery or vein, or they may result from a cholangitis. They may be generally distributed or appear in clusters. If from a malignant endocarditis, they are usually situated under the capsule. (2) Large abscesses occur in two forms,—the large chronic encapsulated abscess surrounded by a pyogenic membrane and the tropical or amœbic abscess. (See Dysentery.) The large abscess is usually single; there may be two or more. The right lobe is usually affected. There is a distinct limiting membrane. The pus is usually of a greenish-yellow color and often of a disagreeable odor. The surrounding substances often show but few changes, except as the result of pressure.

*Sarcoma.*—This may be primary (very rare) or secondary. The most frequent variety is the secondary melanosarcoma following sarcoma of the eye, of the skin, or of the penis. In these cases the liver is greatly enlarged, weighing as much as fifteen pounds, and the secondary nodules, which are of a black or slate color, are usually uniformly distributed throughout the gland. In primary sarcoma of the liver there are but few nodules, and these reach a large size, measuring



at times five or six inches in diameter. Metastases to other organs often occur, though other portions of the liver may escape.

*Other Tumors.*—In addition to carcinomata and sarcomata, the liver is the seat of adenomata, adenocystomata, angiomata, fibromata, and aberrant adrenal tumors similar to those found in the kidney. The cavernous angiomata are usually small in size and, when found, are usually seen on the surface of the liver in elderly persons. They may be injected with colored material by means of any of the hepatic blood-vessels, and then form excellent microscopic specimens for future study. A cystic liver may be associated with a similar condition of the kidneys.

*Parasites and Infectious Diseases.*—*Psorospermia*, *Pentastomum denticulatum*, *Distomum hamatobium*, *Distomum lanceolatum*, *Distomum hepaticum*, and *Echinococcus*. Cases of primary tuberculosis of the liver have been reported, and syphilitic lesions are by no means rare.

*PANCREAS.—Anomalies.*—The tail of the pancreas is sometimes bifid, and peculiar divisions made by septa of connective tissue may occur in all parts of this organ. The pancreatic tissue by surrounding the duodenum may cause an intestinal stricture. Accessory pancreases have already been referred to in the technic of removing the gland from the abdominal cavity.

*Acute Pancreatitis.*—This condition exhibits acute degenerative changes in the parenchymatous cells and an exudation into the interstitial tissue. It is often associated with cholelithiasis, and is usually hemorrhagic or gangrenous. In the lower animals pancreatitis may be produced experimentally by the injection of an artificial gastric juice, but it is impossible to foretell which form of the disease will result. *Hemorrhagic Pancreatitis.*—This variety of pancreatitis is usually associated with gastric or gastroduodenal dyspepsia, slight swelling of the epigastrium, and obstinate constipation. The pancreas is enlarged and infiltrated with blood. There is a cellular and fibrinous exudate present, with a necrosis of the parenchyma; also disseminated necrotic foci are found in the omentum and peritoneum. This condition often ends in *gangrenous pancreatitis*, where the organ is enlarged, swollen, soft, friable, of a color varying from mottled red and gray to dark brown and black, and gives off a foul odor. The extension of the disease to the neighboring tissues may result in almost complete sequestration of the pancreas. In some cases it has been found that the organ has entirely disappeared, its place being taken

by an abscess-cavity containing a foul-smelling mass, which may discharge through the intestine. Disseminated fat necrosis often follows. Gangrenous pancreatitis may be the result of a perforating inflammation of the gastro-intestinal or biliary tracts, arterial sclerosis, and hemorrhagic pancreatitis. *Pancreatic Hemorrhage*.—The pancreatic vessels may rupture from trauma, or the hemorrhage may accompany tumors, cysts, purpura, eclampsia, and acute infections. Apoplexy in this organ is seldom associated with arterial disease. Extensive fat necrosis now and again accompanies it. Opie considers that pancreatic hemorrhage and hemorrhagic pancreatitis represent a single pathologic process.

*Chronic Interstitial Pancreatitis*.—This disease is related to diabetes mellitus, is secondary to morbid changes in the intestines, the bile-passages, and the liver, and is associated with arterial sclerosis, syphilis, tuberculosis, and abuse of alcohol. The islands of Langerhans are frequently not affected. *Chronic Pancreatitis*.—In the interlobular form of this disease the gland is hard, dense, and nodular, with a granular surface. On section the tissue is compact and homogeneous, the loose areolar tissue being replaced by scar-like bands. The islands are unaltered, and the acini have atrophied nuclei and dilated lumina. Lymphoid cells are present in great numbers. In the interacinous variety the gland is tough and shows newly formed connective tissue in the lobules. The lesions are diffuse and irregular in distribution; in one place thickening of the connective tissue and of the network supporting the acini may occur, while elsewhere are found compact bands or small masses of stroma. Lobulation is observed, and at times is associated with general pigmentation. The change in either case may be only microscopic. Fatty infiltration may obscure both types. Chronic pancreatitis may be due to obstruction of the pancreatic duct, to pancreatic or biliary calculi, to malignant growth compressing or invading the organ, to an ascending infection from the duodenum, to alterations of the blood-vessels, to arterial sclerosis, to association with chronic tuberculosis of other organs, and at times to alcohol and to cirrhosis of the liver. It is very rarely due to syphilis.

*Congenital Syphilitic Pancreatitis*.—In this form of the disease the organ is enlarged and firm. There is a diffuse interstitial proliferation of the interlobular and interacinous tissue, first with atrophy and finally ending in destruction of the parenchymatous elements. The arteries are the seat of a syphilitic periarteritis and the adventitia is

infiltrated with lymphoid cells. Finally the capillary network around the acini disappears. The islands of Langerhans are not involved, but are surrounded by newly formed stroma.

*Hyaline Degeneration.*—This especially attacks the islands of Langerhans, and is often found in cases of diabetes, destroying the islands and obstructing the vascular supply of a large portion of the parenchyma. Hyalin is deposited between the capillaries and the parenchymatous cells. The affected areas are larger and more numerous in the tail of the pancreas and may involve two-thirds of the tissue. Epithelial cells are found arranged about a lumen, particularly at the periphery of the altered tissue, and show that the acini are also affected. In the head and body of the gland the areas are smaller and fewer.

*Fat Necrosis.*—This consists of small opaque white areas found in the fat around the pancreas, which are made up of necrotic fat-cells. In disseminated fat necrosis small foci are widely scattered in the fat of the abdomen. Large foci occur, especially in the fat of the omentum. Both a subperitoneal and a retroperitoneal fat necrosis indicate some grave alteration of the pancreas. These areas are frequently surrounded by a narrow hemorrhagic zone. While these lesions are usually limited to the fat in the abdominal cavities, they are found, as Hauseman has observed, in the subcutaneous fat corresponding in location to the reddish areas visible during life upon the overlying skin. They are probably caused by the action of the fat-splitting ferment upon living fat.

*Diabetes Mellitus.*—A constitutional disease characterized by the continued secretion of large amounts of pale cloudy urine, of high specific gravity, containing glucose and, at times, acetone, diacetic acid, and beta-oxybutyric acid. It is, as a rule, associated with excessive hunger and thirst and sometimes with increase in fat, and at other times with progressive emaciation. It occurs most frequently in adult males, Hebrews being especially predisposed. It is due to some failure properly to utilize certain carbohydrates in metabolism. There is a tendency to destructive changes in the tissues and to death from coma. It may be caused experimentally by ingesting phloridzin and by puncture of floor of fourth ventricle. Glycosuria is seen with exophthalmic goitre, certain neuroses, some diseases of the liver, cirrhosis, lesions of the pancreas, injuries to the nervous system, destruction of gray matter in the floor of the fourth ventricle, extirpation of cervical ganglion, pancreatic calculi, atrophy, carcinoma, necrosis, fatty degen-



eration, cysts, acute and rarely chronic interlobular pancreatitis. Some cases of the disease show an absolutely healthy pancreas, yet it occurs often with chronic interacinous pancreatitis. Diabetes is closely related to destructive lesions of the islands of Langerhans, especially with hyaline changes and interacinous pancreatitis. In diabetes the number of islands may be diminished and the pancreas be nearly always atrophied. Arterial sclerosis accompanies many cases of diabetes; also, acromegaly. It is occasionally associated with tabes. The celiac ganglion is atrophic in this disease (Orth). Neuroretinitis is very common, and there may be hemorrhages in the retina and opacities in the vitreous. The most usual change is a thickening and congestion of the membrane. The blood generally appears normal, but contains an increased amount of glycogen, and may be loaded with finely divided fat which floats on the surface in a cream-like layer. There may be lipæmic clots in the vessels. Fat embolism of the pulmonary vessels has been described. The myocardium is pale and soft; rarely it may be hypertrophied. Advanced fatty degeneration of the muscular fibres is the characteristic change in long-standing cases of diabetes. Croupous pneumonia and bronchopneumonia, chronic interstitial pneumonia, and tuberculosis are common complications; many of them terminate in gangrene. The lung may soften (malacia) and, becoming mixed with stomach secretions *post mortem*, form the so-called *pneumomalacia acida*. It has a sour but not a gangrenous odor. The spleen is usually small, pale, and soft, but may be enlarged and congested. Diffuse nephritis with fatty degeneration, and frequently glycogenic degeneration, most marked in the pyramids, may occur. Boils, carbuncles, onychia, eczema, and gangrene of the extremities are common. The liver is usually enlarged, often congested, abnormally firm to the touch, and gives the glycogen reaction; fatty degeneration is common. Do not mistake diabetes mellitus for alkaptonuria; in the latter disease pigmentation of the cartilages or ochronosis may occur.<sup>1</sup> Some alkaptonuriacs do not show ochronosis, but thus far all the cases of ochronosis exhibit alkaptonuric changes in the urine. Established facts concerning diabetes: (1) Considerably more than one-half of all cases are due to a destructive disease of the pancreas. (2) When due to disease of the pancreas, injury to the islands is responsible for the disturbance of metabolism in the

<sup>1</sup> OSLER, *Lancet*, January 2, 1904, p. 10.

carbohydrates. (3) Common lesions injuring the islands are chronic interacinous inflammation and hyaline degeneration. (4) Other lesions of the pancreas do not attack the islands of Langerhans, but produce diabetes by destroying the interacinous islands along with the secreting parenchyma. (Opie.) According to the experiments of Sauerbeck<sup>1</sup> upon guinea-pigs, total extirpation of the pancreas or the tying of its secretory duct gives rise to atrophy of the islands of Langerhans and the subsequent development of diabetes.

The Bremer-Williamson reaction of diabetic blood may be obtained a considerable time after death; the procedure is as follows: Forty cubic millimetres of water are placed in a small, narrow test-tube; to this are added twenty cubic millimetres of blood, one cubic millimetre of a one to six thousand aqueous solution of methylene blue, and forty cubic millimetres of liquor potassæ. The test-tube is placed in boiling water for four minutes, at the end of which time, if the blood is diabetic, the blue color will have disappeared and a dirty-green color will have taken its place. Williamson obtained the reaction in forty-three cases of diabetes tested and thinks it is due to an increase of glucose in the blood. The reaction is of special value in coma where urine cannot be obtained.<sup>1</sup>

*Hæmatochromatosis (Bronzed Diabetes).*—This disease is due to a disturbance of iron metabolism. It affects various organs. Brown pigment is deposited within certain tissues and gives a microscopic pigmentation. Most of the glands are deep brown and their secreting cells are reddish-yellow or tinged with ochre-colored granules. The parenchymatous cells and Kupffer's cells of the liver contain pigment. A yellow fine pigment is seen in the smooth muscle-cells of the stomach, intestine, blood- and lymph-vessels, rarely in the urinary bladder, ureter, vas deferens, Glisson's capsule, or splenic trabeculæ. This second pigment contains no iron. These symptoms associated with cirrhosis of the liver are called Recklinghausen's hæmatochromatosis. Pigmentation is seen also in the liver and spleen in cases of anæmia.

<sup>1</sup> *Ergebnisse der allg. Path. u. path. Anat.*, 1904, eighth year, part II, p. 691.

<sup>2</sup> BROWN, *International Clinics*, January, 1903, p. 266.

## CHAPTER XIII

### EXAMINATION OF THE SKULL AND BRAIN

THE body is placed in the supine position on the side of the table nearest the operator, with the head, elevated by a block placed under the neck and occiput, projecting slightly beyond the end of the table. If the cadaver be in a coffin or box, it may be drawn to the upper end thereof, the head being raised and placed upon a board laid across the top, the back supported by a head-rest, a block of ice, or any convenient bundle of rags or paper. Of the various forms of support employed, the Cornell head-rest (Fig. 45) is peculiarly well adapted for holding the head steady.

Note any anomaly in the size or shape of the head. (See page 361.) The scalp should be subjected to the same careful preliminary scrutiny for evidences of disease or injury, remote or recent, as the other parts of the body. It is then divided by an incision extending from one mastoid process to the other (Fig. 121), passing over the vertex when the hair is abundant and about midway between the vertex and the external occipital protuberance when it is thin. If the hair be long, it should be parted along the proposed line of incision, in order that as little of it as possible may be cut (Fig. 122). For the same reason and to guard against damage to the knife, the cutting edge of the scalpel or cartilage-knife should be directed from the skull when the scalp is being cut. When all the tissues overlying the skull have been separated, the scalp is reflected backward and forward by force, the calvarium being exposed from the occiput to or slightly beyond the frontal eminences. The eyes and nose should be protected by pledgets of cotton placed beneath the anterior flap. Care should be taken to avoid tearing the scalp at the extremities of the incision behind the ears, especially if the posterior incision with a large anterior flap be made. Indeed, it is for this reason that the incision is begun and ends behind and not in front of the ears, for a tear behind the ear would hardly be noticed, while one in front would cause considerable disfigurement. The scalp may be so adherent to the cranium—a condition more apt to occur in the posterior segment than in the anterior—as to necessitate its removal by dissection with the knife or scraping with a chisel. Whatever instrument is used, guard against its slipping, lest



injury be done to the operator or to the subject. Avoid undue traction of the scalp, which would cause it to present a baggy appearance when replaced.

The skull should next be examined in detail. Fractures and other evidences of injury may now be revealed which could scarcely have been discovered in the preliminary examination. Note should be made of the presence of atrophy, hypertrophy, or softening of the bone, of premature or delayed synostosis and supernumerary bones, of tumors, syphilitic or tuberculous abrasions or openings, marks of previous trephining, of asymmetry and abnormal coloration, the "greenish-yellow" discoloration due to osteomyelitis or the "citron-yellow" due to tertiary syphilitic lesions, etc. (For cranial measurements and pathologic types of skull see Chapter XXIV.)

There are two methods of removing the calvarium,—the angular, in which the skullcap is sawed in two intersecting planes meeting behind the ear, and the circular, in which the bone is divided in a single plane. The former method is usually to be preferred, as it permits more secure reposition of the skullcap, but the latter is easier of application and will, therefore, be considered first.

**THE CIRCULAR METHOD.**—The path of the saw, which may be marked with a pencil or the point of a knife, traverses a plane cutting the skull from half an inch to an inch above the glabella anteriorly, an inch or an inch and a half above the external auditory meatus laterally, and passing just above the inion posteriorly. This line will cross the temporal muscles obliquely, and they and their fascia should be divided with a knife instead of the saw, in order that their edges may be accurately approximated for suturing when the skullcap is replaced.

Sawing the skull is no easy task; it may be greatly facilitated by the employment of an electric or dental engine. For this part of the operation it is a decided advantage to be ambidextrous. While the sawing is being done with one hand, the head must be steadied with the other, placed either on the vertex or on the face and protected by a towel, for the saw is liable to slip, especially when first applied. The scalp, especially of a female, should be protected from "sawdust" by wrapping towels about it. Proffered assistance should be declined, because, while it is natural to look out for one's own fingers, it is impossible effectively to guard another's. The reason I often give for not accepting aid is that "I am reasonably supposed to know



FIG. 121.—While the right ear is held back with the left hand an incision is started directly over the mastoid process. The remainder of the incision over the vertex will be made from within outward, thus avoiding dulling the knife and cutting the hair.



FIG. 122.—After the initial incision has been made behind the ear, the hair may be parted when it is long, so as not to injure it when incising the scalp.



FIG. 123.—Method of sawing the skullcap. The temporal muscle has been cut through with a knife in the direction of the future sawing, and a pencil mark shows the posterior line along which to saw. The hand is protected with a towel.



FIG. 124.—Angular method of removing the brain. The saw markings in each case pass close to the ear and meet an inch or so above it. The left hand is covered with a towel to protect it from injury.





FIG. 125.—Method of breaking up the inner table with an old knife after sawing. (There are also various forms of chisels made especially for this purpose.)



FIG. 126.—Method of drawing off the skullcap with a retractor after the sawing is completed.



FIG. 129.—Appearance of the dura mater after removal of the calvarium, showing the superior longitudinal sinus and the meningeal vessels.



where my hand is, but not where yours may be." The saw may be carried entirely through the bone or, better, only to the inner table, this being divided with chisel and hammer. In no case, however, where it is suspected that the skull may have been fractured should the latter procedure be adopted, as the force of the blow required might be sufficient to split the bone. While a post-mortem fracture may be recognized by the absence of extravasated blood, the enlargement of a pre-existing fracture is more difficult to differentiate. A receptacle should be placed beneath the head to catch the cerebro-spinal fluid and the blood that escape when the skullcap is removed and the meninges are opened, and care must be taken to prevent spattering. The calvarium is loosened by twisting a chisel or the sharp end of a hammer in the kerf, and removed with a blunt hook. If instead of an instrument the fingers be used for the purpose, they must be well protected, as they are liable to slip and be abraded by the sharp edges of the bone. Traction should be made steadily and not in jerks, lest from a sudden giving way the calvarium be damaged by falling on the floor or surrounding objects be soiled by the spattering of blood or other fluid. When, as is sometimes the case, the calvarium does not readily yield to traction applied in front, it may often be easily detached by inserting the hook posteriorly. If the dura be adherent, as not infrequently happens in cases of chronic alcoholism, old injuries, or sunstroke, it may be loosened with a blunt instrument, or it may be divided along its margin with a pair of blunt-pointed scissors or a curved, probe-pointed bistoury cutting from within outward, the falx cerebri being incised close to the corpus callosum. In children under seven years of age this must always be done, as up to this time of life the dura is normally adherent to the osseous structures of the skull.

**THE ANGULAR METHOD.**—In this method the skull is sawed in two planes which by their intersection form an obtuse angle at a point a little below and slightly posterior to the apex of the ear. Always try to saw above the line of the hair in front. Although this makes the anterior fossa deeper and consequently the removal of the brain more difficult, it obviates the ugly ridge on the brow so liable to be made by the inexperienced. It is necessary too that the angles be well sawed through and carefully broken, because if spicules of bone remain the brain may be caught and injured during its removal. (For this method of opening see Figs. 123 to 126 inclusive.)



In the French method of opening the adult skull with a hammer,<sup>1</sup> the anterior and posterior flaps are made in the usual manner. A line one centimetre above the soft tissue is drawn around the skull with a soft pencil or with ink, the temporal muscles being cut through with a knife; by means of blows with the hammer the skull is then fractured along this line. The sound tells you when the bone is fractured and warns you to proceed to a new place. (Fig. 127.) This method is



FIG. 127.—French method of opening the skull. (After Letulle.)

much employed in France, and in the hands of experienced operators gives good results, though it is most difficult of performing in the region of the exterior occipital protuberance. It must not be used in children, in cases of fractures, bone lesions, etc. The dura is opened along the circular incision, or, more frequently, crucial incisions are made on either side of the longitudinal sinus and each side is incised by a perpendicular cut running from the vertex down to the upper margin of the bone. The four pieces are then turned down and the falx cerebri is cut anteriorly just behind the crista galli and with a portion

<sup>1</sup> J. DEJERINE, *Anatomie des centres nerveux*, 1895, p. 13.

of dura on each side of the longitudinal sinus pulled backward. It will be seen that the dura mater thus covers the sawed portions of the bones (Fig. 128) and affords a protection to the hands in the subsequent removal of the brain. Aseptic compresses may also be used for a similar purpose.

The thickness of the skull is next noted. It varies much, being usually greater in negroes and, at times, in syphilitic subjects. It also varies in different parts of the same skull, being thinnest in the temporal region and thickest at the occiput, and is often unequal




FIG. 128.—French method of opening the dura. (After Letulle.)

in corresponding points of the opposite sides. The diploe may be entirely absent in some places, in which case the bone-dust will lack the reddish color commonly observed in recently sawed bone. The skull is usually from two to six millimetres thick. In rare cases the frontal sinus may extend high up and be of unusual thickness; in one of my subjects it measured half an inch across at the top after removal of the calvarium in the usual manner. Note the relations of the external table, internal table, and diploe. Pay especial attention to the amount of blood in the latter; if abundant, suspect fracture. At

times it is entirely bloodless. The skullcap should be held up to the light so that any inequality in its thickness may be perceived. The Pacchionian granulations often give rise to small nodular depressions in the inner table, which are of course perfectly normal and should not be mistaken for pressure atrophy. They sometimes cause perforation of the bone, or permit of the passage of an external infection into the interior of the skull.

The grooves of the middle meningeal artery must be looked for on each side. In one of my cases of acromegaly the inner table resembled worm-eaten wood: the bone was soft and pliable and offered no resistance to the saw. It is necessary to be familiar with the normal yellowish-gray color of the inner table in order that changes in it may be readily detected. Whenever blood is found between the inner table and the dura, careful search must be made not only in its vicinity but also on the opposite side for a fracture by *contrecoup*. In the examination of the dura mater note its thickness, the degree of distention, its color, which is normally gray and never very red, and the amount of blood contained within it. As all liquid naturally gravitates downward, those portions of the dura which cover the most dependent parts will be most distended, unless, as often happens, an injury of this membrane has allowed the fluid to escape.

The arteries lie between the two veins. The larger arteries usually contain more blood than the veins. The dura is supplied with but few capillaries and these rarely become inflamed.

In the examination of the outer surface of the dura mater (Fig. 129) note alterations in color and gloss. The latter is often lost in consequence of tumors, hemorrhage, hydrocs, abscess, and other conditions that cause increase of intracranial pressure. Search for hemorrhages (which at times are profuse and depress the brain) and their points of origin, Pacchionian bodies (which must not be mistaken for tubercles), bulging tumors, and external pachymeningitis (ossified, purulent, syphilitic, or tuberculous), etc. The degree of tension due to fluid, etc., may be determined by puncturing or by pinching up the dura.

The brain may be exposed, but not dissected, before the heart is incised, as the quantity of blood in the cerebrum may be modified by venous oozing during the examination of the thorax. If the brain is to be injected, it is best not to remove the dura, as by its detachment usually some of the veins entering the longitudinal sinus are torn, and





FIG. 130.—Appearance of the brain after removal of the dura, which has been left attached at its posterior extremity.



FIG. 131.—Method of removing the brain after it is severed from the body.



FIG. 132.—Dissection of the brain; commencement of initial incision.



FIG. 133.—End of initial incision.





FIG. 134.—Exposure of the central portions of the brain.



FIG. 135.—Method of removing the cerebellar lobes from the pons Varoli and the medulla oblongata.

this permits the escape of the injecting fluid when under pressure. It has been shown that this operation can be performed without external disfigurement while the brain is *in situ* by forcing the fluid through a cannula introduced by way of the nostrils or the orbits.

The longitudinal sinus is opened throughout its entire length with a pair of probe-pointed scissors, and the condition and quantity of the contained blood are noted.

The dura is divided parallel with and slightly above the sawed edge of the skull, with a pair of blunt-pointed scissors, which may be introduced through a chance nick made by the saw or through an opening made with a knife for the purpose. The incision is carried completely around the skull except at the poles of its anteroposterior diameter, where it is necessary to sever the falx cerebri. The arachnoid surface of the two lateral flaps of the dura may be examined by reflecting them to one side. The character of the blood in the membranes of the brain and in its cortex, the fluid in the subarachnoid space, the character of the sulci and convolutions, and the presence of lymph are all to be noted.

To detach the falx grasp both folds of the frontal dura with the left hand, and with the right insinuate the blade of a knife along the outer face of the left fold of the dura to its attachment to the ethmoid bone. This is severed by turning the cutting edge of the blade inward towards the falx and detaching it along the line of its insertion from before backward, as near the crista galli as possible without injury to the olfactory bulbs. As the knife reaches to the anterior genu of the corpus callosum, the index-finger may be gently introduced into the longitudinal fissure so that a view may be had of the portion to be cut. It is no unusual thing to leave behind a thin strip of the dura just above the corpus callosum, a mistake which may cause annoyance to the operator or injury to the brain during its removal.

The dura may now be drawn backward and cut off posteriorly or left *in situ* in order to protect the hands of the operator and the brain in its removal (Fig. 130). The portion of the pia mater dipping down to the genu and splenium of the corpus callosum may be detached with forceps, and that overlying the surface of the cerebrum with the fingers. The handling of this delicate membrane can be greatly facilitated by allowing a stream of water to flow gently over it during its removal. The pia is colorless when normal, but may be gray or grayish white when thickened, yellow when pus is present, or red from hyperæmia or hemorrhage.

The anterior extremities of the frontal lobes are gently raised with the tips of the fingers of the left hand, and any remaining shreds of dura are severed to prevent injury to the cerebral tissue in the frontal region or corpus callosum. With the handle of a scalpel the olfactory bulbs are now shelled from the grooves in the cribriform plate of the ethmoid bone in which they lie, and the entire brain is gently turned outward while supported by the left hand. The various nerves and vessels are divided, as near as possible to their respective foramina, with a sharp, narrow-pointed scalpel, always cutting towards the bone. The ophthalmic artery and optic nerve are now severed close to the optic foramen, first on one side then on the other. Next the dura enclosing the pituitary body is cut with a sharp knife near to the bone (sella turcica) at all points except posteriorly near the infundibulum, great care being taken not to injure the delicate hypophysis, which then may be shelled out and the remaining portion of the dura behind be excised with scissors. The internal carotids are cut long, especially if the brain is to be injected. Next cut the common motor oculi, the trigeminal, external motor oculi, facial, auditory, hypoglossal, glosso-pharyngeal, pneumogastric, and as the temporosphenoidal lobe leaves the middle fossa of the skull, the tentorium cerebelli is divided with blunt-pointed scissors, or a knife with a broad flat back made especially for this purpose, along the superior border of the petrous portion of the temporal bone, preferably passing from the median line towards the sides. In making this incision care must be taken not to injure the cerebellum.

The brain mass being now supported on the left hand, cut the cord as low down as possible by a transverse incision. Pick's myelotome is a very convenient instrument for this purpose. Orth thrusts the knife through the centre of the cord and severs first one side and then the other. Any attachments of the spinal cord, medulla, and vertebral arteries can readily be loosened by introducing the forefinger into the cavity of the spinal column and through the foramen magnum. Of course, if the cord has already been removed, it remains only to cut the vertebral vessels.

The brain is now entirely free, but the cerebellum still remains in the posterior fossa, from which it is best removed by holding it firmly to the cerebrum with the fingers of the right hand and turning the brain first to one side and then to the other (Fig. 131). The brain with its pia ~~and still~~ attached is now weighed. A towel pre-



viously rolled up into the form of a turban makes an excellent temporary resting-place for the inverted brain.

During this entire procedure, which has taken longer to describe than it does to perform, the secant has been searching the exposed parts for any lesions or abnormalities, as their presence may modify subsequent processes.

Examine the external surface of the brain, the adherence of the pia-arachnoid being tested in several places, not forgetting the fourth ventricle, the circle of Willis, and the course of the middle cerebral artery lying in the fissure of Sylvius. With the latter the island of Reil and the retroinsular convolutions are also exposed.

*Quick, but not Accurate, Methods.*—Some operators do not even take the trouble to remove the brain from the skull, but merely make a number of transverse incisions across the cerebral structures. This method is only mentioned to be condemned, though it may diagnose a hemorrhage, a tumor, or an abscess.

In the coroner's work it is often necessary to make a diagnosis between heart-disease and apoplexy, when, because of baldness of the individual or for lack of time, it is impracticable to open the head. In such cases I have found it feasible to trephine just above the ear and from this point tap the ventricles and other situations liable to be the seat of hemorrhage, using an instrument resembling an apple-corer to remove brain substance for examination, though enough clotted blood may be brought out attached to a long thin brain-knife passed into the places where hemorrhage usually occurs—*i.e.*, the ventricles and the cerebellar lobes—for the purpose of establishing a probable diagnosis.

**EXAMINATION OF THE BASE OF THE SKULL.**—The base of the skull and its sinuses are next to be examined. Study the dura at its base for (1) inflammation resulting from fracture or caries, (2) tubercles, (3) gummata, (4) thrombosis of lateral sinus, (5) pachymeningitis and leptomeningitis, and (6) tumors. A fracture may be hid by the dura, but its situation will usually be shown by the presence of hemorrhage. The dura must be stripped off, though this often consumes considerable time, so that the surface of the bone may be exposed. Unless this is done, a linear fracture—one near the foramen magnum, for example—might easily be overlooked. Special examinations should now be made of the orbit, internal ear, and nasopharyngeal cavities.

## INTERNAL EXAMINATION OF THE BRAIN.

The brain may be sectioned either immediately upon its removal or after first being hardened, each method having its advantages. If an immediate diagnosis is required or colleagues are present to give unusual interest to a discussion of the findings, the sectioning will probably be done at once. If any hemorrhagic lesion is suspected, it is more conspicuous in the recent state, and a wholly unexpected bacteriologic investigation might be demanded by the revelations of the incisions. If none of these considerations prevail, the brain is hardened in a medium which will not interfere with any microscopic work that may be desired after the sectioning. Since hardening in certain fluids is necessary for certain stains and entirely precludes others, we must first of all decide what staining methods will be used before a choice of hardening fluids can be made. A two and one-half per cent. solution of bichromate of potassium or Müller's fluid will develop color contrasts between the white and gray matter and furnish material for Weigert and Golgi work, but the later methods for ganglion cells and neuroglia are precluded. Formalin is suitable for all special staining methods, including Nissl's, though the best results are obtained when the tissues are hardened in alcohol.

The brain may be hardened entire in a ten per cent. solution of formalin in a week or ten days and be suited for general topographic work. For finer histologic methods the parts should be serially incised, the sections being not more than three millimetres thick and remaining *in situ*, or, if the material to be studied is not superficial, the brain may be incised according to the methods herein to be given and then hardened. The advantages of hardening the brain in most pathologic cases are so obvious that they do not require mention. It should always be done unless contra-indicated, and when the fresh brain is sectioned and examined, the incisions should be so made that all the segments will fold together like the leaves of a book,—uninjured, undisturbed in their structural relationship, and fit for the most exhaustive microscopic examination.

Whether the brain is sectioned first or after hardening, the choice of a method will be somewhat determined by the situation of the lesion and the desire to preserve intact all its structural relations. Morbid changes in the cortex which we might wish to trace down through the internal capsule would be studied only with the greatest

difficulty after sectioning by Meynert's method, whereas if the lesions were bulbar or situated anywhere in the brain-axis this method would be very advantageous, since it permits of examining the whole of the brain-axis by serial sections.

The centrum ovale is well studied by Pitres's method, but future microscopic investigation is impossible. The same is true of Nothnagel's method, and to examine lesions of the internal capsule we must have horizontal sections. For exposing suspected or unsuspected lesions, for gaining a good idea of the general condition of the brain, and for ease and rapidity of routine work, probably no method is more useful than that of Virchow. Unfortunately, it does not favor microscopic examination and therefore is rather sweepingly condemned by some authors.

Dejerine makes a special effort so to section the brain that it may be sufficiently exposed without in any way interfering with future investigation.

**VIRCHOW'S METHOD.**—A long, sharp knife should be used in the dissection, which should be kept clean and moist by frequent washing, so that the cut surfaces will be even and smooth. A dull knife tears the brain substance more or less, thus distorting the delicate structures. Virchow insisted strongly upon the necessity of a long, clean, smooth incision being made at one stroke, and said that he would rather have a wrong incision rightly made than a right incision wrongly performed.

The brain is placed on its base with its occipital lobe towards the operator. Laying the left hand upon the left hemisphere, with the thumb in the longitudinal fissure and the fingers upon the convexity, raise this hemisphere slightly and pull it away from the median line so as to expose the corpus callosum. Insert the point of a thin narrow knife into the roof of the lateral ventricle, which lies immediately below the corpus callosum, well forward and two or three millimetres externally to the median raphe of the corpus callosum (Fig. 132). Make a concave incision—concavity directed outward—through the roof back to the posterior cornu, being careful not to injure the floor of the lateral ventricle. Note the character and quantity of fluid present, which normally is perfectly clear and about three cubic centimetres in amount. Connect the two extremities of the first incision by a second and third incision meeting at an angle of 45 degrees just outside the basal ganglia. In this manner the greater portion of



the cerebral cortex on the left side will be removed away from the basal ganglia for future sectioning (Fig. 133). The right hemisphere may be turned half around and sectioned in the same way.

The knife is then introduced into the foramen of Monro and the anterior fornix is brought forward, exposing the vena interposita and the choroid plexuses, which with the body of the fornix are carried back, thus exposing the third ventricle (Fig. 134). Then examine the corpus fimbriatum, the lyra, the anterior, posterior, and middle commissures, the corpora quadrigemina, the pineal body, and the commencement and lumen of the *iter a tertio ad quartum ventriculum*. The pineal gland is often infiltrated with salts, as may readily be determined after sectioning by rubbing a small portion of it between the thumb and index-finger. If it be desired to examine the fifth ventricle, an incision is made directly in the median line into the septum lucidum, parallel to the corpus callosum, the anterior fornix being elevated by the left hand and thus put on a stretch.

The crura are then severed by transverse incisions joining at about a right angle in the median line. The cerebellum, the medulla oblongata, and the pons Varolii are next to be removed. This may be done at the start, if preferred, reversing the order here given as to the other parts of the brain. After examining for dilated veins, tumors, and cysticerci, transverse incisions are made in the cerebellum on one side through the centre of the arbor vitæ, and then on the other side. The cerebellum may, however, be removed before these incisions are made by severing the medulla oblongata and the pons Varolii and dividing the cerebellar hemispheres in the median line into two parts. The pons, the medulla, and the commencement of the spinal cord may now be cut transversely by incisions one-fourth to three-eighths of an inch apart, and all pathologic changes carefully noted, but these portions are preferably hardened previous to examination, which is best accomplished by the preparation of serial sections. (Figs. 135 to 139 inclusive.)

Both Nauwerck and Orth, before making transverse sections of the pons and medulla, fold the sections of the brain together as you would the pages of a book in order that it may be turned. Then, pushing the fingers of the left hand under the pons and medulla, the transverse cuts may be made. In case of tumors or metastatic conditions simpler methods may be used; thus, only one longitudinal or one transverse section may be made through the diseased as well as the healthy tissue, while the arachnoid is left intact.



FIG. 136.—Segmented brain. The central portion has been divided into two parts. The cerebellum, pons Varoli, and medulla have been everted with the right and left cortical portions.



FIG. 137.—Method of sectioning the cerebellum.



FIG. 138.—The whole brain after it has been sectioned.






FIG. 139.—Section of the brain. The lines and arrows show the position and direction of the various incisions. (After Nauwerck.)




FIG. 140.—Basal ganglia, with cerebellum, pons Varolii, and medulla oblongata attached, in Meynert's method of dissecting the brain. The twelve cranial nerves are shown. C, cerebellum; F, flocculus; M, medulla; P.V., pons Varolii; T.L., temporal lobe; F.L., frontal lobe; P., peduncles; C.a., corpora albicantes; C.c., central commissura; R.I.C., retrosular convolution; O.C., optic commissura; P.S., posterior roots of olfactory nerve; I, insula. (After Dejerine.)




FIG. 141.—Sectioning of the brain. *AB*, incision practised by Flechsig; *CD*, that of Brissaud; *EF*, that of Dejerine. The hemisphere to be incised is placed on its external surface, the occipital lobe towards the operator in case of the left hemisphere, and the frontal lobe for the right hemisphere. (After Dejerine.)




FIG. 142.—Incisions made by Dejerine in a case of cortical lesion previous to hardening.



**MEYNERT'S METHOD, SLIGHTLY MODIFIED BY BLACKBURN.**—The brain is placed with its base upward and the cerebellar end towards the operator. The cerebellum is elevated and the pia mater cut through above the corpora quadrigemina, around the crura, and along the inner margins of the temporal lobes until the middle cerebral arteries are reached. The Sylvian fissures are opened to their entire extent, the opercula are raised, and the insular lobes exposed to their limiting furrows.

The apices of the temporal lobes are now elevated, and, with the knife held nearly horizontal, their junction with the base is cut through until the anterior extremities of the descending cornua are opened. The knife is inserted in the descending horn, and the incision is carried backward as far as the posterior angle of the insula, or even some distance beyond it, severing some of the convolutions at the posterior extremity of the Sylvian fissure.

The next incision is made to separate the basal piece from the posterior extremities of the frontal lobes. It connects the anterior boundaries of the islands and opens the anterior horns of the ventricles. The incision may be a slightly curved, transverse one, connecting the anterior border of the islands; or, by a little care and a double crescentic cut, the exact boundaries of the convolutions may be followed.

The cerebellum is now raised, the knife entered at the posterior angle of the island, and the incision carried along the outer limiting furrow until it meets the cut previously made through the anterior border. Care must be taken to keep the knife in the angle between the roof of the ventricle and the basal ganglia, to avoid injuring the latter. The basal piece is now lifted until the anterior crura of the fornix and the septum lucidum may be severed, and the basal section thereby completed.

The basal piece thus separated includes the island of Reil, the basal ganglia, the crura, pons, medulla, and cerebellum. (Fig. 140.)

**PITRE'S METHOD.**—The lateral ventricles are exposed as in Virchow's method. The hemisphere lies on its under surface and a series of six transverse vertical sections are made parallel to the fissure of Rolando. These are called the prefrontal, pediculo-frontal, frontal, parietal, pediculo-parietal, and occipital. Pitre's method is very useful for localizing lesions in the centrum ovale, but not at all adapted to studying the internal capsule nor for subsequent microscopic work. The same is true of the closely similar method of Nothnagel. Flech-





FIG. 137.—Method of sectioning the cerebellum.



FIG. 138.—The whole brain after it has been sectioned.




FIG. 139.—Section of the brain. The lines and arrows show the position and direction of the various incisions. (After Nauwerck.)




FIG. 140.—Basal ganglia, with cerebellum, pons Varolii, and medulla oblongata attached, in Meynert's method of dissecting the brain. The twelve cranial nerves are shown. C, cerebellum; F, flocculus; M, medulla; P.V., pons Varolii; T.L., temporal lobe; F.L., frontal lobe; P, peduncles; Ca., corpora albicantes; C.c., central commissura; F.I.C., fronto-insular convolution; O.C., optic commissura; P.S., posterior roots of olfactory nerve; I, insula. (After Dejerine.)

When certain that all attachments are secure, the stopcock is gradually opened, allowing the tubes to become filled and the fluid to percolate slowly through the brain. Care should be taken that the cannulas do not bend the arteries short upon themselves, thus occluding their lumina. The first fluid which passes through will be mixed with blood and should not be used again, but when it has become clear it may be used over and over. It usually runs through very quickly, and the tank should be refilled at least every day for the first week, and oftener if convenient. The brain should always be in an excess of the fluid and a vessel provided for the overflow. For refilling the tank it is best to draw some of the liquid out of the jar with a siphon, which will not disturb the brain or the position of the cannulas.

A week or two will suffice in urgent cases, but the longer the brain remains in the fluid the better will be the hardening. Some of my most beautiful specimens are those which were kept in Müller's fluid for five or six months. Haste and thoroughness are incompatible in this process. No padding should be used to keep the organ in position, the best and surest agent for this purpose being a plentiful excess of the liquid and an occasional change in its position.

If it seems unnecessary to inject the vessels, the following method may more easily be carried out and gives most excellent results. An open jar, bucket, or wash-basin is one-quarter filled with absorbent cotton, and Müller's fluid—to which one per cent. of formalin may be added with great benefit—is poured in until the vessel is about half full. The brain, after being weighed, is carefully placed in the centre of the vessel and more fluid is added until the organ is well covered, when it is placed in a refrigerator. If this be done, even though the arteries have not been injected nor any incision made into the ventricles, there is no danger that the brain will decompose, even in summer. On the next day the position of the brain is altered and the fluid changed. The renewal of the fluid can best be accomplished with a siphon, only a part of it being removed at one time.

The fluid is changed again on the third day, then every other day for three times, twice a week for the next three weeks, and once a week for the final three weeks. Remember that the jar is uncovered, and this allows of the evaporation of the fluid and possible spoiling of the specimen. The brain can then be thoroughly washed and put in 80 per cent. alcohol, or the Müller's fluid can after the fifth or sixth week be diluted with one-fifth alcohol, then with one-quarter, one-



third, one-half, and finally three-quarters alcohol, where the brain can be kept for several months until it is transferred to the alcohol of 80 per cent. strength. Instead of Müller's fluid a 2.5 per cent. solution of bichromate of potassium may be employed. It is important to remember that nervous tissue preserved for the purpose of study by the Nissl method should not be placed in Müller's fluid, but in alcohol or formalin. About two thousand cubic centimetres of a 10 per cent. formalin solution are used and changed every third day. The solution should be kept in a tightly closed jar to prevent the escape of the formalin.

GIACOMINI'S METHOD.<sup>1</sup>—This is well adapted for the macroscopic study of the brain, but, on account of the zinc chlorid used, the tissue is rendered unfit for microscopic work. If the specimen is a brain tumor, a small portion of it may be placed in a hardening fluid for microscopic study and the remainder then treated by this process.

The brain, in as fresh a state as possible, is put into the *Liquor zinci chloridi* (U. S. P.). It will be found to float at first and should be turned several times the first day. On the second day the pia and arachnoid, which until now have been useful in keeping the brain intact, are removed while the organ is under water or floating in the fluid; if allowed to remain longer, they become so adherent to the cortex as to be separated with difficulty and more or less damage to the cortical substance. The brain is left in the fluid for from six to ten days, then removed, well washed with water, and put in 95 per cent. alcohol for ten days or two weeks and next in glycerin for another ten days or more. After this it is placed in absorbent cotton and exposed to the air in a dark place free from dust. Any exudation should be carefully removed, and when no more appears (which may be in from several weeks to as many months) the surface is to be well coated with the best mastic varnish applied with a soft camel's-hair brush. To prevent flattening of the surface upon which it rests, it must be well packed in absorbent cotton and its position frequently changed.

KAISERLING METHOD.—See page 343 for the preparation of brains with the object of preserving their natural coloration.

---

<sup>1</sup> *Gior. di r. Accad. di med. di Torino*, 1883.

## CHAPTER XIV

### THE SPINAL CANAL AND CORD

THE spinal cord may be removed either anteriorly or posteriorly,—*i.e.*, by excising the bodies of the vertebræ through the thorax and abdomen freed from their viscera or by severing the laminæ and spinous processes of the vertebræ through an incision posteriorly. The latter route is decidedly the more convenient and is used whenever possible. Generally it is best to remove the cord before the abdomen is opened, this being a much cleaner operation, an important factor in private practice.

The cadaver is placed prone upon the table close to the side at which the operator stands, with the head hanging over the end or, better, with a block under the chest and neck and, if desired, one under the lumbar region. Beginning at the external occipital protuberance, an incision is carried along the spinous processes to below the fourth lumbar vertebra, dividing all the tissues down to the bone. (Figs. 143, *A B*, and 144.) The incision is made low in order to allow for room for future manipulations, as the tissues here are thick and the future sawing is to be done in a hollow, as it were. The superficial and deep structures are then dissected from the bones, exposing the vertebral groove on either side of the spinous processes. Or, after incising the skin over the spinous processes, insert the knife, with its back down, at the lower end of the incision and cut upward along the column, keeping the blade pressed against the spinous processes. In this way the fibrous attachments are cut close and the vertebral groove is clean and free from troublesome soft tissues. The soft parts should be very thoroughly removed, as they would interfere considerably with the subsequent sawing. This can be quite well done by scraping with a chisel or an old knife.

In cases of luxation, fracture, Pott's disease, etc., it may be desirable to remove portions of the vertebral column *en masse*. This can readily be done by the proper use of a saw after severing the intervertebral cartilages above and below the lesion. The space is then filled by inserting a stick and pouring plaster upon it.

The canal is easily opened with Luer's rhachiotome, an adjustable,




FIG. 143.—Lines for removing the spinal cord and the brain, the latter through a small triangular occipital incision. *AB*, initial incision for removal of the cord; *CD*, curved incision for the purpose of avoiding division of the skin above the dressed portion of the body; *EAF*, angular incision in the occipital bone through which to remove the brain without elsewhere opening the skull.







FIG. 144.—Position of the body in removal of the spinal cord. The primary incision is being made.



FIG. 145.—Removal of spinal cord. The primary incision has been made and the vertebral column freed from muscle, fascia, etc. The angle at which the saw should be held is well shown.



FIG. 146.—Removal of spinal cord. The sawing having been completed at a convenient point in the dorsal vertebrae, the sawed bone is elevated with the fingers or, if strongly attached, with the forceps.




FIG. 147.—Removal of spinal cord. The spinal canal is well shown with the dura mater still intact. The sawed portions of bone both above and below are left attached.





FIG. 148.—Removal of spinal cord. The atlas and the axis are being severed with the cordome.



FIG. 149.—Removal of the spinal cord with its dura. The lower end is held with the left hand while the nerves are being severed. FIGS. 145 to 149, inclusive, are from photographs taken directly from above.



FIG. 132.—Appearance of the parts after wiring of the lateral halves in Harke's method of examining the nasopharynx. Removal of a triangular portion of the occipital bone.



FIG. 131.—Harke's method of examining the nasopharynx. Appearance of the parts after the wiring has been completed and the lateral halves have been pried apart. The tip of the epiglottis is plainly seen in the photograph. A mirror may be used to illuminate these parts in order to bring them more perfectly into view.



double-bladed saw devised for the purpose (Fig. 25). It does the work more quickly, but has the serious fault that it is liable to become impacted and injure the cord in its release. The same object may be accomplished with a single-bladed saw having curved ends (Fig. 21). The lamina should be sawed close to the transverse process, with the saw teeth held away from the spine at an angle of about thirty degrees (Fig. 145). Unless this direction is taken there is some danger that the canal will be missed or that the blade may enter it and the cord be injured. Orth calls attention to the fact that one can tell when sufficient sawing has been done by the mobility of the spinous processes. Other instruments which may be used are the double chisel of Esquirol, the knife-shaped chisel of Brunetti, and the rhachiotome and hammer of Amussat, the latter being much preferred in France to Luer's rhachiotome, which is not approved of. If the rhachiotome is used as the seventh cervical vertebra is approached, both from above and below, the incisions are made more and more towards the side, as the canal is wider here, owing to the increased size of the cord at this spot. After the canal has been opened in the dorsal region with the saw, a pair of bone-nippers is used to pry up the portions of vertebra thus loosened, and the dura is exposed (Figs. 146, 147). The sawing can then be continued in both directions until the entire canal is opened, except the atlas and axis, which had better be cut with bone-forceps (Fig. 148). In using either bone-forceps or pliers be very careful not to produce artefacts of the cord. The cord at the first dorsal vertebra is then tied with a string, so as to have the situation accurately determined, or the first dorsal nerve may be dissected out and left attached to the cord.

The spinal cord covered with its membranes may now be studied *in situ*, after which the dura and the spinal nerves are divided *below* the cauda equina. The dura being elevated with the fingers or forceps and pushed to one or the other side, the spinal nerves are cut, with a long, thin, narrow-pointed, sharp knife, close to their points of entrance into the intervertebral foramina. (Fig. 149.) The dura at the foramen magnum can best be severed from the bony margin above after the brain has been removed. The cord may be taken away with the brain attached if so desired. The spinal ganglia may be extracted with the nerves and cord by cutting away the articular processes and gently pulling the cord, by the dura, to the opposite side and severing the nerve as far in the foramen as possible.



By making a median incision in the dura mater the cord is exposed, and can, of course, be removed. This procedure, however, is more liable to cause injury to the cord than the method given above.

After freeing all points of attachment the cord must be very gently transferred to the table or tray for further examination. Study the dura for (1) thickness, (2) color, (3) blood, the cerebrospinal fluid for (1) pus, (2) blood, and the pia for (1) expansion, (2) thickness, (3) contained blood, and (4) color. Gentle palpation may reveal areas of softening or sclerosis. The further manipulation of the part will depend upon the extent of the examination required. If the cord is to be preserved for future study, the dura is opened in the median line throughout its entire extent, the blade being inserted at the lower end, and transverse incisions about one inch apart down to the pia are made in the cord. It may be hardened at the same time and in the same jar as the brain by curling it around that organ; but it is better to suspend it by the dura, with a small weight attached, in a long jar, or it may be kept in such a jar lying upon its side. In summer the jar should be placed in the refrigerator.

If the examination is to be completed immediately, the cord is laid out on the table, with its anterior surface resting preferably on a towel or piece of cheese-cloth, and the dura opened throughout its entire length as already directed. Note is made of the conditions observed. Much valuable information can be obtained by the macroscopic examination, especially if a hand-glass be used and diagrams made at the time. Then, with a sharp, thin knife, which should be moistened with water after several incisions, transverse sections about an inch apart are made through the cord and membranes; the under surface of the dura, however, is left uncut, in order that the cord may be replaced in its entirety. A careful operator may hang the cord over the index-finger of the left hand, keeping it in place with the thumb, and make the incisions there, the dura being sufficient to protect the finger. A microtome knife is admirably adapted for making the incisions. Areas of softening should not be incised, because of the inevitable disturbance thus produced in the relations of component parts. Froriep's incision of the spinal cord, one long longitudinal incision throughout the entire extent of the cord, is severely criticised by Virchow.

Where the avoidance of disfigurement above the parts covered by clothing is a matter of great importance, sufficient room for opening the cervical canal can be obtained by making a crescentic incision from

the centre of one shoulder to the other, with the concavity towards the head, and dissecting up the skin. (Fig. 143, *C D*.)

Sometimes it is advantageous to open the canal by removing the vertebral bodies through the long anterior incision with the body resting on its back. Brunetti's chisels were devised for this purpose. After removal of the thoracic and abdominal viscera, the pointed guard is inserted in the vertebral canal, and the instrument, held parallel with the long axis of the spinal column, is driven forward with a mallet, thus severing the pedicles and removing the bodies or anterior wall. By this method the spinal ganglia are said to be rendered more easily accessible. The remaining steps are about the same as those described for the posterior incision.

## CHAPTER XV

## CONCLUSIONS OF THE BRAIN AND SPINAL CORD

BRAN.—There is a circumscribed collection of soft, white, granular substance, with or without purulent membranes, and containing organisms.—e.g., *Staphylococcus pyogenes*, *Streptococcus pneumoniae*, gonorrheal and cerebrospinal meningitis, the bacillus of typhoid, influenza, etc.

(b) Extension of disease from the middle ear or mastoid bones. (c) Septic emboli from distant foci.

(d) Ulcerative endocarditis, purulent bronchitis, etc. (e) Actinomycosis and other mycotic collection. (f) Primary, pure or secondary pyogenic from extension or metastatic).

Color of an abscess is usually greenish-yellow, usually in the sphenoid-sphenoidal cleft, but especially in middle-ear discharges. Pus vessels are minute, and are very delicate, dense and firm, and are superficial or subcutaneous, from three to five millimeters in diameter, and may have a yellowish tint, undergo fatty degeneration.

[illegible]



one of my cases there was found after death a sarcoma of the pituitary body; in another, all of the glands of the body appeared to be hypertrophied. I have removed *post mortem* the pituitary body through the orbit. Under acromegaly may also be classed OSTEITIS DEFORMANS, an affection which causes softening and distortion of the long bones of the body; HYPERTROPHIC PULMONARY OSTEO-ARTHROPATHY, where there is antecedent lung disease and the bones of the skull are not involved, and LEONTIASIS OSSEA, an overgrowth of the bones of the cranium. In MICROMEALY the condition is the reverse of that found in acromegaly.

ANÆMIA CEREBRI.—A condition in which the brain is temporarily or permanently deprived of part of its blood-supply. Due to: (a) Mechanical obstruction to the circulation,—*e.g.*, valvular heart-lesions, thrombosis, embolism, or ligation of a vessel. (b) Hemorrhage. *Classification*.—(a) General or local. (b) Acute, subacute, or chronic. (c) Partial or complete. The membranes are pale; small arteries over the gyri are empty, though large veins are full. The brain substance is anæmic, the surface moist, few puncta vasculosa are seen, and the cerebrospinal fluid is increased.

ANEURISM OF CEREBRAL ARTERIES.—*Classification*.—(a) Single or multiple. (b) Large or minute. *Seats*.—(a) Most frequent in branches of the middle cerebral artery, especially those of anterior perforated spaces. (b) May be cortical. The aneurisms are usually very small, varying in size from that of a pea to a cherry-stone (seldom larger), multiple, and may resemble bunches of grapes. If hemorrhage occurs in basal aneurisms, the internal capsule and basal ganglia are injured, the lesion usually being extensive. On the cortex the result of hemorrhage is much less grave.

APOPLEXIA NEONATORUM.—A form of hemorrhage of the brain occurring in the new-born, usually the result of traumatism. (a) Accidents during labor, from forceps, etc. (b) Congenital defects in blood-vessels, brain, or skull. (c) May result from prolonged and severe normal labor. *Seats*.—(a) Meninges (piaarachnoid) most frequently, often bilateral, and usually at the base. (b) May be between dura mater and skull; is accompanied by cephalæmatoma. (c) May occupy the ventricles. (d) May occur in brain substance about basal ganglia. (e) Sometimes found in parietal region and Sylvian fissure. (1) Generally the hemorrhage is meningeal primarily, producing brain-lesions secondarily, such as atrophy and softening, by pressure.

(2) Cortical hemorrhage is represented by a clot, which may be encysted, softened, or organized, causing more or less injury to the brain. (3) When the hemorrhage is between dura and skull, fracture is said to be always present.

ATAXIA, HEREDITARY (FRIEDREICH'S).—A form of ataxic paraplegia occurring especially in young children. (a) The disease is sometimes hereditary, and it is not uncommon to have several members of the same family affected. (b) More frequent in males than in females. (c) A specific lesion of the cord. (1) There is a gliosis of the posterior column of the spinal cord, due to developmental errors (Osler). (2) Talipes equinus occurs in both feet. (3) Lateral curvature is common.

ATAXIA, LOCOMOTOR (TABES DORSALIS).—A chronic disease of the nervous system, characterized by sclerosis of the cord and brain, and by incoördination, with motor, sensory, and trophic disturbances. (a) Male sex. (b) Adult life. (c) Syphilis. (d) Wet and cold. (e) Sexual excesses, etc. (1) *Spinal Cord*.—Externally the meninges are thickened and adherent. Posterior roots are atrophic and of a grayish tint. Internally sclerosis of the cord begins in the posterior-root zone, involving the outer layers of posterior columns in the lumbar region. The sclerosis gradually extends inward, involving successively the columns of Burdach and Goll; when the process reaches the upper dorsal region, it is confined to the column of Goll. The cord presents a flattened appearance posteriorly, the sides being somewhat contracted. The diseased areas are firm, grayish or grayish red in color, and the whole cord is often firmer in consistency. (2) *Brain*.—Changes of less consequence than in the cord may be sclerosis in restiform bodies, inferior peduncles of cerebellum, and certain cranial nerves,—the oculomotor, optic, and auditory. Atrophy of the optic nerve and hemiplegia may occur. Some recent writers consider paralytic dementia to be such a disease of the brain as locomotor ataxia is of the cord. (3) Peripheral nerves may show degeneration or even neuritis. (4) In later stages occur dermopathies and arthropathies,—*e.g.*, perforating ulcer of foot, herpes, etc., Charcot's joint, etc. There may be evidences of loss of control of sphincters. The essential lesion is a dystrophy attacking the peripheral sensory neuron. Erb<sup>1</sup> has found out of a total of 1100 cases of tabes that there was

---

<sup>1</sup> *Berliner klinische Wochenschrift*, 1904, vol. xli, nos. 1, 2, 3, and 4.

an unmistakable history of syphilis or chancre in 89.45 per cent., syphilitic antecedents were probable in all but 2.8 per cent., and even here syphilis is suspected by Erb. In 96 cases of *tabes* Lesser<sup>1</sup> found an aneurism in 18, and speaks of these processes as the quartan manifestations of syphilis.

**CAISSON DISEASE.**—A peculiar nervous affection, the result of a sudden reduction of atmospheric pressure. Occurs in bridge-builders, divers, etc., who, after working for hours under a pressure of two or three atmospheres, have suddenly returned to air of normal density. In fatal cases there is a marked destruction of nerve tissue in the posterior columns and the posterior portions of the lateral columns, forming fatty detritus and compound granular cells. Free gas bubbles of nitrogen are said to exist in the circulatory system of those affected.

**CHOREA, ACUTE.**—(a) Female sex. (b) Early life (before the fifteenth year). (c) Heredity. (d) Bad hygiene. (e) Fright. (f) Bad habits. No constant lesions are found. Vascular changes, usually of a congestive type, such as hyaline degeneration, leucocytic infiltration, minute hemorrhages, and thrombosis of small arteries, have been described. Possibly due to a specific organism.

**CONGENITAL ANOMALIES.**—Cranioschisis, rhachioschisis, hydro-meningocele, encephalocele, myelomeningocele, hypoplasia of different parts, as of the cerebellum, micrencephaly, hydrocephalus, internal and external porencephaly, idiocy, cretinism, micromyelia, total absence of parts, and anomalies of distribution.

**CRETINISM.**—A low form of idiocy, either congenital or acquired during the early years of life, and associated with anatomic changes in the thyroid gland, as absence, hypoplasia, atrophy, or goiter. It is endemic in certain localities, notably Switzerland, where goiter is prevalent. Heredity bears a causative relation. The condition usually appears at birth. The child is stunted and dwarfish in appearance. The trunk is large in proportion to the development of the head, hands, and feet. The head is flat, the face broad and expressionless, the eyes are dull and stupid, the nose is flat and depressed, the lips are thick, and the tongue is large and usually protrudes. The teeth are carious; the hair is thin, brittle, and harsh to the touch; the skin about the hair is dry and scurfy. The abdomen is prominent; the legs are short and

<sup>1</sup> *L. c.*, no. 4, p. 80.



thick, the hands and feet undeveloped. The skin is yellow, leathery, and rough to the touch.

**DELIRIUM, ACUTE.**—The post-mortem findings are usually negative. There may be great venous engorgement of the meninges, and the cortex and blood-vessels may show exudation and leucocytic infiltration into the lymph-spaces and sheaths. Careful examination of the lungs and ileum should be made in fatal cases.

**ENCEPHALITIS, ACUTE.**—Due to: (a) Acute infectious disease. (b) Traumatism. (c) Intoxications. The minutest foci of inflammation are not recognizable by the unaided eye; later stages have a pinkish appearance or are represented by clusters of small dark-red hemorrhagic foci. When suppuration follows, these areas take the form of yellowish-white patches whose tissue soon liquefies and becomes purulent.

**ERYTHROMELALGIA.**—In this condition there is arteriosclerotic thickening of the blood-vessels with diminution of their lumen (obliterative arteritis) and some involvement of the peripheral nerves.

**HÆMATOMYELIA.**—Hemorrhage into the cord. (a) Traumatism. (b) Exposure. (c) Convulsions. (d) Tumor. (e) Syringomyelia. (f) Myelitis. (g) Male sex. (h) Middle life. The cord is usually enlarged, occasionally lacerated. The blood is generally confined to the gray matter, but may escape beneath the membranes.

**HEMIPLEGIA IN CHILDREN.**—Causes: (a) First or second year. (b) Traumatism. (c) Embolism or thrombosis. (d) Congenital defect. *Classification.*—(a) Embolism, thrombosis, or hemorrhage. (b) Atrophy and sclerosis. (c) Porencephalon. (1) The results of embolism, thrombosis, or hemorrhage depend on the extent and rapidity of the formation and on location. When the process is an acute one and extensive, it is either immediately fatal or leads to more or less extensive destruction of the brain substance; there is a tendency to softening or suppurative change. (2) Atrophy and sclerosis may involve a group of convolutions, an entire lobe, or even a whole hemisphere. The affected gyri are firm, hard, and atrophied, contrasting sharply with the normal tissue. They may be uniform in appearance or there may be nodular projections. In porencephalon there is loss of substance, with the formation of cavities or cysts at the surface of the brain.

**HEMORRHAGE, CEREBRAL.**—The most common cause (sixty per cent.) is rupture of the lenticulostriate artery. *Classification.*—(a)

Basilar. (*b*) Cortical. In basilar hemorrhage section of the brain substance frequently shows miliary aneurisms, which are seen as small dark bodies along the course of the blood-vessels penetrating the anterior perforated spaces. Aneurism of a branch of the circle of Willis may be found. Endarteritis and periarteritis are found in the cerebral vessels. At the seat of a recent hemorrhage the brain has a dark-red, softened appearance, the tissue being reduced to a coagulated or pulpy mass of detritus. When the hemorrhage has been extensive, the remainder of the brain is anæmic. The gyri are more or less flattened, from extravasated blood, and the sulci are indistinct. Hemorrhages are most common near the corpus striatum towards the outer section of the lenticular nucleus. They may be small and limited to the lenticular body and internal capsule or may break into the lateral ventricle. Ventricular hemorrhage is rare. It is usually bilateral. Meningeal hemorrhage is usually caused by fracture of the skull or rupture of a blood-vessel. The hemorrhage may be small or large. It may be above or below the dura or between the pia and the arachnoid. The hemorrhage may be primary into the fourth ventricle.

HEMORRHAGE INTO THE SPINAL MEMBRANES.—Extrameningeal hemorrhage may be extensive, without compression of the cord. Rupture of an aneurism into the spinal canal may produce profuse and rapidly fatal loss of blood. There may be little demonstrable morbid change. Intrameningeal hemorrhage usually occurs in scattered areas as the result of acute infectious fevers. More extensive hemorrhages result from epilepsy, tetanus, and strychnine poisoning. Occasionally hemorrhage into the spinal meninges may ascend to the brain.

HYPERÆMIA, CEREBRAL.—This may be: (*a*) Active. (*b*) Passive. (1) The cerebrum is congested, the blood-vessels are somewhat distended, and petechial hemorrhages are numerous. On section the gray substance contrasts very markedly with the white; the former is of a brick-dust color; the latter shows many punctate hemorrhages. (2) In passive congestion the veins of the cortex are distended; the gray matter has a deeper color and its vessels are full. The gray matter shows distention of the smaller veins, which on section allow their contents to exude as drops of blood of various sizes. Excessive passive hyperæmia may result in cerebral œdema.

LEPTOMENINGITIS, ACUTE CEREBROSPINAL.—Acute inflammation of the pia and arachnoid of the brain and spinal cord. Causes: (*a*) Acute infectious fevers. (*b*) Injury or disease of the base of the skull.

(c) Extension of disease from nose, ear, or Eustachian tube. (d) Pyæmia. The organisms most commonly found are the meningococcus, the pneumococcus, the tubercle bacillus, and the cocci of inflammation; more rarely, the bacilli of influenza and of typhoid, the colon bacillus, and the gonococcus. *Classification*.—(a) Simple or traumatic. (b) Purulent. (c) Tuberculous. (1) In simple or purulent meningitis the membranes are thickened, the blood-vessels dilated, and there is more or less exudation, which may be serous, serofibrinous, or purulent. The exudation may be so extensive as to cover up the convolutions. The inflammatory process is most marked in the basilar portions. It may be unilateral or bilateral. In the former the condition is due to extension from neighboring parts. (2) The tuberculous form of the disease is usually cortical as well as basilar. It begins as a miliary tuberculosis, and in the early stages exudate is not extensive. The ventricles also may be involved and present considerable distention and softening; they seldom suffer in other forms of the disease.

**MENINGITIS, ACUTE CEREBROSPINAL.**—An acute infectious disease, especially of early life, characterized by inflammation of the membranes of the brain, with an exudation of fibrinopurulent material, chiefly towards the base, and due to the *Diplococcus intracellularis*. (1) *Membranes of the Brain*.—In acute fatal cases there is intense injection of the pia and arachnoid, with a little exudate. In more chronic cases there is a formation of fibrin or of pus, or of both; this is most marked at the base of the brain. The meninges are much thickened and opaque. The larger blood-vessels are overfilled and many of the smaller ones are obliterated. Sometimes the entire cortex is covered with a thick purulent exudate, and there may be much lymph along the larger fissures and in the sulci. In acute cases the ventricles are dilated, the ependymæ are inflamed, and the cavity may contain pure pus. (2) *Cranial Nerves*.—The nerves usually involved are the second, fifth, seventh, and eighth. They are often embedded in the exudate. Micro-organisms may be found in the fibrin. (3) *Brain Substance*.—This is softer than normal, has a pinkish color, with foci of hemorrhage and of brain softening. (4) *Lungs*.—Pneumonia and pleurisy may occur. The lungs are often congested, with evidences of bronchitis. (5) *Abdominal Organs*.—The liver is rarely altered. *Acute* ~~is~~ **is sometimes present**, and the intestines may show ~~it~~ **it**.—There may be rose-colored,



hyperæmic spots, resembling the typhoid rash, urticaria or pemphigus, and in rare instances gangrene. (7) *Eye*.—Neuritis is common, and there may be acute papillitis. Purulent chorioido-iritis or even keratitis sometimes occurs. (8) *Ear*.—Otitis media develops from direct extension, and frequently leads to abscesses. In one of my cases the bacillus of tuberculosis was found associated with the meningococcus. In two fatal cases examined by me there was a history of traumatism, though no sign of this was found at the postmortem. During an epidemic domestic animals, as the goat, should be watched for signs of disease.

MENINGO-ENCEPHALITIS; CHRONIC DIFFUSE OR DEEP CHRONIC LEPTOMENINGITIS.—(a) Male sex. (b) Early adult or middle life. (c) Syphilis. (d) Alcoholism. (e) Certain occupations, as those of artists, navy and army officers, etc. The membranes of the brain are thickened and opaque and more or less extensively adherent to the cortex, which is torn on attempting to remove them. The convolutions of the brain are atrophied, especially in the frontal and parietal regions. The gray matter may be obscurely outlined. The white matter is firm in consistency. The ventricles are dilated and the ependymæ granular; frequently there are areas of hemorrhage or softening associated with chronic arteriosclerosis. There is an increase in the cerebrospinal fluid. Usually sclerosis of the posterior columns, with involvement of the lateral, is found. There may be an extraordinary development in the lymph connective system of the brain, with a parallel degeneration and disappearance of the nerve-elements and the axis-cylinders, and finally shrinking and extreme atrophy of the parts involved.

MUSCULAR ATROPHY, PROGRESSIVE (SPINAL).—(a) Male sex. (b) After the thirtieth year. (1) Macroscopically there is great muscular wasting, beginning usually in the thenar and hypothenar eminences and thence extending to the general muscular system. In marked cases the subject may be reduced "to skin and bone." Deformities and contractures result and lordosis is almost always present. (2) Microscopically the muscles undergo fatty and sclerotic change and the terminal ends of the motor nerves are degenerated. (3) Examination of the cord shows the anterior roots corresponding to the diseased muscles to be atrophied. Neuroglial tissues show marked increase, most conspicuous in the anterolateral tracts. The degeneration of the gray matter extends to the medulla. Large ganglion-cells

in the motor cortex may be wasted. In a case at Elwyn which I examined *post mortem* the diaphragm was easily seen through when held up to the light.

**MYELITIS, ACUTE.**—(a) Traumatism. (b) Exposure. (c) Certain infections. (d) Disease of the spine. (e) Disease of the cord. (1) The cord is swollen and soft and the pia injected. On incision a diffuent fluid may escape. The distinction between gray and white matter is often lost. Hemorrhages are frequent. (2) Histologically the nerve-fibres are swollen, the axis-cylinders beaded, myelin droplets abundant, and corpora amylacea may be seen. The ganglion-cells are swollen, irregular in outline, and exceedingly granular and vacuolated. In the removal of the cord in these cases great care must be taken not to produce artefacts.

**MYELITIS FROM COMPRESSION.**—(a) Caries of the spine. (b) New growths. (c) Aneurism. (d) Parasites. (e) Distention of central canal with inflammatory liquid or blood. Changes appear first in the white matter, the fibres of which may within six hours swell up and disintegrate.

**POLIOMYELITIS, ACUTE ANTERIOR.**—(a) Early life. (b) Boys more susceptible than girls. (c) Acute infectious fevers. (d) Probably a specific micro-organism. (1) The seat of the lesion is in the part supplied by the anterior median branch of the anterior spinal artery. Cervical or lumbar portions of the cord are most often affected. (2) In the early stages the lesion is an acute hemorrhagic myelitis, with rapid destruction of the large ganglion-cells. (3) The nerve-fibres of the anterior roots corresponding to the ganglion-cells destroyed break down and disappear. (4) Certain anterior nerve-roots are atrophied, and the muscles innervated by them waste and become fatty and sclerotic.

**RAYNAUD'S DISEASE.**—A form of vasomotor neurosis causing local syncope, cyanosis, and symmetrical gangrene, affecting especially the fingers and toes, caused by spasm and constriction of the small blood-vessels.

**VON RECKLINGHAUSEN'S DISEASE.**—This is a general fibrosis of the peripheral nervous system. Nuthall and Billington<sup>1</sup> report the necropsy of a case.

**SCLEROSIS, INSULAR (DISSEMINATED SCLEROSIS).**—Its cause is

---

<sup>1</sup> *Lancet*, December 27, 1902, p. 1751.

not definitely known. Is more common in the young than in the old. Sclerotic areas are usually small, of a grayish or whitish color, widely distributed in the brain and cord and in the gray and white matter. They are more abundant about the ventricles, the central canal, the pons, the cerebellum, and the basal ganglia. The patches are firm, dry, and sharply defined from the surrounding tissue; in some cases they may be less firm and not so well defined. Microscopically there is a marked increase of neuroglia, the medulla of the nerves is destroyed, and the axis-cylinders persist.

**SPINA BIFIDA.**—There is a congenital defect in the union of the laminae of one or more vertebrae, associated with malformation of the spinal cord or its membranes. It occurs most frequently in the lumbar regions, and persons may live to a good old age thus affected.

**SYRINGOMYELIA.**—Syringomyelia is a chronic affection of the spinal cord characterized anatomically by the pathologic formation of cavities in its gray matter, and clinically by peculiar disturbances of sensibility associated with trophic disorders. Causes: (a) Embryologic malformations. (b) A gliosis. (c) Traumatism. (d) Development of embryonal neuroglial tissue in which hemorrhage or degeneration takes place with the formation of cavities. (1) The characteristic lesion is a cavity which forms in the cord in or near the central canal and extends into the gray matter of the anterior, or more frequently the posterior horns. It is most often situated in the cervical and thoracic portions of the cord. (2) On transverse section the cavity may be oval, circular, or narrow and fissure-like, or it may present the appearance of two or more cavities independent of each other or intercommunicating. (3) The contents of the cavity are usually a colorless liquid. Occasionally it may be a yellow or brown gelatinous substance, or it may consist of blood and the products of its degeneration. The white matter of the cord in moderate cases is unaffected, but where the cavity is large and pressure from the sclerotic tissue has become great, the white matter is in its turn involved, being crowded to the periphery and more or less unable to carry on its functions.

**TUMORS.**—Tumors and cysts of the brain are of common occurrence and of the greatest variety, such as fibro-endothelioma, sarcoma, psammosarcoma, fibroma, osteophytes, perithelioma, lipoma, myxoma, glioma, gliosarcoma, angioma, neurofibroma, and neuroma. Of the granulomata, syphilis, tuberculosis, and actinomycosis are the most common, and of the parasites the cysticercus and echinococcus.



## CHAPTER XVI

### EXAMINATION OF THE NASOPHARYNX, EYES, AND EARS

#### EXAMINATION OF THE NASOPHARYNX.

IN order to expose to view the upper air-passages, nasal, pharyngeal, laryngeal, and accessory cavities, epiglottis, etc., Harke's<sup>1</sup> method has come into general use. If the procedure is properly carried out, the parts when returned to their normal position present no noticeable deformity, though during the examination such a result seemed almost impossible.

**HARKE'S METHOD.**—The brain having been removed and the examination of the skull completed, the anterior skin flap is dissected away from the frontal bone down to the root of the nose, while the posterior flap is dissected away some distance below the foramen magnum. It is not necessary that the primary incision of the scalp behind the ears be made lower than the mastoid process on each side. Next, directly in the median line, the skull is cleft with a small saw into two lateral portions. For the sake of convenience the saw markings may be divided into two sets (Fig. 150), the first starting from the front in the frontal bone, extending down to the nasal bone, and continuing to the foramen magnum (*AB*), and the other starting at the occipital bone and extending to the foramen magnum (*CD*). The atlas and axis are sawed through if much room be desired. The sawed portions are now separated by means of a chisel and hammer, any portions of mucous membrane that may appear being severed with a knife or scissors. By means of strong lateral traction the two segments may be pulled apart, and the entire region down to the vocal cords will thus be exposed. Usually the incision passes to one or the other side of the nasal septum. The walls of the accessory cavities are readily cut away with strong scissors, and a plain view is obtained of the maxillary sinuses as well as the frontal, sphenoid, and ethmoid. Even the epiglottis and vocal cords can be examined by this method (Fig. 151). In order to view the parts better, light may be thrown in by means of a mirror.

Another method is to drill holes just in front of the sphenoid and a little behind and to the right and left of the crista galli, and then with a

---

<sup>1</sup> *Berliner klin. Wochenschrift*, 1892, no. 30, p. 742; *Virchow's Archiv*, 1891, vol. cxxv, p. 410. *Beiträge zur Pathologie und Therapie der oberen Athmungswege*, Wiesbaden, 1895.



FIG. 150.—Method of examining nasopharynx, eyes, and ears. The sawing for opening the nasopharynx is done in the median line from the frontal bone, *A*, to the anterior portion of the foramen magnum, *B*, and from the occipital bone, *D*, to the posterior portion of the foramen magnum, *C*. The sawing can best be accomplished by standing on the table directly over the head, the finger-saw being especially useful at the beginning and the end of the operation. *E* and *F*, lines of incisions for the removal of the eyes; *G*, situation of the ear-ossicles; *K*, *I* and *L M N O*, lines for removal of the ear-ossicles; *P* and *Q*, drill-holes for saw-markings in the oval method of examining the nasopharynx.





saw or a chisel make an ovoid incision extending almost to the foramen magnum, and remove the portion of bone which hides the nasopharyngeal cavities. (Fig. 150, *P. Q.*) The two lateral halves are then brought together and wired as in Fig. 152. Nasal obstruction may cause deformities of the upper jaw, teeth, and palate.<sup>1</sup>

## EXAMINATION OF THE EYES.

For this purpose a triangular piece of the orbital plate of the frontal bone is broken through with a hammer or chisel, care being taken not to injure the optic nerve in the optic foramen, the remaining portion of the eye and the nerve being well protected. (Fig. 150, *E* and *F.*) The direction of the nerve can be determined by observing the situation of its exposed portion, and the chiselling done a small distance on either side of its normal position. The pieces of bone are removed with the nippers and the optic nerve is carefully dissected out, its cut end being held with the fingers or forceps. The capsule of Tenon and the fat are removed, and the entire eye is excised or, if this is not permitted, an incision is made in the sclerotica posterior to the conjunctival attachment. This requires a very sharp knife, as the tissue is extremely tough. A circular incision is made around the entire eye, and the fundus is exposed. A piece of dark cloth or cotton dipped in ink is placed in the remaining portion of the eye in order to hide any disfigurement, and the cavity is packed with cotton.

If only a macroscopic examination of the retina and other structures is desired, the retina may be floated out in normal salt solution and then separated from the choroid. If the retina is to be fixed for microscopic examination, the incision should be as nearly equatorial as possible and the fundus placed immediately in Orth's or Müller's fluid or ten per cent. formalin, or fixed by exposing for three minutes to the fumes from a one per cent. osmic acid solution heated just to the boiling point. The eye is then put for twelve hours into Lindsay Johnson's mixture:

Potassium bichromate, two and one-half per cent.....	70 parts.
Osmic acid, two per cent.....	10 parts.
Platinic chlorid, one per cent.....	15 parts.
Acetic or formic acid (to be added just before using)....	5 parts.

The gloss of the cornea disappears as soon as death comes on. After twenty-four or thirty hours, and often earlier, the bulbus softens

<sup>1</sup> COLLIER, *Lancet*, October 18, 1902, p. 1038.

and the cornea and retina become dull. The conjunctiva is now removed more easily from the cornea, and the sclera which is not covered by lids becomes brownish black and dry. (Orth.) According to Runge, several days after death a diffuse redness occurs in the transparent media of the eyes of a foetus that has died *in utero*. The redness affects first the cornea and then the lens,—extending from without inward,—in this way indicating approximately the date of death.

The position of the eye may be altered. Exophthalmus, or protrusion of the eye, may be caused by a retrobulbar tumor, œdema, hypertrophy of fat, collection of blood from hemorrhages, emphysema, inflammatory exudates, Basedow's disease, etc. Enophthalmus, or retraction of the eye, may take place in atrophy of the fat, loss of liquid, as in cholera, deformity from scars, etc. The consistency of the eye varies, being increased in glaucoma and diminished in certain forms of degeneration. Frontal empyema may occur.

#### EXAMINATION OF THE EARS.

A fair idea of the condition of the middle ear may be obtained simply by chipping away the roof with a chisel or biting it off with bone-forceps, but if a closer inspection is desired the petrous portion of the temporal bone and the mastoid process had better be removed together.

Extend the incision from its original point back of the ear and along the anterior border of the trapezius about half-way down the neck. Reflect the flaps with their soft tissues so as to leave the bone clear. Begin at the apex of the petrous portion of the temporal bone and with a chisel laid flat break through the petrobasilar suture to the jugular foramen, and chisel or saw through the skull on a line from the jugular process of the occipital to a point about five centimetres posterior to the base of the mastoid process. Anteriorly chisel or saw through the skull on a line from the apex passing posterior to the spinous process of the sphenoid; or between the foramen ovale and the foramen spinosum and well anterior to the external meatus, just cutting off the root of the zygoma.

Woodhead<sup>1</sup> uses the following method:

“The temporal bone, with its petrous portion containing the internal ear, may be taken out and examined after removal of the brain, by stripping off the dura mater from the base, dissecting off the skin

---

<sup>1</sup> *Practical Pathology*, 1892, p. 28.

and muscle, detaching the external ear from the bone, and disarticulating the jaw; then, taking the margins of the temporal bone as the base of a pyramid, the apex of which is a little beyond the inner extremity of the petrous portion, two saw-cuts are carried almost vertically downwards so as to bound the pyramid, and then with a bone-chisel and mallet the whole temporal bone may be removed, after which it may be softened in a decalcifying fluid; or the internal ear may be dissected out with a small saw, a pair of sharp well-fitting bone-forceps, and a sharp gouge and chisel. The internal ear or tympanic cavity and mastoid cells may also be opened up with the aid of the above instruments."

By sawing or chiselling as in Fig. 150, *K J I* or *L M N O*, the ear-ossicles and internal ear may readily be reached. The method of




FIG. 153.—Chain-saw used in removing the auditory apparatus, the lower jaw having first been removed. (After Letulle.)

using the chain-saw is seen by referring to Fig. 153. In this illustration is also shown what is known as the T incision, which consists in



membrane and the articular ligaments become dark red in color, swollen and infiltrated, and covered with pus; later the cartilage and lastly the bone (molecular necrosis) are attacked, often causing disarticulation of the ends of the bones. Perforation may be primary or secondary. A purulent effusion, remaining a long time without serious destruction, is called catarrhal synovitis.

*Chronic Inflammations.*—Chronic articular dropsy (hydrarthrosis) is a serous or serofibrinous inflammation, usually seen in the knee, wrist, or elbow. The synovial membrane is thickened, indurated, and may have patches of fatty degeneration. In the knee the patella may be lifted and the bursæ distended by a thick or thin or gelatinous secretion, and the synovial membrane may protrude through the fibrous bands of the capsule. The synovial tufts become large and projecting; the joint-cartilages degenerate and proliferate. Adhesions or destruction of the joint may follow.

*Chronic purulent arthritis* is usually associated with tuberculosis or is due to extension from adjoining parts. The capsular ligaments and synovial membranes are infiltrated and covered with fibropurulent deposits; the cartilages are cloudy, fibrillated, or necrotic, the marrow suppurating, and the joint filled with pus or numerous abscesses forming around the joint. Ankylosis or dense fibrous adhesions are found in cured cases.

*Chronic, dry, ulcerative arthritis* occurs in old age, accompanying neuropathic disorders, or as a sequel to rheumatism. It consists in a proliferation of the synovial membrane, forming fringes, and in sclerotic thickening of capsule and ligaments. Fibrillation and cleavage of the cartilages, with patches of calcareous, amyloid, or fibroid degeneration, may occur. The denuded bone of the articulating surfaces may ulcerate or become sclerotic and waste, the capsule becoming so large that dislocations may occur. In the senile form the hip is usually affected, but the shoulder, elbow, phalangeal joints, and the patella of the knee may be involved. In tabes dislocations are very common in knee, shoulder, and elbow.

*Arthritis deformans*, chronic gout, or rheumatoid arthritis, is a chronic disease of the joints, characterized by degenerative changes in the cartilages and synovial membranes, by periarticular formation of bone, and great deformity (Osler). It is often associated with infectious diseases, as gonorrhœa, gout, and rheumatism. In all forms the articular surfaces are hyperplastic and softened. Later, absorption

takes place, the ends of the bones becoming eburnated and polished; the head of the femur has entirely disappeared; generally it becomes conical, flat, or broad. The bone marrow liquefies, forming cysts; subchondral cysts and deep-seated bone cavities occur. At the edges, where the friction is less, irregular nodules (osteophytes) develop and calcify. Capsules, synovial membranes, fringes, and ligaments thicken and become infiltrated with lime salts. There is always a complete absence of uric acid. Great deformity, not infrequently ankylosis and dislocation, occurs. There is often marked atrophy of both bones and muscles. Arborescent lipomata are found. The smaller joints of the hands and feet are usually first affected, the fingers being deflected to the ulnar side. In severe cases all the joints may be more or less involved. In old people the disease is apt to attack the hip, knee, shoulder, or spine. *Spondylitis deformans* is due to the formation of osseous bridges between the vertebræ. Heberden's nodosities, a form of the disease, consist of small nodes or tubercles about the dorsum of the phalanges; this form very rarely affects large joints, as the knee.

*Gouty arthritis* is the deposition of urates in the articular structures, usually in the metatarsophalangeal joint of the great toe (podagra) or a finger-joint (cheiragra). The periosteum, tendons, ligaments, and skin are more or less inflamed. The joint contains a clear fluid, with crystals of sodium urate, sodium chlorid, calcium carbonate, and calcium phosphate, hippuric acid, and other uric-acid compounds. Chalky, mortar-like, nodular masses, tophi, are found in the matrix of cartilage and ligaments. In old cases these are also found in the bone, periosteum, tendons, and bursæ. Fibrillation and erosion of the cartilages cause abscess-like cavities, which may open externally.

*Gonorrhæal arthritis* occurs in one knee-joint, between the third and the sixth week of the disease. It has also been seen in an ankle and hip. There is a fibrous metaplasia, also a fibrous or osseous ankylosis. Ulceration of the cartilage, bone, and capsular tissue may occur, as a rule, with a purulent effusion.

*Rheumatic arthritis* occurs in several joints at one time, generally in the hip, shoulder, and jaw. It begins as a hyperæmia of the synovial membranes, with an increase of fluid, followed by thickening and elongation of the ligaments and later by absorption or ossification of the interarticular cartilages, which become rough, fibrillated, and often converted into a tough, felted mass. Finally, there are induration and eburnation of the bony extremities. It may involve joints in succes-

sion, and, in rare cases, all the joints. It always causes ankylosis, by fibrous adhesions of the ligaments and bony deposits in and around the joint.

*Chronic ankylosing arthritis* is the most common anatomic feature of chronic rheumatism, and is due to a vascularization and fibrous metamorphosis of the articular cartilages, with coherence of the opposed cartilages.

*Spinal or neurogeneous arthritis*, usually associated with tabes dorsalis, syringomyelia, degeneration of the anterior horns of the gray matter, arising from section of spinal nerves, consists in a rapid destruction of the articular ends of bones, thickening and ulcerative destruction of synovial membranes and ligaments, and a serous effusion into the joint, with swelling of the surrounding tissue and spontaneous dislocation.

*Syphilitic and tuberculous arthritis* are described elsewhere. I have had one case of *pneumococcal arthritis* of the knee, with streaky hyperæmia of the skin, boggy swelling, purulent infiltration, and hemorrhagic effusion. *Toxic arthritis*, due to alcohol, occurs, usually in small joints, and always associated with enlarged liver, spleen, and lymphatic glands.

Degenerations generally occur in the cartilage of a joint. After hemorrhage into a joint, hæmatoidin is often found as crystalline and amorphous masses in the superficial cells. Ochronosis produces diffuse brown patches in cartilages, due to saturation of the matrix with some unknown coloring matter. Mucoid degeneration of the matrix produces a turbid fibrillar appearance, which may go on to complete disintegration, and is often associated with fatty change in the cells. Fatty degeneration, a translucent gray material, appears, in senile softening, associated with calcification; and in chronic inflammatory disease. It attacks mainly the costal cartilages, but may occur in margins of the articular cartilages, and in places where the matrix is already in a process of fibrillation and degeneration. Hyaline and amyloid degenerations occur in the capsules and cartilage-cells. Amyloid degeneration may also affect the matrix.

**ATROPHY.**—Acquired or true atrophy is: (1) eccentric, the bone being normal in size, but on section showing great increase in the cavities and in the amount of the cancellous tissue, and decrease in the compact tissue; (2) concentric, the bone being slender and the external compact tissue showing local defects (osteoporosis), or being exces-



sively thin and brittle (osteopsathyrosis, fragilitas ossium). The medullary canal is always contracted.

Atrophy may follow trauma, such as fracture, luxation, or epiphyseal injury, and disuse, as seen in old stumps and unset fractures. Pressure often thins the bones markedly. In hydrocephalus the internal surface of the skull may be rough, or the inner table may be entirely absorbed. The Pacchionian bodies often make deep pits in the temporal bones. The vertebra, sternum, and other bones may be found deeply eroded and perforated by aneurisms and tumors; even scars may cause atrophy of the bone upon which they make pressure. Tumors of the marrow, periostitis, and osteomyelitis also cause atrophy of the bone by their pressure. A peculiar type of atrophy is seen in the aged, affecting those bones with only a slight muscular covering, as parietal, maxillary, and pelvic bones. The external table of the skull may be entirely resorbed. Perforation of the entire thickness has occurred, bony deposits being often found at the same time on the inner table. The bone is rough, dull, and lustreless, with shallow erosions, not uniform in shape or position. Nervous diseases (neuropathic and neuromyopathic), infantile paralysis, inflammation, rickets, and many other pathologic conditions are often associated with osseous atrophy.

**CHONDRITIS.**—This affection is often a sequela of severe arthritis; it usually occurs in articular cartilages, which become turbid and disintegrate. Erosions, caries, and more or less extensive necrosis are not uncommon. Hypertrophic proliferation, general or local, may occur with any productive inflammation of the cartilage or fibrous tissue of a joint. It is common in arthritis deformans and tuberculous arthritis. In cartilage it is nodose or tuberculous, while in the capsule or synovial membrane it appears as a diffuse thickening or as a papillary excrescence. Loose bodies, usually found in the knee, elbow, and wrist, rarely in the hip, shoulder, elbow, or ankle, may be single or numerous, a knee with 1047 of these bodies having been reported. These may be entirely free or attached by a slender stalk. They may be composed of fibrin, the remains of hemorrhage into the joint, or may be a proliferation of the synovial or fatty tufts, pieces of bone or cartilage detached by violence, foreign bodies which have penetrated the bone, loosened nodular masses, cartilaginous, osteomatous, fibrous, lipomatous, or lipoma-arborescent, or displaced semilunar cartilages. These bodies are usually oval, lenticular, or devious in outline, often faceted.

They are associated with arthritis deformans and rheumatoid arthritis, and in many cases appear without signs of previous inflammation.

Dislocations of the semilunar cartilages are due generally to separation of the anterior attachment of the cartilage from the tibia, which may be torn transversely through the edge of the meniscus or split longitudinally. A central tear has also been described. Cartilages, infrapatellar pad, and ligamentum alare are usually thickened. The displacement is inward, towards the centre of the joint, so that the leg cannot be extended.

**FRACTURES AND DISLOCATIONS.**—Fractures, the most common injuries to bones, are either complete or incomplete. A certain amount of repair callus will be found in all cases. Contiguous bones, as tibia and fibula, may coalesce during repair and lead to a synostosis. In cases with great displacement, with soft parts between the fragments, with existing debility or other unfavorable conditions, there may be no union, or simply a firm, fibrous adhesion (syndesmosis), or a false joint (pseudoarthrosis). Compound fractures, affording a favorable opportunity for the introduction of pyogenic organisms, are often associated with caries, necrosis, or osteomyelitis.

Diastasis, a pathologic separation of the epiphysis from the diaphysis, occurs usually as a sequela of accident or ulceration, the epiphysis being pushed off by the resulting granulations. It is most common in the upper part of the femur, in the lower portion of the humerus, and in the tibia. In rare cases the dislocation may take place between the manubrium and the gladiolus, or the head of the femur may be found loose in the acetabulum.

Congenital dislocation of hip, single or double, is associated with softening of ligaments, effusion, fungous synovitis, hydrarthrosis, caries, arthritis, and arrest of development. The acetabulum is narrowed, elongated, less concave than normal, and occasionally filled with fat, connective tissue, or exostoses. The head of the femur is flattened, but larger than the acetabulum; it lies on the dorsum of the ilium or obturator foramen; the neck may be wanting; or it may be atrophied. If the patient has walked, there is usually a depression of the ilium and some lordosis. The gluteal muscles are contracted, the unused muscles atrophied, and the pelvis is contracted above and expanded below. Congenital luxation of the sternum has been reported. If acquired dislocations with rupture of the capsule, tendons, ligaments, muscles, and other structures around the joint be not reduced, the

muscles and ligaments may atrophy, the synovial fluid be destroyed, and the bone be partially absorbed, or the bones may unite by firm, fibrous, cartilaginous, or bony adhesions (ankylosis). This may also follow non-use of a part, articular disease, trauma, phlegmonous erysipelas, burns, or as a sequela of tuberculous, gonorrhœal, gouty, rheumatic, syphilitic, neurotic, or puerperal affections. A form of ankylosis of the spine is sometimes seen in typhoid fever. Congenital ankylosis of the entire skeleton has occurred. False ankylosis is due to bands of cicatricial tissue, adhesions of the ligaments and capsule, or organization of inflammatory deposits.

**INJURIES.**—Cuts, gunshot wounds, and stabs may result in acute inflammation, effusion of blood, empyema, ulceration of articular capsule, necrosis of cartilage and bone, or a more or less complete disorganization of the entire joint. Septicæmia and pyæmia may supervene. Crushing injuries to joints almost always end in abscess, usually associated with bony or cartilaginous fragments, which remain as foreign bodies in the joint.

**MARROW.**—The marrow in children is soft, bright red in color, rich in cells and blood-vessels. In middle life there is an increase in fatty tissue, giving a yellow or yellow-red color of oily lustre, while in old age it atrophies, becoming gelatinous, with clear mucinous fluid, and there is a diminution of fat and a decrease in the number of cells present. This atrophy may follow chronic emphysema, phthisis, chronic disease of the kidney, or starvation. Lymphoid marrow is gray red or dark red, according to the amount of blood it contains. In pernicious anæmia the marrow of the long bones resembles raspberry jelly, while in leukæmia it has a flesh-pink to a gray-yellow color, like that of pus. Fatty degeneration occurs in the cells and capillaries of the marrow, sometimes with necrotic foci, in cases of typhoid, typhus, and relapsing fever.

In osteomyelitis there is often a purulent inflammation, frequently complicated by transformation of the marrow into a vivid red, tough, fibrinous material, and with effusion into the cavity of a joint. Necrotic changes in the bone follow. Hypertrophy of the marrow-cells is seen in oligæmia, leukæmia, chronic pulmonary tuberculosis, chronic suppurative osteitis, cancerous cachexia, typhoid fever, croupous pneumonia, septic affections, acute endocarditis, and smallpox; while hypertrophy of the fatty tissue occurs in cases of general atrophy of the skeleton, sometimes involving the entire bone.



—

rickets or in men who have lived laborious lives. The external articular surface of the tibia or femur is retarded in its growth or depressed, so that these bones form with each other an obtuse angle. This may be associated with separation of the epiphysis, caries of the external condyle of the femur, or arthritis deformans.

Club-hand, hallux valgus, club-foot, talipes varus, talipes valgus, talipes equinus, talipes calcaneus, talipes cavus, and talipes planus are, as a rule, due to perverted development of tendons.

Contractures are associated with poliomyelitis, caries of the spine, trauma of the cord or peripheral nerves. They are the result of fixation of a joint in a deformed position, the character depending upon the group of muscles paralyzed. Dupuytren contracture is a scar-like contraction with fingers flexed, due to trauma, rheumatism, or gout, affecting the palmar fascia. Small, hard, nodular fibromas are found along the course of the contraction. Late in the disease the skin is affected.

Spinal curvatures are associated with pleural effusion, large tumors, unilateral contraction of the thorax, cirrhosis of the lungs, oblique fixation of the pelvis, rickets, tuberculosis, weakness of the muscles, occupation, etc. According to Bradford and Lovett, scoliosis, or lateral curvature, is most common in the thoracic region, usually to the right, with compensatory curve, in the lumbar region, to the left. Kyphosis may be a rotated lateral deviation or a result of disease, as tuberculosis. It forms a posterior protrusion of the vertebral spines; if due to rotation, it is usually to the convex side. The vertebræ become wedge-shaped. Ossification of the ligaments is at times found. Lordosis is always associated as a compensatory curve. Synostosis, or ossification in a situation not normally ossified until advanced life, causes a marked deformity, especially of the pelvis, shortening in the base of the skull, craniostenosis, microcephalia, etc., and depressions of the bridge of the nose.

OSTEITIS is almost invariably associated with periostitis. It may follow trauma, such as fracture, amputation, and gunshot wounds; or infections, as pyæmia, scarlatina, measles, typhoid and relapsing fever, dysentery, smallpox, mumps, gonorrhœa, and acute articular rheumatism. Four forms of osteitis exist: (1) Rarefying (osteoporosis), a chronic form associated with wounds, syphilis, or tuberculosis, which consists in resorption of the spongy bone, with the formation of cavernous excavations. (2) Osteosclerosis, a reparative reaction, occurs in

the same diseases and consists in the eburnation of the entire bone.

(3) *Osteoarthropathies* (hyperplastic osteitis), associated with chronic tuberculous lung diseases, involves the terminal phalanges, which are swollen like drum-sticks, with their articular ends irregularly thickened; it is also seen in rhachitis, osteomalacia, and osteitis deformans. (4) *Osteitis caseosa* is always tuberculous. Purulent osteitis may follow typhoid, scarlatina, measles, and pyæmia, and is usually due to a secondary staphylococcic or streptococcic infection. It arises spontaneously in the femur and tibia; it is in almost every case associated with osteomyelitis, and at times with gangrene.

In *osteitis deformans* we have an inflammatory disease of old age, consisting in a wide-spread absorption of the bone and the deposition of new bone. It may be limited to the femur, cranial bones, or spine, or involve the greater part of the skeleton. The resorption is marked in the cancellous and cortical regions, where the osseous trabeculæ may be replaced by gelatinous or fibrous tissue. This softening allows the long bones to bend at abrupt angles and gives rise to many deformities, of which kyphosis is the most frequent. Cysts are often found. The deposition of new bone, which is especially seen in the skull, starts from the periosteum and from the marrow, causing thickenings of the bone.

**OSTEOMALACIA.**—Osteomalacia (*mollities ossium*) is a rare disease, occurring especially in pregnancy, and is characterized by a rapid and general resorption of the inorganic salts, advancing from the centre outward and including all except a thin layer next to the periosteum. The marrow is increased and splenified, or replaced by a dark semifluid material. On section, the spaces contain a reddish gelatinous mass, which later becomes yellow and fatty. Cysts containing a clear, turbid, or hemorrhagic fluid have been seen in the interior of the bones. The bones are very light, bend and break readily, or may even be cut with a knife. Fractures are commonly multiple, occur spontaneously or from very slight injury, and tend to repair, even in the active stage of the disease, the callus, however, remaining free from bone salts. At first the disease is limited to the pelvis; later the entire skeleton may be involved. In the so-called non-puerperal form the disease starts in the spongy bones of the vertebræ and thorax, extending to the extremities and finally even to the cranium. The sacrum is pushed downward by the weight of the body and the acetabula upward and inward by the femora, producing a characteristic pelvic



deformity. The disease is associated with general cachexia and often with pneumonia.

**PERIOSTITIS.**—The normal periosteum presents a yellowish-gray color, while in suppuration it is distinctly yellow. It may be raised or inflamed by traumatism, perforation of compound fracture, abscesses, tumors, infectious granulomata, or extension of inflammation from neighboring structures. Blood under the periosteum, particularly near the epiphysis, is seen in children with Barlow's disease. Simple acute inflammation, usually local, produces a reddening, thickening, and a greater adherence of the periosteum to the bone. Suppurative periostitis, generally associated with osteomyelitis, affects growing bones, and is rare after the union of the epiphyses. The exudate, or the hemorrhage beneath the periosteum, rapidly separates the membrane from the bone, causing stretching, occlusion, or thrombosis of the blood-vessels passing into the bone; hence necrosis of the superficial osseous layers results. Pyæmia and infective fat embolism may occur before the abscess is opened. A diffuse form, attacking the long bones in those presenting a strumous diathesis, often ends in a rapid suppuration.

Fibrinous, ossifying, or productive inflammation follows chronic inflammation of the joints, syphilis, rickets, and tuberculosis, and results in osteoses, or bony thickenings. The periosteum is hardened, and a projecting node is formed beneath it, which may become fibrous or calcified (periostitis ossificans). This calcification begins as a vertical process at the surface of the bone, at first distinguishable from the old tissue, but later blended with it. In syphilis the subperiosteal nodes show a marked tendency to suppurate, and in rare cases suppuration occurs, but due to other causes, producing a malignant purulent periostitis. Typhoid bacilli have been recovered from these cases. Albuminous periostitis, a mild inflammation with a ropy, albuminous exudate, is found only in the bones of the young. Tuberculous periostitis also occurs, most often in young patients, and has more or less sharply defined granulomatous foci, containing tubercles which become caseous and soften, and give rise to peripheral caries, sacculated cold abscesses, consecutive abscesses, sinuses, or fistulous tracts. The caseous nodes are surrounded by a zone of induration and granulations, which may be so luxuriant that they form mushroom-like excrescences over the external orifice of the sinus. Simultaneous with increase of the caries there are proliferation of the periosteum and the formation of con-

siderable new bone. This is usually absent in the cranial bones, where resorption alone more often occurs.

**OSSEOUS TISSUE.**—**Osteitis**, or **osteitis**, is a constitutional disease of childhood, characterized by alterations in the conversion of cartilage into imperfect osseous structure. Congenital osteitis is rare, the so-called fetal osteitis being merely a disturbance of growth closely resembling myxodema. Ossific changes often occur in the periosteal and medullary spaces of the long bones, especially between the shaft and the epiphyses, where a soft and irregular mass of proliferation, 5 or 10 millimetres thick is found. The bony tissue is softer and more vascular than normal; the marrow, wider and darker in color. The periosteum strips off easily, revealing a spongy tissue which looks like decalcified bone. Large osseous firmations occur under the periosteum at the insertions of the tendons and apophyses. The shafts of the long bones are usually bent and shortened and the short bones flattened. The cranial bones are thin and atrophied at the sites of pressure *craniotabes*; the frontal and parietal bones often have flattened swellings, and the fontanelles remain so large that the head suggests hydrocephalus. A swelling may occur around the hip, simulating coxalgia. Associated with the bone lesions are anemia, enlarged spleen, changes in the liver, muscular atrophy, and mucous inflammation of the mucous membranes, especially of the intestines and respiratory tract. The lungs and heart often present changes due to the deformity of the chest. Dentition is delayed. After the active stage of the disease the bones become very hard, heavy, and deformed.

**TENDONS, SHEATHS, AND BURSE.**—Acute tenosynovitis, simple or hematogenous, results from wounds, bruises, strains, or excessive exercise. Gonococci, pneumococci, or pus cocci infect the tendons usually of the dorsum of the hand, producing a purulent exudate, which shows a marked tendency to burrow between the sheath and the tendon, sometimes for considerable distances. In the dry form deposits of fibrin are found upon the inner surfaces of the sheath, giving rise to a rubbing or creaking sensation. The tendon is cloudy and swollen, the intervascular substance often suppurating or necrotic. Chronic tenosynovitis is generally gouty or rheumatic or the result of healing wounds. Calcareous deposits or gouty urates are particularly common in this form, and may cause necrosis, inflammation, or the formation of new fibrous tissue. Tuberculous tenosynovitis occurs in the walls of the sheath, with exudation. In advanced stages there may be

fungous granulations on the tendon. Tubercle bacillus may be found in the arborescent lipoma, a papillomatous, fatty outgrowth of the synovial lining, it being a debated question as to whether or not this organism causes the lesion. Hygroma or ganglion, due to chronic irritation, is a cystic mass which contains rice-like bodies in a serous fluid beneath the sheath of the tendon.

Acute bursitis, acute hygroma, is a fluctuating tumor with serous, serofibrinous, or purulent exudate, the result of injury or hæmatogenous infection. The walls are generally thin, but may be greatly thickened. In the chronic form, hygroma, hydrops bursarum, or housemaid's knee, the contents, in the early stages, are mucilaginous and viscid; later, thin and limpid. Loose bodies are frequently found in these cysts. Tubercles may develop in the walls of the sac, associated with serous effusion, or the walls may become thickened and permeated by fungous granulomatous masses which may undergo caseous degeneration. A ganglion is a round, oval, or lobulate cyst, varying in size from that of a pea to a pigeon's egg, and containing a reddish-yellow crystalline jelly or colloid material, probably the result of a recurring slight injury. It appears on the dorsal aspect of the intercarpal joints.

TUMORS, CYSTS, PARASITES, ETC.—True osseous tumors occur, which may be primary or secondary, myelogenic or periosteal. Osteoma, exostosis, osteophytes, and enostosis are found, the latter arising from the periosteum or cartilage, during the period of growth at the diaphyses of the long bones. Two kinds of osteoma are seen,—the cancellous, or spongy, and the compact osteoma. On the skull they are usually small, round, conical, or mushroom-shaped. *Chondromata* usually arise from congenital, misplaced islands of cartilaginous tissue, though often not until late in life, and are found especially in the hands and feet of children and young adults, sometimes producing marked and grotesque deformities. They are nodose or tuberous excrescences which, especially on the scapulæ, long bones, or ribs, reach considerable size. They may soften and form bone cysts. The myelogenous are at first covered with a shell of bone. Cartilaginous exostoses start from the epiphysis, as long or rounded bony projections, the apex or a greater part of their surface being covered with cartilage. These new growths undergo fatty, calcareous, and mucoid degeneration. *Lipomata* and *angiomata* are rare. Fibromata, nodular and highly vascular growths, occur on the facial and cranial bones and



in the buccal and nasal cavities. A more rare tumor is the encapsulated *myxoma*, which arises simultaneously in the periosteum and marrow, the myelogenous form having no capsule and destroying the bone rapidly. Both varieties give rise to cysts, single or multiple. The *sarcoma* is the most common primary tumor, and the cells comprising it may be round, spindle-shaped, or giant-celled. It is often telangiectatic. The myelogenous sarcoma, usually occurring in the epiphysis of the tibia, humerus, etc., is, even until it reaches considerable size, covered with a bony shell, which may fracture spontaneously. On section, a milky fluid may exude. Periosteal sarcomata occur anywhere and are generally mixed tumors. A special variety, chloroma, green and yellow in color, is seen in the facial and cranial bones. *Carcinoma* is always secondary and is usually due to direct extension; it is seen in the skull, sternum, and ribs, where it forms either a circumscribed node or diffuse infiltration and is always accompanied by lacunar resorption.

*Cysts* arise from lacunar atrophy, osteomalacia, disintegration, or excessive resorption. They are common on the alveolar processes of the upper or lower jaws, associated with enlarged and tortuous veins, and on the clavicle, usually connected with some solid tumor. A turbid or hemorrhagic fluid exudes on section.<sup>1</sup>

The *Echinococcus* is the most common parasite. It occurs in the long, pelvic, cranial, and vertebral bones as a single sac or as internal or external daughter cysts. There are always associated some resorption and atrophy of the affected bone, and there may be distention, inflation, or spontaneous fracture. *Cysticercus* is very rare.

*Aneurisms* are generally anastomotic, but primary aneurism may occur, usually in the cancellated tissue of the head of the tibia, associated with absorption of the compact bone and periosteum. *Hæmatomata* are occasionally produced by hemorrhage following trauma or rupture of a softening tumor.

---

<sup>1</sup> For a description of the benign dentigerous cysts of bones see BLOODGOOD, Jr., *Amer. Med. Assoc.*, Oct. 15, 1904.

## CHAPTER XVIII

### POST-MORTEM EXAMINATIONS OF THE NEW-BORN <sup>1</sup>

IN performing a postmortem on a child it is sometimes advantageous to remove the viscera *en masse*, scissors being largely used for this purpose in place of the knife, even to the cutting of the clavicles in their central part. In France evisceration is done quite frequently in the adult, and has the advantage of saving time by permitting the removal of the spinal cord while the thoracic and abdominal organs are being examined both from behind and in front. To practise evisceration the trachea and œsophagus are twice tied as high up as practicable, divided between the ligatures, and the lower portion then elevated with the free hand. All the posterior attachments are cut as close as possible to the vertebral column until the diaphragm is reached. The cervical and thoracic organs are then brought out of the body and laid over the opposite costal margin to the side upon which the operator next works. The diaphragm is now excised laterally and posteriorly, adhesions being severed with the knife as before. It is well to pull from below upon the liver, stomach, and spleen, so that these organs will not be injured by the manipulations. The crura being cut loose, the diaphragm is free. The posterior peritoneum having been already incised by the removal of the diaphragm, the kidneys are readily found from above, and when removed the psoas muscles come prominently into view. The common iliac vessels, round ligaments, etc., are next incised. Two ligatures are now applied to the rectum, which is then divided between them. When everything which holds the abdominal organs in place has been loosened with the hand, the organs of both the thorax and the abdomen can be readily removed, leaving only the bladder and organs of generation *in situ*; these may be excised later, in the same manner as that described for the adult on page 15; or, by means of the incisions there given, removed while still attached to the other abdominal viscera. The cervico-thoracic abdominal cavity is then to be carefully inspected.

The body of a child thus disembowelled can be kept for a long time, especially if the thoracic and abdominal cavities are packed with

<sup>1</sup> For additional information see pages 243, 344, and 358.

a mixture of equal parts of bran and salt to which a little white arsenic has been added. The cadaver may then advantageously be surrounded with cotton and a circular bandage applied to the chest and thorax. Parental consent to the performance of an autopsy may sometimes be obtained by giving assurance of the preservation of the body by this mode of procedure.

In the child there will at once be noticed the large size of the liver, appendix, and adrenals; also the perpendicular situation of the stomach, it being more difficult to distinguish the greater and lesser curvatures than in the adult. The dissection of the cardiac plexus should always be made after diphtheria and other contagious diseases attended with cardiac failure. This examination also includes the pneumogastrics and the cervical sympathetics. Wrisberg's ganglion is found by looking carefully in the region of the arch of the aorta, the right branch of the pulmonary artery, and the fibrous remnants of the ductus Botalli. Walckhoff believes that the expansion of the lungs changes the position of the heart to such an extent that the arterial canal is at once twisted, thus stopping the circulation through it. The unexpanded lungs are of firm consistence, do not crepitate, and do not cover the anterior surface of the heart. The color is a brownish slate. The expanded lungs are of light rose tint, somewhat blood-stained, except where the bluish spots of fetal atelectasis persist. The methods of examining the umbilical vessels and the ductus arteriosus are readily seen by referring to Figs. 154 and 155 respectively. Study with care the point of insertion of the gelatin of Wharton to the circular fold of the skin at the umbilicus in a new-born babe.

The removal of the child's brain is more difficult than that of an adult, because, first, it is much softer, and, second, the dura is normally adherent to the cranium. But it is easier in one respect,—the bones and sutures are not ossified. In a new-born child the brain is so soft that its removal without injury is almost impossible. In such cases it is advisable to lay the body for a short time on ice sprinkled with salt, in order that the brain may become firmer by the consequent lowering of the temperature. By another method—and it is the one from which I have obtained the best results—the child is placed in a large basin or tub containing a strong solution of common salt (about half a bucketful to four or five times this amount of water) and held steady by an assistant while the brain is removed by operating beneath the surface of the liquid. As brine of the above strength has a specific





FIG. 154.—Examination of the umbilical vessels. (After Nauwerck.)




FIG. 155.—Examination of the ductus arteriosus. The sound is represented as introduced into the ductus arteriosus Botalli; this duct usually closes about the fourth day after birth. (After Nauwerck.)



FIG. 156.—Removal of the spinal cord of a child.




FIG. 137.—Method of examining for syphilitic osteochondritis of the femur. The initial skin incision is seen over the femur on the right side; in the left limb the bone has been disarticulated. The left hand, holding the bone, is well wrapped in a towel. The longitudinal incision is made with a strong cartilage-knife. Usually this is sufficient even to cut the bone; if not, a small saw may be used.



FIG. 138.—Examination for syphilitic osteochondritis of the femur. After the sawing is completed, one portion of the cut bone is bent back and the osteochondral line is well shown.





FIG. 159.—Method of examining nasal cavities, antrum of Highmore, etc. By means of a knife the uppermost mucous membrane between the lip and the superior maxilla is incised, the upper lip being elevated with the left hand during the incision. Vertical sawing is now done in the median line, and the tooth extracted at the point where the lateral sawing is to take place. The bone-forceps readily bring the desired portion of bone away, or it can be loosened by means of a chisel.



FIG. 160.—Appearance of the part after removal of a portion of the superior maxilla for the purpose of examining the nasal cavities, antrum of Highmore, etc.

gravity slightly greater than that of the cerebral substance, it affords a more general and even support in the subsequent manipulation, thus lessening the liability of damage in the removal of the brain.

The method in detail is as follows: The scalp is incised across the vertex and the flaps are turned forward and backward as in the adult. With scissors having well-rounded points the sutures and dura are cut through well down to the base of the skull. The five flaps thus formed are pulled outward and, if necessary, cut partly across their base with strong scissors. While the brain is being removed the body should preferably be held in the salt solution. Begin by removing the falx cerebri and longitudinal sinus, then the frontal lobes, olfactory bulbs, etc., in the usual order. When the tentorium and falx are cut through, the brain can be pushed out into the solution, where it will float. If it be desired to harden the brain, it is well to place a jar of Müller's or other hardening fluid in the brine and under the brain as it floats therein, care being taken in the transfer to allow as little as possible of the salt solution to enter the jar, though the fluid should afterwards be changed for a fresh supply.

The spinal cord may be removed from the body of a baby with scissors alone, as the parts are easily cut through. The lines for the incisions through the skin and the vertebræ are made in the same manner as in the adult, but neither knife nor saw is required, the scissors being strong enough to penetrate easily the soft bony structures of the vertebral column in a child under fifteen months of age. (Fig. 156.) In babes the spinal cord is relatively much more firm than the brain.

In autopsies on babes suspected of being the victims of hereditary syphilis it is often important to look for the fatty changes produced by that disease at the junction of the cartilage and the bone in the femur. For this purpose a longitudinal incision is made directly over the head of the os femoris and the soft parts are dissected until the bone is reached. The ligaments are then incised and the head is disarticulated. The shaft is held by the left hand securely wrapped in a towel while a perpendicular incision through the cartilaginous head is made down to the bone; should this be much ossified, the incision may be continued with a saw. After sawing for about two inches, a knife is introduced and one segment is broken off. The presence of a yellowish area of fatty degeneration, more conspicuous in the osseous portion than in the cartilage, shows an interference in the nutrition of the part, quite characteristic of hereditary syphilis. (Figs. 157 and 158.)

The centre of ossification of the lower epiphysis of the femur is present at the end of the ninth lunar month of intra-uterine life. In over 700 full-term infants examined by Vibert and Liman it was found to be absent in only sixteen cases. To determine the presence or absence of Bécclard's sign (presence of the centre of ossification in the femur) the knee is forcibly flexed, and the epiphyseal cartilage of the femur is cut in thin sections perpendicular to the axis of the bone until the greatest diameter of the femoral osseous centre is found, which in a child at term will measure from a half to five millimetres across, its red color affording a striking contrast to the gray cartilage, except where putrefactive changes are far advanced. The ossific centre of the tarsal cuboid bone is still more reliable in determining maturity, as this centre appears at the last month of fetal life. Another sign of importance in a full-term child is the presence of eight separate small dental compartments in the inferior maxillary bone, four on each side of the median line, and a large space towards the ramus which has not yet had the partitions divided off, though an attempt at their formation may often be seen. To determine if the child be viable the osseous centre of the os calcis should be examined, as this is first found between the one hundred and ninety-sixth and the two hundred and tenth day of fetal life.

After a careful study of the urine of the new-born, Sabrazès<sup>1</sup> concludes that the secretion of the fetal kidneys possesses hæmolytic properties and is poor in chlorides and phosphates. Some authorities find no phosphates until after two complete days of extra-uterine existence. In the renal tubules are found during the first week of life considerable quantities of precipitated sodium urates, causing in the pyramids yellowish-white lines which converge towards the apex of the papillæ. These are supposed to prove that the infant was born alive. If a child was born before term, meconium is found alone in the small intestine, but if born at term, it is found in the large intestine. The kidneys are lobular, as in ruminants.

The latest investigations lead to the conclusion that puerperal eclampsia arises from a defective excretory power of the mother, usually referable to the kidneys, but that the actual toxin is a fetal product, the added stress of which upon the defective eliminating powers of the mother precipitates the eclamptic attack. Raubitschek<sup>2</sup>

<sup>1</sup> *Hebd. des sciences méd.*, October 5, 1902.

<sup>2</sup> *Z. f. Heilkunde*, vol. xxv, no. 1, p. 10; abs., *Amer. Med.*, March 26, 1904.



believes that the secretion (*Hexenmilch*) frequently found in the mammary glands of the new-born is the result of a necrosis and the separation of epithelial cells in the acini and ducts of these glands, which are thus shown to be at this stage analogous to the sebaceous glands. The secretion of colostrum immediately preceding lactation in the puerperium is of similar origin.

Blummer<sup>1</sup> considers the status lymphaticus to be a definite pathologic entity, probably associated with, if not due to, a condition of intermittent lymphotoxæmia, and at times capable of playing an important rôle in sudden death, fatal anæsthesia, and infection. In some cases sudden death is undoubtedly mechanical and due to asphyxia caused by the enlarged thymus pressing on the trachea.

The following two examples of the many which have come under my personal observation show the necessity of constant vigilance on the part of the practising physician in order that he may draw correct conclusions from the pathologic data presented to his view. Two days after its birth a babe came upon the post-mortem table of one of the hospitals with which I was connected some years ago. The autopsy revealed a completely imperforate rectum, the anus likewise being imperforate. The physician in charge had ordered immediately after birth that a glycerin suppository should be administered morning and evening. The nurse reported each time upon the hospital records the carrying out of the order!<sup>2</sup> In 1896 there came by train to Philadelphia a hard-working colored woman who was about five months pregnant. Feeling her labor pains coming on she took a cab at the Broad Street Station, and was driven to one of our large hospitals, where she was duly delivered of a dead child. The cabman returned to his post, seeking more work. Upon being again employed a five-months' colored fœtus was found on the floor of the carriage. At the coroner's inquest the mother swore that she was unaware of the fact that she had given birth to the child while in the carriage and that no abortion had been performed upon her person. She considered the premature labor as being due to hard work. At the hospital the placenta had not been critically examined, the existence, therefore, of a twin pregnancy not being established at the time of the delivery of the dead fœtus.<sup>3</sup>

<sup>1</sup> *Johns Hopkins Bulletin*, October, 1903.

<sup>2</sup> CATTELL, *Annals of Gynec. and Pædiatry*, September, 1893, p. 759.

<sup>3</sup> CATTELL, *Int. Med. Mag.*, February, 1897, p. 80.

## CHAPTER XIX

### RESTRICTED POST-MORTEM EXAMINATIONS

IN case permission to open the thorax is refused, the diaphragm may be severed from its anterior attachments, and the lungs, the heart, and even the tongue and adjacent parts may be removed *en masse* through an abdominal incision or a laparotomy wound.

Should the avoidance of visible mutilation be imperative, it is possible to examine and, if necessary, to remove both the abdominal and thoracic viscera through the rectum or perineum in males or through these parts and the vagina in females. In the male this procedure is performed in the following manner:<sup>1</sup>

The body is placed on the back, with the buttocks very near the end of the table and the thighs widely separated and flexed upon the body. The scrotum is then well drawn up, and an incision is made from the perineo-scrotal junction to the margin of the anus and down to the bulb. The knife is carried around this and through the subjacent tissue to the pelvic fascia underlying the vesicorectal pouch, without injuring the bladder or rectum. The left arm being bared to the shoulder, the hand is introduced through the incision, and gradually forced up between the parietal peritoneum and the rectus muscles to the diaphragm. The peritoneum may be opened, but the intestines will invest the hand like a tightly fitting glove and make the manipulation more difficult. If unable to perforate the diaphragm with the fingers, a scalpel may be carried up, with the blade flat against the index-finger, and a nick made in the muscle, the knife being then withdrawn and the opening enlarged with the fingers. The lungs may be examined by palpation, any adhesions broken up, and the organs dragged into the abdominal cavity, the roots being severed with a knife, after which they may be removed. The heart can be examined in a similar manner, except that, before it can be moved very far, scissors or a knife will be necessary to sever the large vessels. The kidneys, adrenals, spleen, stomach, etc., may be removed in this manner, but the liver must generally be divided into its lobes in order to get it through the incision. The organs are examined in the usual manner and returned to the body; some wads of oakum may then be pushed into the abdominal cavity and the perineal incision very carefully closed by hidden sutures.

---

<sup>1</sup> H. A. KELLY, *Medical News*, June 30, 1883.

It is also possible to make the examination through the rectum, but the sphincter is left dilated and gaping, presenting a much more conspicuous and unsightly appearance than the perineal incision.

This method is most difficult of accomplishment when the operator's arm measures more than ten or eleven inches around the biceps, especially in subjects of only average size. The work is very arduous, because of the strained and cramped position which the hand and arm must assume in order to pass the promontory of the sacrum. Coplin suggests the use of the photographer's thimble in tearing the tissues within the abdominal cavity.

Access to the interior of the trunk may readily be had from the dorsum by making a longitudinal incision to one side of the spinal column and sawing the ribs close to their vertebral attachments. When the examination is made through the vagina, an oval incision such as is described on page 183 may be made, or a vaginal hysterectomy may first be performed (Figs. 101 to 109 inclusive).

The brain may be removed almost intact (in two or three pieces) by making a transverse four-inch incision across the fifth cervical vertebra, dissecting up the soft tissues, and cutting a V-shaped segment out of the occipital bone by introducing a saw through the foramen magnum and sawing towards the ears and then across transversely. (Fig. 143, *E A F.*) A rapid, but not scientific, method of diagnosing hemorrhage, which also permits of the removal of the brain in small pieces, is referred to on p. 233.

An examination of the bones of the face is sometimes desirable, but the circumstances and conditions under which it may be required are so variable that the method must be left entirely to the judgment of the operator. Disfigurement is so readily noticed that nothing further than a superficial examination should be attempted without the permission of those interested. The simplest and most unobjectionable method of procedure is to introduce the knife through an incision previously made from the ear to the neck and dissect subcutaneously the tissue investing the bony structures. If the bones of the face are to be removed, it may be necessary to make a transverse incision, the point of election being the furrow between the inferior maxilla and the neck.

If the oral cavity must be examined through the orifice of the mouth after rigor mortis has set in, the rigidity may be overcome by placing towels soaked with hot water over the muscles of the jaw. Such appli-



cations repeated for about five minutes usually suffice. Do not use a chisel to pry the jaws apart, as is sometimes recommended, because of the danger of breaking the teeth or knocking them out. As the rigidity rarely returns, it is advisable at the end of the examination to close the mouth with a few sutures through the mucous membrane of the upper and lower lips.

The nasal cavity may be exposed and examined by detaching with a knife the upper lip from the maxilla from within and then removing with a saw such portions of the superior jaw-bone as will afford room for inspection of the parts under consideration (Figs. 159, 160). By the removal of the eye the pituitary body, Gasserian ganglion, etc., are rendered easily accessible. Indeed, it is surprising what extensive dissections may be made in the region of the face and neck in the ways just mentioned, thus affording an opportunity for thorough digital examination of areas not open to ocular inspection.

## CHAPTER XX

### RESTORATION AND PRESERVATION OF THE BODY

WHEN the examination has been completed, the cavities of the body should be thoroughly sponged out, all blood and other fluids removed, and bleeding vessels tied to prevent leakage. The organs should then, as nearly as possible, be returned to their respective positions, and the cavities filled with dry bran, absorbent cotton, sawdust, sea-weed, or shavings, in sufficient quantity to restore the original contour of the body, covering the abdominal contents with old cloth or papers to protect the under surface of the seam. The brain is generally put into the abdominal or thoracic cavity, owing to the great difficulty in returning it to the skull. If several postmortems be made at the same time and place, care should be taken to return the organs to the proper body, nor should a cadaver be used as a convenient receptacle for the disposal of specimens which are no longer of any use. In the case of a child a small bag may be packed with sand or sawdust so as to assume the shape of the brain and placed inside the calvarium; the brain itself, after dissection, is placed in the abdominal or thoracic cavity. It is unwise, however, to permit any member of the family to witness this procedure.

In all private cases it is important to secure the skullcap in position, to prevent the unsightly disfigurement produced when it slips after the scalp has been sutured. A number of efficient methods have been devised, but the one selected usually depends upon circumstances or upon ingenuity. The fossæ of the skull as well as the calvarium may be filled with plaster of Paris, and while the plaster is still soft a short, stout stick of wood is pushed through into the foramen magnum, the upper end extending to the skullcap, which is then adjusted. When the plaster hardens, the calvarium is well fastened in good position. If in removing the skullcap the precaution is taken to crack at least a part of the inner table with the chisel and hammer, projecting pieces of bone are usually left, which interlock and hold the calvaria snugly in position when it is replaced.<sup>1</sup> If the edges of the temporalis have not been too

---

<sup>1</sup> MALLORY and WRIGHT, *Pathological Technique*.

## THE HISTORY OF THE UNITED STATES

The history of the United States is a story of growth and change. From the first settlers to the present day, the nation has evolved through various stages of development. The early years were marked by exploration and settlement, followed by a period of rapid expansion and growth.

The American Revolution was a pivotal moment in the nation's history. It was a struggle for independence from British rule, which resulted in the formation of the United States as a sovereign nation. The revolution was followed by a period of consolidation and the establishment of the federal government.

The 19th century was a time of great change and growth. The nation expanded westward, and the economy began to diversify. The Civil War was a major event that shaped the nation's future. It was a conflict between the North and the South over the issue of slavery, which resulted in the abolition of slavery and the preservation of the Union.

The 20th century has been a period of rapid change and growth. The nation has expanded its influence around the world, and the economy has continued to grow. The Civil Rights Movement was a major event that shaped the nation's future. It was a struggle for equality and justice for all Americans, which resulted in the passage of the Civil Rights Act of 1964.

The 21st century has been a period of rapid change and growth. The nation has continued to expand its influence around the world, and the economy has continued to grow. The 9/11 attacks were a major event that shaped the nation's future. It was a tragedy that resulted in the loss of thousands of lives and the destruction of property.

The future of the United States is uncertain. There are many challenges ahead, including climate change, terrorism, and economic inequality. However, the nation has a rich history and a bright future. The American people are resilient and determined, and they will continue to shape the nation's destiny.

The history of the United States is a story of growth and change. From the first settlers to the present day, the nation has evolved through various stages of development. The early years were marked by exploration and settlement, followed by a period of rapid expansion and growth.





FIG. 181.—Method of sewing up the body.



FIG. 182.—Appearance of body after it has been sewed with horse-hall stitch. The sewing has been done from above downward, and there is no puckering at the point of starting.




FIG. 285.—Slee's method of fixing the skullcap.




FIG. 286.—Author's method of holding skullcap in place. Four holes are drilled in the bones on each side, two to the right and left of the angle in the temporal bone and two in the skullcap just above the angle. Saw-cuts to hold the wire or string are made in the vertex, the string being thrust in and out of the openings and tied at any convenient spot.



FIG. 165.—Method of withdrawing blood from a body previous to its injection with an embalming fluid. *A*, trocar inserted into left auricle of the heart; *B*, stop-cock; *C*, exhaust valve; *D*, outlet valve for air; *E*, syringe. The valve *D* is used when fluid is to be pumped into the body.



FIG. 166.—Injection of body with embalming fluid. *A*, cannula inserted into brachial artery; *B*, tube going down to bottom of bottle containing embalming fluid; *C*, hand-bulb and valve, by the use of which the fluid is forced through the arterial circulation.





FIG. 167.—Refrigerating room. *A*, recording thermometer and middle tier of shelving; *B* and *D*, tiers of shelving; *C*, brine tank; *E*, pipes of refrigeration apparatus.



The skullcap being secured, the scalp is replaced and sutured with glover's or base-ball stitches,—*i.e.*, those made by repeatedly passing the needle from within outward. By careful use of black or dark thread the incision may be so neatly closed as to escape even fairly close inspection. It sometimes happens that by stretching the skin becomes baggy. Should such be the case, a small portion of the hairy scalp parallel to the original incision across the vertex may be removed previous to the sewing.

After the organs are returned, the sternum should be supported by paper, or, still better, by old linen. Bran and fine sawdust are very useful to fill in with, as they absorb the moisture. Oakum makes the sewing difficult. If the organs have been removed through the vagina or rectum, these outlets should be doubly sewed, some absorbent material having first been introduced to prevent leakage.

A round stick or a piece of gas-pipe may be placed in the spinal canal after the removal of the cord, with the upper end pushed through the foramen magnum, especially if any of the vertebræ have been taken away, and plaster of Paris may be poured in until the cavity is well filled. An old cloth or some paper is then placed on top and the whole sewed together. The line of the incision may be covered with a strip of adhesive plaster. As in Gersung's method for the correction of deformities, paraffin at a temperature of about 55° C. may be injected into the body in large quantities and before cooling be moulded into the proper shape.

The abdominal incision is closed by sewing from the pubes to the sternum, passing the stitches from within outward, about three-eighths of an inch from the cut edges and about half an inch apart, alternating on the two sides so that each needle-hole on one side will be midway between two on the opposite side. The twine should be about half a millimetre thick. Both ends of the suture should be securely tied. For the closing stitch it is well to cut the thread near the needle, withdraw one end, and tie in a surgeon's knot. Roughly estimated, the thread required is twice the length of the incision to be closed. Carefully crowd in any extruding fascia and avoid puckering of the part. (Figs. 163 and 164.)

If the mouth has been opened, or any of the tongue removed with the structures of the neck, the lips may be held together by a few sutures passed through the oral mucous membrane.

If any portions of bone have been excised, their place may be sup-

plied by using a properly shaped piece of wood, which is held in position with sutures, wire, or strong cord, or by plaster of Paris.

Lastly the body should be very carefully cleaned and returned to the place and position in which it was found.

The characteristic "post-mortem odor" is very persistent and defies all kinds of soap. It usually results from handling the intestines, and can best be removed by washing the hands with aromatic spirit of ammonia or, in the absence of that, by rubbing them with dry mustard and then washing with soap and water, or, still better, with some of the newer liquid antiseptic soaps.

Ammonia or the aromatic spirit thereof will remove iodine stains, while carbol-fuchsin and other aniline stains yield to a weak solution of sodium hypobromite.

**EMBALMED AND FROZEN BODIES.**—Embalming may interfere with the work of the pathologist, the bacteriologist, and the toxicologist. Fortunately, the old zinc, mercurial, and arsenical combinations have been very largely superseded by formalin, a much more desirable preparation, although it may irritate the eyes, deaden the sensibility of the finger tips, and even produce an eczema of the hands. Arterial embalming is at present more used by undertakers than any other method of preserving the body after death. It is usually practised by opening one of the superficial arteries, as the femoral, carotid, radial, or brachial, and injecting at least two quarts of fluid slowly into the vessel (arterial embalming). (Fig. 165.) It is customary among many undertakers to first aspirate the heart and remove as much blood as possible (Fig. 166), though this procedure is no longer of such importance as formerly, owing to the improvement in the preservative powers of the embalming fluids now in use. Removal of the contents of stomach and intestines is also sometimes practised and materially aids the preservation.

Next in importance is cavity embalming or injection of preserving fluid into the three body cavities. In the abdomen, the instrument is thrust, preferably, through the umbilicus, so that the wound of entrance will not be conspicuous, and efforts are then made to puncture the heart, lungs, intestines, liver, and other abdominal organs. The gas escapes, any blood exuding is withdrawn, and the fluid is afterwards injected, the stream being directed towards the liver. Some embalmers urge avoidance of any injury to enclosed viscera, and remove the contents *per rectum*.



The disadvantages of this method are: first, in cases of abortion with peritonitis there may be considerable difficulty in determining whether the openings were made before or after death; secondly, such punctures may also complicate matters by opening up abscess-cavities, cysts, aneurisms, etc.; and thirdly, in cases of poisoning, besides allowing the stomach contents to escape, the fluid may contain the same substance as that which caused death. Even when formalin has been employed, as in the recent Haines case in New Jersey, the syringe may have been previously used for injecting an arsenical preparation.

The thorax is best filled by injecting the fluid through a long curved trocar passed down through an opening in the trachea. The fluid should first fill both lungs, and afterwards the trocar should be pushed down so that the point pierces the pleura, and these cavities are then to be filled with fluid. The brain cavity is best filled by passing a needle up the nose, breaking the cribriform plate, and then injecting the cerebrospinal cavities and the sinuses. Any excess of fluid passes through the jugular vein to thoracic organs.

The trocar may be passed through the inner canthus of the eyes by the sphenoidal fissure, or through the foramen magnum, thence injecting the brain and cephalic cavity. Fluid may also be previously introduced by lumbar puncture.<sup>1</sup> These methods, though preserving the nervous tissue, are not as efficient as the arterial embalming for the preservation of the body and are apt to discolor or cause swelling around the face.

Nauwerck employs: an injection-syringe having a capacity of five hundred cubic centimetres; long cannulae of different calibers, with pear-shaped ends and with stopcocks or, preferably, with double stopcocks; strong twine; scalpels, scissors, forceps, grooved director, hæmostats, an aneurism-needle, and ordinary needles; basins and buckets; several packages of absorbent cotton; cloths and sponges; and ten litres of a one per cent. watery solution of corrosive sublimate, which may be kept in one-litre bottles. His method of embalming is begun by exposing the lower part of the abdominal aorta and the two iliac arteries. Two ligatures are placed beneath the aorta, about two finger-breadths apart, and the aorta is obliquely incised to allow the entrance of the cannula, which is secured by

<sup>1</sup> *OSUR, Med. Record*, July 9, 1904.

tying the distal ligature over it. The injection into the upper part of the body is then begun carefully and slowly, pausing occasionally when the counter-pressure becomes too great. About three litres are injected, more or less, depending upon the appearance of swelling of the face, seen first about the eyes and chin. The cannula is removed, both proximal and distal ligatures are tied, and the aorta is cut through. In like manner a litre of the solution is injected into each leg through the common iliac artery. A cannula with a double stopcock can be used to inject both the upper and lower parts of the body at the same time. The mesentery is ligated, and the intestines, from the beginning of the jejunum to the end of the sigmoid flexure, are removed, opened, washed out, and put in a one per cent. solution of bichloride of mercury, and later replaced in the abdominal cavity, wrapped in sublimated cotton, or, where practicable, disposed of by cremation. The stomach, duodenum, and rectum are cleaned out with sublimate solution and packed with sublimated cotton. The bladder, vagina, external ear, and nose are similarly treated. The abdominal cavity is carefully wiped with a cloth wrung out of the bichloride solution and dried, and the abdominal incision is sewed. The surface of the body, with the exception of the hair, is also wiped with the solution and dried. If this method fails, Nauwerck injects into the carotid and axillary arteries.

Hewson<sup>1</sup> recommends the following injection for the preservation of human bodies for the dissecting-room:

R Sodium arsenate .....	2 kilogrammes.
Potassium nitrate .....	1 kilogramme.
Carbolic acid .....	150 cubic centimetres.
Boiling water.....	7850 cubic centimetres.
Boil until complete solution, then add	
Glycerin .....	2000 cubic centimetres.
Formalin (40 per cent. solution).....	100 to 150 cubic centimetres.
Thymol, as much as will go into solution, a piece the size of the end of the thumb being sufficient for a carboy of solution.	

About two and one-half gallons of this fluid are introduced into an artery—say the common carotid—by gravity, openings having previously been made in the toes or in several of the veins if they be distended with blood. After injection the body is thoroughly greased,

---

<sup>1</sup> *Phila. Med. Jr.*, October 27, 1900; *Amer. Med.*, February 27, 1904.

covered with paper, bandaged, and placed in cold storage until wanted for dissection.

Frozen bodies should not be thawed hastily by the addition of warm objects, but should be allowed to remain in a warm room for some twelve hours previous to the post-mortem examination. Figs. 167 and 168 show the refrigeration room of the Medical Department of the University of Pennsylvania, planned by Dr. Holmes, in which when teaching in that institution I kept the cadavers used in illustrating my lectures. The bodies were removed during the afternoon preceding the performance of the autopsy the next morning. At the Paris morgue the bodies are frozen at  $-12^{\circ}$  C. to  $-14^{\circ}$  C. immediately upon their arrival; at the end of one day they are placed in a separate apartment having a temperature of  $-4^{\circ}$  C., so as to facilitate their thawing in fifteen to twenty hours. (Letulle.)

**DEATH MASK.**—In the making of a death mask about five pounds each of plaster of Paris and of modelling clay are employed. Nearly all of the clay is rolled out until it reaches thirty inches in length. The head is placed perpendicular to the body upon an old pillow, and then the face and any hairy portions which are to be included in the mask are thoroughly anointed with olive oil or liquid vaseline. Crumpled towels or pieces of paper are now arranged so that when the potter's clay is entwined around the head it will be supported and prevent the liquid plaster of Paris from escaping. Any openings are now filled in with the remainder of the potter's clay, and the interior of the potter's clay anointed with olive oil. Two quarts of water are placed in a bowl and the plaster of Paris is slowly added, the whole being constantly stirred with a large spoon until of a gruel-like consistency, then more of the plaster of Paris is added until it begins to thicken, when it is immediately poured into the hollow cavity to secure the mould of the face. If the plaster of Paris is good it will set in about twenty minutes; the potter's clay is removed, and the cast can be lifted in a single piece from the face. It is then carefully packed in cotton, removed to the laboratory, coated with mastic varnish, and oiled. A plaster-of-Paris cast is now made of the mould in a similar manner, or the mould itself may be sent to an Italian worker in plaster, usually to be found in a city of any size.



## CHAPTER XXI

### DISEASES DUE TO MICRO-ORGANISMS, PARASITES, AND HÆMATOZO.

THE number of diseases known to be due to vegetable and animal parasites is constantly on the increase, the study of tropical disease especially having in recent years received marked attention and added much to our knowledge on this subject. The lesions which are produced by these agencies and found *post mortem* are varied, though rarely characteristic, and require special bacteriologic and histologic training for their study and elucidation.

ACTINOMYCOSIS.—A chronic, infectious disease, which occurs most frequently in cattle (as “lumpy jaw” or “wooden tongue”), but is found also in man; it is characterized by the formation of small nodules, which break down and infiltrate the surrounding tissue. The exciting cause, the *Streptothrix actinomyces* (ray-fungus), is found in the form of yellowish opaque granules,—called sulphur balls,—which measure from one-half to two millimetres in diameter. When these masses are crushed and placed under the microscope, they give the appearance so beautifully depicted (in 1856) by Lebert in his Atlas. The organism is introduced into the body with food, often through the medium of carious teeth. In one case reported the patient had been accustomed to pick his teeth with a straw. The most common locations of the lesions are: I. Alimentary canal. II. Lungs (lesions are usually unilateral). (a) Chronic bronchitis. (b) Miliary nodule formed by masses of fungi surrounded by granulation tissue. (c) These nodules may fuse, forming abscesses and finally cavities. (d) Bronchopneumonia. III. Heart, emboli and localized parenchymatous myocarditis. IV. Thorax. (a) Erosion of vertebræ. (b) Necrosis of ribs and sternum. V. Skin. (a) Subcutaneous abscesses. (b) Chronic ulceration, which may last for years. VI. Primary infection of the brain, liver, and vermiform appendix have been described. The characteristic primary lesion is a small nodule resembling that seen in an anatomical wart. Later there occurs, especially in the lower jaw proliferation of cells into surrounding tissues similar to those seen in osteosarcoma; this is followed by suppuration. The abscesses are at first multiple, spherical, and discrete; later they coalesce and give

reticulated and honeycombed appearance to the part affected. Metastases may occur.

**ANTHRAX.**—An acute, infectious, contagious disease, more common in the lower animals than it is in man, caused by the *Bacillus anthracis*, and having for its characteristic lesion a pustule. Certain animals are predisposed, especially sheep and goats, though the Angora sheep is apparently immune. In man the disease is contracted in certain occupations, as wool-sorting, tanning, etc., and by the ingestion of the flesh or milk of an infected animal. The *Bacillus anthracis* is a rod-shaped micro-organism, from two to twenty-five microns in length, non-motile (thus distinguished from the similarly shaped but motile *Bacillus subtilis*), often united, and grows with great rapidity. Characteristic cultures may be made on gelatin plates at ordinary temperatures. The bacillus is easily killed, but the spores are very resistant. For seven successive years Ziegler was able to produce anthrax in mice by inoculations from similarly prepared pieces of dry catgut which contained the spores. Two sets of lesions are found, depending upon the method of invasion,—by skin or mucous membranes. I. *External Anthrax.*—(1) Malignant pustule. At the site of inoculation appears a papule which rapidly becomes a vesicle; later a brown eschar is formed, surrounded by small vesicles and an extensive area of brawny induration. The neighboring lymphatics are swollen, tender, and hard. (2) Malignant anthrax œdema. This is an extensive œdema affecting the eyelids, the head, arm, and often the entire upper extremity. It may terminate in gangrene, enteritis, peritonitis, or endocarditis. II. *Internal Anthrax.*—(1) Thorax. Very soon after death the upper extremities, both anteriorly and posteriorly, become dark purple, the nails are blackish blue, and dark chocolate-colored fluids issue from the mouth and nose. The cellular tissues of the upper part of the chest are emphysematous and crackle on pressure. On opening the thorax these tissues are often found infiltrated with blood and a gelatinous effusion. The pleuræ contain much serum (two or three pints), the right more than the left. The pericardial fluid is also increased (six or eight ounces). The lungs are engorged with dark-colored blood. Some portions are œdematous, others harder than normal and of a darker-red color. The bronchial glands are swollen, hemorrhagic, and friable. The heart-muscle is dark colored, soft and flabby; the heart may be empty or contain dark, semifluid blood in all its cavities. The lining membranes of the heart and larger blood-vessels are stained a

1. The first step in the process is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the problem.

2. Once the problem is identified, the next step is to define the objectives and goals of the project. This helps to clarify what needs to be achieved and provides a clear direction for the team.

3. The third step is to develop a plan or strategy to address the problem. This involves breaking down the problem into smaller, manageable tasks and determining the resources needed to complete each task.

4. The fourth step is to implement the plan. This involves putting the strategy into action and monitoring progress to ensure that the project is on track.

5. The final step is to evaluate the results of the project. This involves assessing the outcomes against the objectives and goals and identifying any areas for improvement.

1. The first step in the process is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the problem.

2. Once the problem is identified, the next step is to define the objectives and goals of the project. This helps to clarify what needs to be achieved and provides a clear direction for the team.

3. The third step is to develop a plan or strategy to address the problem. This involves breaking down the problem into smaller, manageable tasks and determining the resources needed to complete each task.

4. The fourth step is to implement the plan. This involves putting the strategy into action and monitoring progress regularly to ensure that the project is on track.

5. Finally, the fifth step is to evaluate the results of the project. This involves assessing the outcomes against the objectives and goals and identifying any areas for improvement or further action.

1. The first step in the process is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the problem.

2. Once the problem is identified, the next step is to define the objectives and goals of the project. This helps to clarify what needs to be achieved and provides a clear direction for the team.

3. The third step is to develop a plan or strategy to address the problem. This involves breaking down the problem into smaller, manageable tasks and determining the resources needed to complete each task.

4. The fourth step is to implement the plan. This involves putting the strategy into action and monitoring progress regularly to ensure that the project is on track.

5. The final step is to evaluate the results of the project. This involves comparing the actual outcomes with the objectives and goals to determine the effectiveness of the project and identify areas for improvement.



closed, and the abdomen scaphoid. There is cyanosis of the skin. (1) In very acute cases the intestinal lesions are not characteristic, but the bowel contains large quantities of "rice-water." In more protracted cases the bowel presents a mapped appearance, — some areas hyperæmic and some anæmic, some hypertrophic and others ulcerated. The inflammation is well marked in the Peyer's patches. The serous membrane is sticky and of a rosy color. The blood-vessels are prominent and the body looks thin and shrunken. The mesenteric glands are swollen, soft, and of a reddish color. (2) The stools are largely serous and contain masses of columnar epithelial cells and almost pure cultures of the micro-organism. (3) The kidney is swollen, of a violet hue, and shows the changes of acute diffuse nephritis. (4) The liver shows little alteration except cloudy swelling, with minute areas of focal necrosis. (5) The heart is flabby. Its right side is usually distended with tarry blood. The left heart is usually empty. (6) The lungs are collapsed and show marked congestion at their bases. Pneumonia and pleurisy may develop, and abscesses are not uncommon. (7) There is a decided tendency to the formation of diphtheritic exudate on mucous membranes, particularly in the throat. (8) The coeliac ganglion is hyperæmic or even hemorrhagic (Rokitansky). (9) All the abdominal organs are very dry.

DENGUE.—An acute infectious disease, prevalent in our Southern States, and generally known as "break-bone fever." It is bacterial in its origin; a therapeutic serum being now made like the antitoxin of diphtheria. The large and small joints become red and swollen. There is commonly a rash, but this has no distinctive character. General enlargement of the lymphatic glands is not uncommon. Being rarely fatal, no detailed observations have been made regarding the pathologic anatomy of this disease.

DIPHTHERIA. — An acute infectious, contagious disease, characterized by the presence of the Klebs-Löffler bacillus and of a false membrane. This bacillus is a non-motile micro-organism which, when grown on blood-serum, assumes a great variety of shapes. It is easily cultivated on albuminous media in from twelve to sixteen hours. The bacillus is fairly resistant, and will live for months under favorable conditions. Many other organisms produce a similar membrane, and the identity of this organism with the pseudobacillus of diphtheria, the bacillus of scleroderma, and the organism of ozæna is believed by many, but the subject is still *sub judice*. The presence of the organism

in well persons is a fact of great interest. The forms of the disease are nasal, pharyngeal, laryngeal, and cutaneous. The characteristic lesion of diphtheria is a false membrane, beginning early as a slightly raised, opaque, whitish-yellow spot on the mucous membrane. As a rule, it grows rapidly, becoming thicker, of a grayish or greenish hue, and firmly adherent to the underlying tissues. In the early stages if an attempt be made to remove it, there is left behind a raw bleeding surface. In the later stages the membrane becomes less firmly adherent, soft, shreddy, and somewhat easily detached. The diphtheritic patches may become hemorrhagic, the color being then dirty brown or grayish green. The blood not only infiltrates the submucous layer but also the pseudomembrane. When the submucous layer and the surrounding connective tissue become markedly infiltrated, the inflammation is said to be phlegmonous. There is great swelling and pus soon forms. A retropharyngeal abscess may be of diphtheritic origin. In nasal diphtheria the membrane may be slight in extent or may entirely block up the nasopharynx. It is apt to lead to extension of inflammation to the membranes of the brain. In the pharyngeal form the exudate is usually first seen on the tonsils. It is apt to be very extensive and extend into the mouth, the œsophagus, and even the stomach. In the laryngeal form the amount of exudate is often very great: it may entirely occlude the air-passages and extend to the lungs and the bronchial tubes, even to those of the third and fourth dimensions, but as it extends it gets softer and thinner. In this form the pharynx may be entirely free from membrane. The cutaneous form is somewhat less common; it is apt to occur about wounds, the false membrane being seldom extensive. In nearly all cases of diphtheria there is marked inflammation of the neighboring lymphatic glands and often of the salivary glands. There is apt to be a bronchopneumonia. There are small atelectatic patches surrounded by areas of inflammation. Should the diphtheritic membrane become gangrenous, the process is liable to extend to the lung. Klebs-Löffler bacilli are usually not found, but cocci of various kinds are numerous. Endocarditis is extremely rare, but changes in the fibres of the heart-muscle are comparatively common. The serous membrane often shows ecchymoses. The kidneys always show more or less diffuse inflammation, which may be hemorrhagic, and albuminuria is a constant symptom of the disease. The other organs show the ordinary febrile changes. In malignant cases the micro-organisms may be found in the bladder and the internal

organs. As a rule they do not penetrate below the submucosa at the site of the lesion. Orth describes an enteritis nodularis in which the follicles and Peyer's patches are markedly swollen and hyperæmic. Growths may occur in various mucous membranes, as in the eye, the œsophagus, the vagina, in exstrophy of the bladder, etc.

**ERYSIPELAS.**—An acute contagious disease, characterized by a rash, and due to the *Streptococcus* or *Diplococcus erysipelatis*. The micro-organisms gain entrance through a wound or abrasion of the skin or mucous membrane. Three types of erysipelas are noted,—simplex, ambulans, and phlegmonosum. In uncomplicated forms little more than an inflamed œdema is seen. The micro-organisms may be found *post mortem* in the lymph-spaces and in the zone of spreading inflammation. In severe forms the face is enormously swollen, the eyes are closed, the lips œdematous, the ears thickened, and the scalp swollen. Blebs and vesicles often appear upon the eyelids, ears, and forehead. Small cutaneous abscesses about the cheeks, forehead, and neck are common, while beneath the scalp large quantities of pus may accumulate. There is enlargement of the cervical glands, but this is masked by the œdema. Erysipelas of the phlegmonous type may extend to the intermuscular fascia. It is then likely to be gangrenous, particularly when following hemorrhagic contusions. This form, besides being the cause of acute purulent œdema, may result in emphysematous inflammation when gas-producing germs are associated. Infarcts often occur in the lungs, spleen, and kidneys; these are usually septic in character. Endocarditis ulcerosa is particularly common. Albuminuria is a constant complication, but true nephritis is only occasionally seen. Septicæmia, septic pericarditis, and pleuritis are of comparatively frequent occurrence. Acute atrophy of the liver sometimes occurs.

**FEVER, GLANDULAR.**—An infectious disease of childhood, characterized by marked enlargement of the cervical glands. It is bacterial in origin and occurs between the ages of one and ten years. The disease is rarely fatal. The cervical glands are swollen and softened; they seldom suppurate, and the adjacent skin and mucous membrane show no marked changes.

**FOOT-AND-MOUTH DISEASE.**—Stomatitis aphthosa epizootica is an acute contagious disease, occurring most frequently in cattle and sheep, but found also in persons who come in contact with the disease in animals. It begins as a small vesicle (which is at first clear, later grayish)



on the lips, cheeks, or pharyngeal mucous membrane. When the vesicle reaches a diameter of from one and a half to three centimetres, it bursts, leaving a shallow ulcer, with oval, circular, or irregular edges. The affected mucous membranes are inflamed, swollen, and œdematous, and there is considerable exudate. The lesions are also found on the udder and feet, usually appearing after the eruption in the mouth. The post-mortem appearances are most varied, consisting in œdema, hemorrhagic infiltrations, fatty changes in the parenchymatous organs, etc. Löffler and Frosch consider the disease to be due to an organism so minute that it passes through the finest filters and is not visible with our present methods of staining. A colored illustration of the lesion is seen in Kitt's *Atlas der Thierkrankheiten*, 1896.

**FRAMBÆSIA.**—Yaws is a contagious disease of the skin, characterized by an indefinite period of incubation and the presence of dirty or bright red-raspberry-like tubercles. It is presumably of microbic origin. The eruption begins as a papule, usually at the site of an old wound. In a few days the papules are scattered over the body; they rapidly enlarge and become tubercles, which are generally circular in shape, and vary in size from that of a pin's head to a small apple. The epidermis splits or cracks, exposing a raw granulating surface, which rarely ulcerates. The disease is by some supposed to be a modified form of syphilis. In his excellent work on *Tropical Diseases*, Manson states that the question of their identity is certain to be debated until the respective germs of yaws and syphilis have been separated, cultivated, and inoculated, though he considers them to be specifically distinct diseases.

**GLANDERS.**—A contagious disease occurring most frequently in horses and asses, the exciting cause being the *Bacillus mallei*. Two forms are recognized: (a) Glanders proper. (b) Farcy. (1) Glanders proper is an acute disease, essentially a necrotic alteration (Unna), occurring most frequently on the mucous membrane of the nose and upper respiratory tract. Its characteristic lesion is a node or tubercle, which is at first spherical, later becomes flattened, then breaks down and presents more or less extensive ulcerations which tend to run together. The mucous membrane is swollen, is of a purplish or dark-red color, and there is considerable exudate from the ulcerating surfaces. The process may extend to the lungs, the most prominent lesion being a catarrhal pneumonia, in which the diseased areas show a marked tendency to break down, with the formation of abscesses. An

eruption of papules, which soon become pustular, frequently appears upon the face and about the joints. The cervical glands are usually much enlarged. A dirty-yellow pasty mass of pus in the gastrocnemii is probably due to glanders. Chronic glanders usually occurs in the nose and is often mistaken for a chronic coryza. There are frequently ulcers about the turbinated bones. (2) Farcy may be acute or chronic. The acute form is of the nature of a phlegmonous inflammation at the point of inoculation. The process may be very extensive and lead to rapid suppuration of the surrounding parts. Metastasis to the surrounding tissues is common, accompanied by the formation of abscesses in the muscles. In chronic farcy localized tumors are found, usually in the skin, the subcutaneous tissue, and the muscles. These tumors result in abscesses and may form deep ulcers. The disease in man has been described as a chronic specific pyæmia, characterized by eruptions on the skin and nasal mucous membranes, with frequent intramuscular abscesses.

GONORRHOËAL INFECTION.—Lesions due to the presence of the gonococcus. That organism has been found in the blood, which after death may be fluid or semiliquid and tarry-black in color. Manifestations of the infection include: (1) *Arthritis*.—The inflammation is acute, periarticular, and extends along the sheaths of the tendons. It is a synovitis which rarely becomes purulent. (2) *Conjunctivitis*.—This occurs most frequently in the new-born. It leads to thickening and ulceration of the conjunctivæ; erosions or entire destruction of the cornea may result. The skin of the lids may be destroyed. (3) *Endocarditis*.—An acute form of simple or ulcerative endocarditis, from which pure cultures of the gonococcus have been made. (4) The results of gonorrhœal infection are periurethral abscess, prostatitis, vaginitis, salpingitis, iritis, pericarditis, pleurisy, etc. All these lesions show a marked tendency to suppurative change.

HYDROPHOBIA.—Rabies is a convulsive disease due to the action of the toxins of the bacillus of hydrophobia on the higher nervous centres. The cerebrospinal system shows congestion of the blood-vessels. There are minute hemorrhages, most numerous in the medulla. The mucous membrane of the pharynx is congested and not infrequently covered with blood-stained mucus. This is true of the larynx, trachea, and larger bronchi, also of the lungs, œsophagus, and stomach. Experiments have shown abundant virus in the spinal cord, brain, and peripheral nerves, but it has not been found in the liver, spleen,

When a dog that is supposed to be mad has bitten a human being, the animal should not be at once killed, but permitted to live and kept under close observation until it shows unmistakable signs of rabies. It should then be killed and its body sent to a competent bacteriologist for microscopic study and inoculation experiments on rabbits. While the recent so-called rapid method of diagnosing rabies is not absolutely characteristic of the disease, it affords a most valuable and early means of tentative diagnosis, to be confirmed or disproved by subsequent animal inoculation. The method employed is that of Babès, van Gehuchten, and Nélis, and is as follows:<sup>1</sup> Several intervertebral ganglia or a portion of the bulb are put at once into absolute alcohol, in which they are left for twenty-four hours. They are then transferred for one hour to a mixture of absolute alcohol and chloroform, next put for one hour into pure chloroform, then for one hour into a mixture of chloroform and paraffin, and lastly for an hour into pure paraffin. The sections are put in the oven for a few minutes, then passed through xylol, absolute alcohol, and ninety per cent. alcohol, after which they are stained for five minutes in methylene-blue, according to Nissl's formula, differentiated in ninety per cent. alcohol, dehydrated in absolute alcohol, and cleared in essence of cajuput and xylol. Other methods of preparing the tissues may be used, as the rapid fixation with ten per cent. formalin, subsequent freezing, and staining with hæmatoxylin and eosin. The microscopic changes are chromatolytic and capsular. The "rabid tubercle" of Babès consists in the pericellular accumulations of the embryonal cells described by Kolesnikoff. The prolongations of the cells of the bulbar nuclei are shortened, the nuclei are altered or even obliterated, and the nerve-cells are invaded by the embryonal cells and small corpuscular elements. Atrophy, invasion, and destruction of the nerve-cells of the intervertebral and plexiform ganglia of the pneumogastric take place by cells newly formed from the capsule, which appear between the cell body and its endothelial capsule, in advanced cases the field even resembling an alveolar sarcoma.

INFLUENZA.—The grippe is an acute, epidemic, contagious disease, due to Pfeiffer's bacillus, and characterized by abrupt onset, great depression, and many sequelæ. The bacillus is found in the nasal and bronchial secretions. It is one of the smallest organisms known, non-

---

<sup>1</sup> RAVENEL AND MCCARTHY, *Proceedings of the Pathological Society of Philadelphia*, 1901, p. 93.



motile, and stains well with Löffler's methylene-blue. On culture media it grows best in the presence of hæmoglobin. (1) Lesions of the respiratory form are those of an acute inflammation of the mucous membrane of the upper respiratory tract and bronchial tubes. Lobular pneumonia is common, and is probably due to a mixed infection. Pleurisy is more rare, but may lead to empyema. Tuberculosis is apt to be exaggerated by an attack of influenza. (2) In the gastro-intestinal form the inflammation extends to the mucous membrane of the stomach and the intestines. It is seldom of a severe type. The spleen is usually enlarged in this form. The recent large number of cases of appendicitis is attributed by some to the wide-spread prevalence of this disorder. (3) In the nervous form mild degrees of meningitis and encephalitis are not uncommon. Abscesses of the brain have occurred in severe acute cases. In some epidemics accumulations of pus in the nasopharynx are exceedingly common. *Complications.*—Acute diffuse nephritis is quite frequent. Endocarditis, pericarditis, and thrombosis have been reported. Occasionally purpura is seen and also catarrhal conjunctivitis and iritis. In an autopsy on a child dying from meningitis following the grippe Dr. Kneass isolated for me the influenza bacillus.

**LEPROSY.**—Leprosy is an infectious disease characterized by the formation of a node or nodule, and due to the leprosy bacillus. The *Bacillus lepræ* has many points of resemblance to the bacillus of tuberculosis. It, however, stains more readily, is more easily decolorized, and is present in far greater numbers in the lesions which it causes. (1) The tubercular form starts as a small red spot in the corium, which either disappears or gives rise to the formation of inflammatory nodules of a brownish-red color, somewhat soft in consistency, and resembling a strawberry. The primary lesion is found most frequently in the skin of the face and on the surfaces of the knees, the elbows, the hands, and the feet. It may also involve the conjunctiva and the mucous membrane (particularly the nasal), the cornea, and the larynx. This form of the disease is apt to be exceedingly chronic, the surrounding tissues showing marked fibroid changes. The tubercles at times undergo fatty disintegration and in this way become swollen. (2) In the anæsthetic form the leprosy process gradually involves the peripheral nerves, first causing a perineuritis, then obliterating them and producing marked trophic changes, consisting in necrosis and ulceration with extensive loss of substance, as of fingers, toes, and even

limbs. There is great loss of hair and the face often shows marked ravages of the disease. Death results not infrequently from laryngeal complication or aspiration pneumonia. That leprosy may be cured in the sense of the lesions not advancing is now an established fact. Van Houtum<sup>1</sup> claims to have cultivated successfully the *Bacillus lepræ*, while several investigators have recently given promising reports of the discovery of a curative serum.

**MALTA FEVER.**—Mediterranean fever is a chronic disease, resembling in its clinical course typhoid fever and malaria, occurring most frequently in the Mediterranean region, and due to the *Micrococcus melitensis*. It is often followed by swellings of the joints, profuse diaphoresis, anæmia, orchitis, and neuralgia. Young and previously healthy adults who are unacclimated are most frequently attacked, and it is a serious disease in the British garrisons. The micrococcus is found in large numbers in the spleen. The visceral changes are those common to all infectious diseases with high temperature. The small intestine is usually anæmic except in the upper part, where it may be intensely congested. The mesenteric glands show little change. The spleen is much enlarged and dark in color; its pulp is soft and friable, and sections show an increase in the lymphoid elements. The average weight is eighteen ounces. The liver is congested and its surface on section is pigmented. The kidneys are usually congested and may be slightly hemorrhagic. The agglutinative reaction can be obtained with the micrococcus and the blood of a patient affected with Malta fever. It should be remembered that this disease occurs in our new possessions, and that soldiers and sailors on their return home may bring the affection with them.

**MEASLES.**—Morbilli or rubeola is a markedly contagious disease, attended with a skin eruption and catarrh of the mucous membranes, and due to a micro-organism the identity of which is not yet definitely settled. This affection, as well as scarlet fever and German measles, must be distinguished from Duke's fourth disease, a malady having characteristics in common with all three disorders. Lesage, Canon and Pielicke, Czajkowski, and others have described organisms as causes of the disease. The post-mortem appearances in measles are chiefly those of its complications and sequelæ. The skin, especially about the face, may be swollen and slightly œdematous, and may show

<sup>1</sup> *Journal of Pathology and Bacteriology*, September, 1902, p. 260.



the remains of the characteristic rash, especially in the hemorrhagic type. Desquamation, when present, is in the form of fine branny scales. The gastro-intestinal mucosa is usually hyperæmic; Peyer's patches are frequently swollen, sometimes markedly so. The lungs invariably show evidence of bronchitis, and almost invariably lesions of broncho-pneumonia with areas of collapse; less frequently lobar pneumonia may be found. The bronchial glands are invariably swollen. Pleurisy is less common. In debilitated infants severe stomatitis, cancrum oris, or ulcerative vulvitis may develop. In the middle ear catarrhal inflammation, which may go on to abscess formation, is not uncommon. Of the sequelæ tuberculosis is the most important; it is either miliary or a caseous pneumonia. Severe forms of conjunctivitis and ulcer of the cornea are not uncommon. Nephritis is exceedingly rare. There is cloudy swelling of the organs.

**MUMPS.**—An acute, infectious, contagious disease, characterized by a marked cellular infiltration of the parotid glands, which do not tend to suppurate or to become fibroid, and frequently complicated with metastases to the ovaries and mammary glands in females, and the testicles in males. (*a*) Probably due to a coccus infection. (*b*) Childhood and adolescence. Very young infants and adults are seldom attacked. Uncomplicated mumps is rarely fatal. Of the complications meningitis, acute mania, endocarditis, gangrene, and optic atrophy are the most important.

**PLAGUE.**—An acute, infectious, contagious, epidemic disease, due to the *Bacillus pestis*, occurring usually in the far East, but at present (1904) widely distributed over the earth's surface, and characterized by marked glandular enlargements which tend to suppuration and by a general septic condition. The bacillus was discovered by Kitasato and Yersin. It is a short rod with rounded ends, and is found in the blood, glands, and viscera. Hossack found no buboes in thirty per cent. of his cases in Calcutta in 1900. *Varieties.*—(*a*) Bubonic. (*b*) Pneumonic. (*c*) Septic. (*d*) Intestinal. (*e*) Meningeal. (*f*) Carbuncular. *Lesions:* (1) At the point of inoculation, which usually occurs on the lower extremities, there appears a small spot (plague-corpuscle) which soon becomes a vesicle and then a pustule. (2) Following primary inoculation, the inguinal glands become swollen, succeeded in order by the axillary, cervical, popliteal, and then any of the glands in the body may become affected. The diseased glands swell rapidly and are at first tense and firm to the touch, but soon undergo a



suppurative change, and in rare cases gangrene ensues. It may be stated that it is the periglandular tissue which becomes œdematous and undergoes septic inflammation. (3) Carbuncles may develop in the skin of the legs, hips, and back. Subcutaneous hemorrhages are very common and may also occur in the mucous membranes. (4) The central nervous system, especially the brain, is deeply congested. The brain substance may become softened and the blood-vessels, especially the veins, are engorged. (5) The lungs are deeply congested, especially posteriorly, and are at times the primary seat of the disease. (6) The pericardium contains an excess of blood-stained fluid. The right heart is dilated with black, imperfectly coagulated blood, and the whole venous system is engorged. The heart-muscle is pale and somewhat softened. (7) The stomach and small intestine contain blood or blood-stained fluid. There may be ulceration, but Peyer's patches are not affected. The spleen is greatly enlarged in all cases. (8) The dorsum of the tongue is coated, but the edges, the tip, and often the median raphe remain pink and clean; sometimes, however, becoming red and dry (Hossack). The disease must be distinguished from puerperal fever, septicæmia, pyæmia, smallpox, influenza, cerebrospinal meningitis, diphtheria, erysipelas, measles, gonorrhœa, syphilis, mumps, malaria, scrofulous glands, Hodgkin's disease, etc. In the case of a Chinaman suspected of having the plague, the writer found almost complete occlusion of the prepuce, with a discharge containing the gonococcus, and in the suppurating bubo a fat diplo-bacillus which did not stain by Gram's method.

RELAPSING FEVER.—An acute, epidemic, contagious disease, not found at the present time in America unless imported, occurring in the same class of persons as typhus fever, giving rise to a fever which lasts from five to seven days, followed by relapses, and due to the *Spirochæta* of Obermeier, which are found in the blood only during the paroxysms of fever. This very motile organism is only rarely to be discovered at the postmortem. No characteristic or constant lesions are found after death. The following are sometimes present. (1) If death occurs during the paroxysm, the spleen is large and soft; the pulp is purple. The follicles are enlarged and often obliterated, though they may be gray or whitish yellow in color. Infarcts are not uncommon. (2) The heart is flabby, of a pale dirty-gray color, and very friable. (3) The liver is more enlarged in this than in any other infectious fever. Its color is uniform gray-red. Fatty degeneration

may be marked. (4) The kidneys may retain their normal weight. The renal parenchyma is soft and flabby; the cortical substance is increased and shows cloudy swelling. Hemorrhagic spots or lines radiating to the pyramids are often observed. (5) The lungs may be the seat of pneumonic infiltration, bronchitis, or bronchiectasis. (6) Hyperplasia of the bone marrow has been found. *Complications.*—(a) Pneumonia is frequent. (b) Rupture of the spleen. (c) Nephritis and hæmaturia. (d) Ophthalmia in certain epidemics. (e) Abortion usually takes place. (Osler.)

**RHEUMATIC FEVER.**—(a) Follows exposure to cold and wet. (b) Usually regarded as a coccus infection, though a bacillus has also been described as the etiologic factor. (1) The affected joints are swollen, tense to the touch, and somewhat hyperæmic. The fluid in the joint is turbid, and contains albumin, leucocytes, and a few flakes of fibrin, but rarely pus. There may be slight erosion of the cartilages. (2) Endocarditis occurs in about sixty per cent. of all cases. The verrucose variety is most common. The mitral valve is most frequently involved. (3) Pericarditis may occur, with or without endocarditis. It may be fibrinous, serofibrinous, or, in children, purulent. (4) Myocarditis occurs most frequently in association with endopericarditis. It leads to weakening and dilatation of the heart-muscle, and is the most common cause of sudden death in rheumatic fever. (5) Pleurisy and pneumonia occur in about ten per cent. of all cases. (6) Rheumatic nodules, varying in size from a small shot to a large pea, are found on the fingers, hands, and wrists. They may also occur about the elbows, knees, spines of the vertebræ, and scapulæ. (7) Meningitis is extremely rare. (8) Purpura may be present.

**RHEUMATISM, CHRONIC.**—(1) The synovial membranes are injected. There is usually not much effusion. The capsules, ligaments, and sheaths of the tendons are thickened. There may be erosion of the cartilages. As a result of these changes, the joints are often deformed and ankylosis may occur. (2) Muscular atrophy, especially about the joints, frequently follows. (3) Valvular heart-lesions, due to sclerotic changes, are of common occurrence.

**RUBELLA (RÖTHELN, GERMAN MEASLES).**—This disease is rarely fatal in uncomplicated cases. There is no distinctive lesion other than the rash, which may fade entirely after death.

**SCARLET FEVER.**—(a) The majority of cases occur before the tenth year. (b) Infants and adults are usually exempt. (c) Cocci are fre-

quently found in the throat-lesions and in the blood. Class, of C  
 cago, claims to have isolated a specific coccus, which has also b  
 described by Baginsky. (1) Rigor mortis is usually well mark  
 Decomposition may set in early and develops with exceptional rapid  
 cadaveric lividity usually appearing before death. (2) The blood  
 dark in color, thin, and coagulates imperfectly. The vessel-walls  
 usually stained. (3) Except in the hemorrhagic form the skin at  
 death rarely shows a trace of the rash. (4) In the throat follicu  
 tonsillitis, diphtheritic membrane, or suppuration may be prese  
 Punctate hemorrhages, especially about the mouth, are always  
 served. (5) Catarrhal inflammation of the gastro-intestinal muc  
 membrane is not uncommon. The follicles of the small intestines  
 swollen, red, and may even be hemorrhagic. (6) In severe cases  
 intense lymphadenitis, with much inflammatory œdema, is found  
 the neck. This may lead to suppuration or even gangrene, and in r  
 cases to ulceration of the carotid artery and fatal hemorrhage. (7  
 The kidney lesions are most important. Acute diffuse nephritis  
 present in a majority of cases. It is frequently of the glomeru  
 type and may be hemorrhagic. This lesion is not infrequently f  
 lowed by the changes observed in chronic parenchymatous nephrit  
 (8) Endocarditis, which may be either simple or malignant, is r  
 infrequent. Pericarditis and myocardial changes are less commo  
 (9) The spleen is often enlarged, and shows the changes which cha  
 racterize acute splenic tumor. (10) Hemorrhages into the subser  
 tissues beneath the pericardium, endocardium, and pleura are qu  
 frequent. There is more or less cloudy swelling of all the organ  
*Complications.*—(a) The most important is nephritis. The urine  
 small in quantity, of a high specific gravity, cloudy, and of a da  
 blood-color. It contains large amounts of albumin, free blood, a  
 epithelial cells, with hyaline and epithelial tube-casts. Œdema m  
 be slight or marked; in a few cases œdema of the glottis has caus  
 sudden death. (b) Heart complications are next in importance. The  
 may be endocarditis, pericarditis, or myocarditis. (c) Catarrhal pne  
 monia, more rarely croupous pneumonia or pleurisy, may occur. (d  
 Involvement of the middle ear may lead to thrombosis of the later  
 sinus, meningitis, abscess of the brain, or necrosis *en masse* of t  
 middle ear. (e) Adenitis may result. The glands of the neck are the  
 most frequently involved. There may be great destruction and loss  
 tissue. (f) Arthritis of a rheumatic type or more closely resemblin



the gonorrhœal variety may be found. In the latter affection only one joint is involved and suppuration may supervene. The toxin seems to act especially on the epithelial cells. In one of my cases death occurred in convulsions twenty-four hours after the onset of vomiting and without the appearance of any rash. The diagnosis was confirmed by a sister being attacked with the disease later on.

**SCLERODERMA (HIDE-BOUND SKIN).**—(1) *Circumscribed Form.*—On the skin are found patches varying in size and of a waxy or dead-white appearance. They are brawny, hard, and inelastic. (2) *Diffuse Form.*—This form usually occurs in the extremities or on the face. Gradually a diffuse brawny induration develops. The skin becomes firm, hard, and so closely united to the subcutaneous tissue that it cannot be picked up or pinched. The color may be natural. The skin is commonly glossy, drier than normal, and unusually smooth.

**SMALLPOX.**—(a) Bad hygiene. (b) Improper vaccination. (c) Season, fall or winter. (d) Streptococci are found in the characteristic lesions. Councilman<sup>1</sup> has announced the discovery of a protozoön. (1) The characteristic lesion of smallpox is a rash. On the skin may be seen papules, umbilicated vesicles, pustules, and crusts. A shot-like feel of the papules upon the forehead and wrist is quite characteristic. (2) The rash may also be found upon the mucous membranes from the mouth to the rectum, but on account of the moisture the pocks are not quite so characteristic in these situations as upon the skin. In some cases there is deep ulceration, especially in the larynx, which may be followed by necrosis of the cartilages. (3) Swelling of Peyer's patches is not uncommon. (4) In the hemorrhagic form of smallpox extravasations of blood are found on the serous and mucous surfaces, in the parenchyma of the organs, in the connective tissue, and about the nerve-sheaths. They have also been observed in the bone-marrow and in the muscles. (5) As a rule, the spleen is markedly enlarged, but it may be small, very dark, and firm. The liver shows evidences of parenchymatous inflammation. (6) The heart is flabby and pale. The myocardium shows cloudy swelling and fatty degeneration. It is often dark brown in color and may be firm to the touch. The cavities contain little or no clotted blood, and the arterial trunks are nearly

<sup>1</sup> See ZIEGLER'S *General Pathology*, translation by CATTELL, 1895, p. 39: "It is not impossible that other infectious diseases—for instance, smallpox—are caused by parasites that belong among the protozoa."

empty. (7) Lesions of the kidney are not common. It may show cloudy swelling and areas of focal necrosis, or the pelvis may be blocked with dark clots which sometimes extend into the ureters. (8) Absence of the scar resulting from vaccination is very often noted. (9) The epidermis of the hands and feet may be shed entire. The skin is sometimes plum-colored. (10) The face may be swollen. In black smallpox there may be found hemorrhages in all the numerous membranes and in joints. The cornea may be sunken. *Complications.*—(a) Bronchopneumonia is almost invariably present in fatal cases; lobar pneumonia and pleurisy less commonly. (b) Albuminuria is frequent, but true nephritis rare. (c) Purulent changes in the arteries, bones, conjunctiva, and middle ear are common. (d) Ulcerative laryngitis with œdema sometimes causes death. (e) Myocarditis, endocarditis, and pericarditis are comparatively common. At the postmortem the odor is so characteristic that the disease may be recognized by this means alone. The physician should always vaccinate himself both before and after making an autopsy on a smallpox case.

**SPRUE (PSILOSIS).**—A chronic remittent inflammation of the whole or part of the mucous membrane of the alimentary canal, occurring principally in persons residing, or who have resided, in tropical or subtropical climates. Apparently nothing is known of its origin. At postmortem the thoracic organs, the abdominal viscera, and the tissues generally are found to be much wasted, giving the body a mummified appearance. The bowel is exceedingly thin, and on opening it a thick layer of dirty viscid gray, tenacious mucus is seen. On removing this, areas of congestion, ulceration, pigmentation, or thickening may be found. The mesenteric glands are generally enlarged.

**SYPHILIS.**—Lustgarten and van Niessen have described specific organisms, neither of which has been definitely accepted. *Classification.*—I. *Acquired Form.*—(a) Primary. (b) Secondary. (c) Intermediate period. (d) Tertiary. II. *Hereditary Form.*—(a) Primary. (b) Secondary. The following lesions should be looked for in making a postmortem: (1) The initial lesion or its scar. (2) Lymphatic enlargement, especially of the groin, neck, and elbow. (3) Various skin lesions and thinness of the hair. (4) Mucous patches. (5) Onychia and dactylitis. (6) Gumma in the viscera, skin, subcutaneous tissues, muscles, etc. (7) Parotitis. (8) The bones for periostitis or osteomyelitis. (9) The eye for iritis or choroiditis. (10) The bowels for stricture, especially the rectum. (11) The nervous system for tabes,

dementia paralytica, and other forms of sclerosis. I. The lesions found in the *primary stage* are: (1) The chancre. This begins as a small red papule, usually situated at the junction of the skin and mucous membrane. It gradually enlarges and breaks in the centre, leaving a small ulcer with indurated edges and base. (2) The neighboring lymphatic glands are enlarged and hard. II. *Secondary Stage*.—(1) Cutaneous eruptions of all forms. As a rule, the syphilide is polymorphous, varying in form from an erythema to a pustular eruption. It is symmetrically distributed and of a reddish-brown or copper color. It appears most frequently on the chest, abdomen, and flexor surfaces of the arms. (2) The mucous patch is a softened and macerated epithelium, and appears on the mucous membrane or on the moist regions of the skin. It is most frequently found in the mouth, in the throat, and about the anus. The mucous patch is irregularly shaped, non-inflammatory, and does not discharge pus. (3) The hair of the scalp is decidedly thin. (4) Ulcers may be seen on the tonsils and larynx. (5) There may be warts about the vulva and anus. (6) Iritis is common; retinitis rare. (7) The finger-nails may be diseased, forming dry or moist onychia. (8) Periostitis may be present. III. In the *intermediate stage* there are but few lesions: (1) Gumma of the testicles and (2) choroiditis are the only ones found. IV. *Tertiary Stage*.—(1) The late syphilides show a tendency to ulcerate and destroy the deeper layers of the skin, leaving scars. Rupia may develop. (2) The gummata are the characteristic lesions, and may be hard or soft. The former develop in the internal organs and in the mucous membranes. They most frequently terminate in cicatrization, forming stellate scars which often cause marked deformities. Soft gummata are found in bones, skin, etc. They tend to break down and ulcerate, leaving chronic indolent, often serpiginous, sores. (3) When there has been prolonged suppuration, amyloid degeneration of the liver, spleen, and kidneys often occurs. This is especially true with regard to rectal syphilis in women. (4) *Circulatory System*.—The heart frequently shows sclerotic changes of the valves, especially about the aorta. (5) The blood-vessels present arteriosclerosis or atheromatous changes. (6) In the central nervous system sclerosis of the brain and cord and gummata are common. V. *Congenital Syphilis*.—(1) At birth the infant is usually apparently healthy, but it may present well-marked lesions. (2) There is wasting, and pemphigus is noticed on the hands and feet. (3) The lips may be ulcerated and



the mouth and anus fissured. (4) There is inflammation of the nasal mucous membrane; hyperæmia with papillary infiltration is present and necrosis of the bone may occur. (5) The spleen and liver are enlarged. (6) The lungs may present the lesions of white pneumonic or miliary gummata. (7) The long bones usually show characteristic changes, and the epiphyses may be separated. (8) Later the child looks prematurely old. The teeth are wedge-shaped and the cutting edges notched (Hutchinson's teeth). (9) Eye lesions may be seen, interstitial keratitis. (10) Dactylitis is not uncommon.

*Syphilis of the Brain and Cord.*—(1) Gummata are usually multiple, varying in size from a pea to a walnut. In the cerebrum they occur along the sulci. Heubner describes two forms. In the first variety they are grayish or grayish red in color, soft, and not sharply defined. On section they are moist and exude a small amount of juice. In the second form they are quite hard and dry. Their outline is distinct. On section they may be cheesy and look not unlike tuberculous growths. An arteritis around them exists and causes softening. (2) Gummatus arteritis and sclerosis of both arteries and nerve tissue may exist. (3) There may be softening due to obstruction of the blood-vessels. Several years ago a man was condemned to death for killing a person in cold blood. A commission of experts pronounced him sane. The man committed suicide by hanging, and was found at the postmortem numerous gummata of the brain, situated especially in the right temporal and frontal regions.

*Syphilis of the Circulatory System.*—(1) Gummata are rare. (2) Fibrosis of the heart-muscle is common. (3) Sclerosis of the valve is frequent. (4) Arteriosclerosis, aneurism, and endarteritis obliterans are common.

*Syphilis of the Gastro-Intestinal Tract.*—(1) The œsophagus is rarely affected. Ulceration or stenosis may be present. (2) Ulcerophlegmonous inflammations, or abscesses may be found in the pharynx. (3) Ulcers may occur in the small intestine and cæcum. (4) The rectum is not infrequently the seat of cicatricial contraction. This lesion is most often to be seen in women. The lesions that syphilis produces in the gastro-intestinal tract are (a) chancre, (b) ulcers, (c) localized fibrous patches, (d) gummata, (e) miliary nodules, (f) condylomatous masses.

*Syphilis of the Kidneys.*—(1) Gummata are not infrequent. (2) Acute syphilitic nephritis may occur. (3) Chronic interstitial nephritis

tis is more common. This is a localized nephritis caused by the resultant shrinking and marked irregularity of the surface of the kidney. It is sometimes hard to distinguish it from old infarcts, but the change in color, which in syphilis is gray and in infarcts is brown, is a rather good point of differentiation.

*Syphilis of the Larynx.*—(a) Congenital. (b) Acquired, which may be secondary or tertiary. (1) In the secondary form there is erythema, with symmetrical, superficial, whitish ulcers on the cords or ventricular bands. (2) Mucous patches are occasionally seen. (3) In the tertiary form true gummata may appear towards the base of the epiglottis. These break down, producing deep flask-shaped ulcerations, which may heal by connective tissue that shrinks and produces stenosis. (4) Islands of connective tissue commonly appear between the cicatrices and form inflammatory excrescences. (5) The neighboring cartilages may show necrotic changes. (6) A fatal termination may result from perforation of an artery.

*Syphilis of the Liver.*—(1) In diffuse syphilitic hepatitis there is marked fibrous change. The organ is hard, firm, and resistant. The disease usually begins with a perihepatitis, which frequently causes adhesions to the surrounding structures. With contraction of the fibrous tissue great deformities of the liver become manifest. Capillary bile-ducts may be present in abundance in the cirrhotic portion. (2) The smaller gummata are pale-grayish nodules, the larger ones pale yellowish in color. Usually they are multiple (miliary). Although they may be present in any part of the organ, the most common situation is at the junction of the right and left lobes. Great deformity results from healing and contraction.

*Syphilis of the Lung.*—(1) In white pneumonia of the fœtus the affected lung is heavy and airless. On section it presents a grayish-white appearance (white hepatization). (2) Hereditary gummata are small in size, grayish in color, firm in consistence, and more or less symmetrically distributed throughout the lung. (3) Acquired gummata vary in size from a pea to a goose's egg. They are grayish yellow in color and are embedded in connective tissue. The parts around them are hard and brawny and of a glossy lustre. These gummata may break down and form cavities. This condition is called syphilitic phthisis. (4) There may be a fibrous interstitial pneumonia in which the lesions are hard, large, and pale or dark grayish red in color. The middle of the right lung or either apex is the part most frequently

involved. (5) The pleura is thickened. (6) Endocarditis may extend to the hepatic artery and portal vein.

*Syphilis of the Testes.*—(1) Gummatous growths usually involve the epididymis, which becomes a hard mass, from the size of a bean to that of a walnut. It affects the head more commonly than the body of the epididymis. (2) In interstitial orchitis the progress of the disease is slow. The organ is larger than normal and distinctly harder to the touch. The overlying skin is not adherent and there is no tendency to suppuration.

**TETANUS.**—The bacillus of tetanus is a slender rod usually growing in long threads. It is motile, grows on ordinary media at ordinary temperatures, and is anaërobic. It stains readily, but does not retain the stain very well. (1) The bacilli develop at the site of the wound, which is usually of a penetrating character, and do not invade the blood or organs, except very rarely late in the course of the disease. (2) No characteristic lesions have been found. (3) The condition of the wound depends upon the kind and extent of the injury. (4) The central nervous system shows congestion, with perivascular exudations and granular change in the nerve-cells. Some investigators have found swelling and areas of disintegration in the gray matter of the cord, with exudation of a finely granular material and disintegrated blood. (5) In tetanus neonatorum the umbilicus may be inflamed. (6) The rectus muscle has been found ruptured as the result of a spasm. (7) Death may occur from heart-failure or asphyxia.

**THRUSH.**—This disease is due to the *Oidium albicans*, or thrush fungus. Parts affected: (1) The mouth, tongue, cheeks, etc., are more or less densely covered with minute, slightly raised, white spots, which are quite firm and adherent to the mucous membrane. When scraped off and examined microscopically, the characteristic fungus is seen. (2) Occasionally the fungus invades the oesophagus and grows to such an extent as seriously to obstruct its lumen.

**TUBERCULOSIS.**—Any morbid lesion produced by or through the agency of the tubercle bacillus, which is a rod-shaped micro-organism, measuring in length about one-half the diameter of a red corpuscle and in width two-tenths of a micron. It is bent upon itself, grows best on agar containing glycerin, stains with difficulty, but retains the stain tenaciously. The best method of staining is by carbol-fuchsin and Gabbett's solution. When stained it often has a beaded appearance. It is morphologically similar to the bacillus of leprosy and the



smegma bacillus. Tuberculous lesions are: I. *Acute*.—(a) Miliary tuberculosis. (b) Caseous pneumonia or phthisis florida. (c) Tuberculous ulcerations. II. *Chronic*.—(a) Diffuse tuberculosis, ulcerative phthisis, or caseous tuberculosis. (b) Fibroid phthisis. (c) Cold abscesses. III. *Modes of Invasion*.—(a) Aërogenous. (b) Lymphogenous. (c) Hæmatogenous. IV. *Characteristic Lesions of Tuberculosis*.—(a) Miliary tubercle. (b) Caseation. (c) Cold abscesses. (d) Ulceration. *Characteristics of Tuberculous Lesions*.—(1) Miliary tubercle is a small nodule about the size of a mustard-seed, grayish white in color, semi-translucent, raised above the surface, and primarily adherent to the surrounding structures. (2) In caseation or diffuse tuberculosis two or more miliary tubercles agglutinate, isolating the intervening healthy tissue and cutting off its blood-supply. The necrosed area loses symmetry of shape and arrangement and undergoes fatty degeneration. The area is yellowish in color, soft or firm in consistence, and is surrounded by an inflammatory zone. There is an almost complete absence of blood-vessels. (3) Cold abscess is most frequently found in association with tuberculosis of the vertebræ. It is frequently seen as a "psoas" abscess. The capsule of this abscess is more or less imperfect. It does not present the ordinary characteristics of a pyogenic membrane, the limiting wall being composed mainly of broken-down tuberculous tissue with more or less perfectly formed tubercles. The contents of the abscess are pale and of a somewhat watery consistence, composed mainly of broken-down cells, fatty debris, and water. Bacteriologically the contents of the abscess are usually sterile. V. *Distribution of Tubercles in the Body*.—(a) The lungs are most commonly affected. In two hundred and seventy-five cases out of a thousand autopsies, the lungs were, with two or three exceptions, involved in all. Other organs were affected as follows: (b) Intestines in sixty-five cases, (c) peritoneum in thirty-six, (d) kidneys in thirty-two, (e) brain in thirty-one, (f) spleen in twenty-three, (g) generative organs in twenty, (h) liver in twelve, (i) pericardium in seven, and (j) heart in four. (Osler.) VI. *Fate of Tuberculous Lesions*.—Tuberculous lesions may terminate: (a) In resolution, which is rare. (b) In fibroid changes. This sometimes occurs in the small intestine and may cause stenosis. (c) In caseation or supuration. (d) In calcification. (1) Resolution sometimes takes place when the area of tuberculosis is small, the blood-supply good, and the patient under favorable conditions, especially when leading an out-door

existence. (2) In healing by fibroid change the area affected is first encapsulated and then by gradual pressure and absorption the affected area is removed, leaving a scar. (3) Caseation is by far the most common result of all tuberculous lesions. The process has been already described. Suppuration in tuberculous lesions is the result of the introduction of pyogenic organisms. (4) Calcification is the most fortunate ending of the tuberculous process, and it is estimated by careful observers that seventy-five per cent. of all persons who die after the age of forty years show this form of tuberculosis in their lungs or pulmonary glands.

*Tuberculosis of the Alimentary Tract.*—This form may be: (a) Primary in the mucous membranes. (b) Secondary to disease of the lungs or eating infected food. (c) It occurs rarely through extension from the peritoneum. I. *Mouth.*—(1) Primary tuberculosis, which is usually miliary. The tonsils are affected primarily more often than was formerly supposed. (2) Secondary to tuberculosis of the face, larynx, or lung. It may attack the tongue or cheeks and be miliary or caseous. II. *Æsophagus.*—(1) Primary tuberculosis is very rare. (2) Secondary tuberculosis through extension from the lungs or larynx is comparatively common. (3) The lesions may be miliary, caseous, or ulcerative. III. *Stomach.*—Tuberculosis of the stomach is comparatively rare; Orth never saw a case. IV. *Intestines.*—The lesions occur in the ileum, cæcum, colon, and rectum. The most frequent seat is in the ileum, just above the ileocæcal valve, as it is here that stasis of the intestinal contents occurs and a favorable opportunity is given for the growth of the tubercle bacillus. (1) The large bowel is less frequently involved than the small bowel. (2) Small, firm, gray nodules develop, which soon soften and become yellow in the centre. If cut into at this stage, pus does not exude as in an ordinary abscess, but a thick caseous material may be pressed out. The mucous membrane over these nodules finally breaks down and the cheesy material is erupted. There remains an ulcer with swollen cheesy base and edges (primary tuberculous ulcer of Rokitansky), which soon combines with others and enlarges irregularly (secondary tuberculous ulcer of Rokitansky). Miliary tubercles in the form of small gray nodules now appear at the base and edges of the ulcer and its immediate vicinity. Through the caseation of these, the ulcer enlarges both downward and laterally. The round ulcer becomes a long one, with its longer axis usually at right angles to the long axis of the intestine; it may

extend around the bowel. Hemorrhages may occur, particularly at the edges. The submucosa and muscularis are usually involved, and colonies of young tubercles may be scattered over the serous membrane. Perforation is rare. Gangrene may occur in a very rapidly developing ulcer. Healing sometimes takes place. (3) There may be solitary or multiple areas of cicatricial tissue. (4) *Fistula in ano* is quite common. V. (1) The liver is constantly involved in general tuberculosis. It is pale in color, often fatty, and presenting miliary tubercles or caseous masses which may break down into numerous small abscesses, especially about the bile-ducts. (2) There may be an increase in the connective tissues, leading to tuberculous cirrhosis.

*Tuberculosis of the Brain and Cord.*—(a) Acute miliary infection. (b) Chronic meningo-encephalitis. (c) Solitary tubercles. I. *Acute Miliary Tuberculosis.*—(1) This is usually secondary to tuberculosis of the lungs, bronchial glands, or bones. (2) Miliary tubercles occur most frequently in the pia and arachnoid of the cerebellum, next in the cerebrum, then in the pons. They follow the direction of the blood-vessels. They are apt to lead to obliteration of the vessels and thus cause softening and necrotic changes. Serous, seropurulent, or sero-fibrinous exudate is also present. (3) This acute process may result in acute inflammation of the meninges, principally the pia and arachnoid. It is spoken of usually as acute hydrocephalus. This is most pronounced towards the base of the brain and occurs most frequently in children. I have found tubercle bacilli in fluid removed by Quincke's lumbar puncture. II. *Chronic Meningo-Encephalitis.*—The membranes at the base of the brain are most often involved, next in frequency the optic chiasm, the Sylvian fissure, and the interpeduncular space. The membranes are thickened, firmly adherent, and covered with a fibrinous, purulent exudate. The convolutions are flattened and the sulci obliterated. The cerebral substance is more or less cedematous. The lateral ventricles are dilated and contain a turbid fluid. III. *Tuberculous Tumors of the Brain.*—(1) Solitary tubercles are found most usually about the cerebellum. As a rule, they are attached to the meninges, often to the pia mater. (2) Cerebral softening from pressure is not uncommon. The tubercles vary in size from a pea to a small orange. They are grayish yellow in color, caseous, and usually firm and hard, but the centre may be semi-fluid. They may be surrounded by submiliary tubercles, but are, as a rule, surrounded by a soft translucent tissue. (3) They may calcify.



*Tuberculosis of the Circulatory System.*—(1) Primary tuberculosis of the larger vessels is unknown; secondary lesions are not infrequently found if carefully searched for. (2) In the lungs, brain, and other organs the smaller arteries are usually involved in acute infiltration which leads to thrombosis. (3) Tubercles may develop in the walls of the vessels, particularly the muscularis, and undergo softening, which may result in hemorrhage or a wide-spread distribution of the tuberculous infection.

*Tuberculosis of the Genito-Urinary System.*—(a) Most common in males. (b) Age from twenty to forty years. I. *The Kidneys.*—(1) These organs are frequently the seat of an acute miliary infection, which may be primary or secondary. The disease is most marked in the cortex. It may be limited to the areas supplied by a single blood-vessel. Necrosis and caseation rapidly follow. The miliary tubercles may be seen in a row in the direction of the vasa interlobularia. One or both organs may be affected, but at autopsy both are found to be enlarged. (2) Not infrequently one kidney may be completely destroyed and converted into a series of cysts; these contain a cheesy substance, and lime salts may be deposited in their walls. This is a chronic form of the disease and frequently starts at the apices of the pyramids. (3) The walls of the pelvis may be thickened and cheesy, and the mucous membrane converted into a necrotic ulcerating mass. The ureters are usually thickened, caseous, or ulcerated. II. *The Bladder.*—Tuberculosis here is most common in men. (1) Infection of this organ is nearly always secondary to infection elsewhere, particularly in the pelvis of the kidney. The bladder is small, shrunken, thickened, and surrounded by sclerosed tissue. Ulcer formation is most common. It is lenticular in shape and is surrounded with red mucous membrane. Its seat of predilection is the trigone and fundus. Minute gray tubercles may be seen. In advanced cases ulcers are found. (2) To find tubercle bacilli in the urine centrifugation should be employed, and the precipitate stained in the usual manner for showing these organisms. Care must be taken not to get the smegma bacillus; it is, therefore, advisable that the urine be collected with the strictest precautions. III. *The Testes.*—Infection may occur before the second year. It may be secondary to peritoneal tuberculosis. At times the greater part of the testis is destroyed, its stroma being replaced by a softened or still firm caseous deposit, which may be softened in the centre. IV. Tuberculosis of the *ureters* is very rare. V. *Salpingitis.*—

The oviducts are enlarged, the walls thickened and infiltrated, and the contents cheesy. It is usually bilateral.

*Tuberculosis of the Larynx.*—The lesions may be primary or secondary, usually the latter. The lesions found are: (1) Miliary tuberculosis. (2) Diffuse tuberculosis. (3) Ulceration. In early cases the epithelium is intact, the tubercle starting in the mucosa or sub-mucosa.

*Tuberculosis of the Lung.*—I. *Acute.*—(a) Miliary tuberculosis. (b) Phthisis florida, showing itself as bronchopneumonic tubercles, as lobar-pneumonic tubercles, or as a combination of both. II. *Chronic.*—(a) Ulcerative phthisis. (b) Fibroid phthisis. I. *Acute.*—(1) In acute miliary tuberculosis the lesions are usually present in both lungs. They are frequently so small and transparent that they may be overlooked on macroscopic examination. At other times they are aggregated in localized spots or even become diffuse. In the latter case the lung is increased in size, is firm in consistence, in color is a darker shade of red, is heavier, and crepitates. The pulmonary vessels should be opened with the scissors, and seldom in the pulmonary arteries but often in the veins miliary tubercles can be seen, the infection having been brought through the circulation. Such tubercles may, however, be localized near an old caseous mass, the lymphatic system then being the transmitter. Local spots of emphysema are seen if the condition is not very acute. The tubercles may be peribronchial, perivascular, or in the parenchyma. There is a chronic miliary tuberculosis which presents a combination of lesions of both acute miliary tuberculosis and phthisis and is the connecting link between the two. (2) Phthisis florida, or acute phthisis with formation of cavities, presents a varied appearance. One lobe only, or more or less of the whole lung, may be affected. The organ is heavy; the implicated portions do not collapse and are firm and airless. The pleura is covered with a thin exudate. On section the condition may resemble red or gray hepatization or an intermediate stage between them. In other instances the lung presents a mottled appearance, some areas being intensely congested, others exhibiting a characteristic pale-gray gelatinous exudate, others caseous degeneration and not infrequently cavity formation. Recently affected areas of pulmonary tissue with croupous pneumonia are often seen. II. *Chronic.*—(1) In ulcerative tuberculosis apical involvement in relation to implication at the base exists in the proportion of five hundred to one, according to Kidd.

There are varied lesions. First, there are caseous nodules, which are grayish, white, or yellow in color. Second, cavities may exist, which, if the case is acute, have walls made up of soft caseous masses. In the more chronic cases these walls are replaced by pyogenic membranes of greater or less density, at times covered with granulations. Frequently trabeculæ are seen in the walls; these are the blood-vessels, branches of the lung artery, which have resisted the tuberculous process. The arteries sometimes become aneurismal. Their rupture may be followed by hemorrhage severe enough to cause death. Frequently they are contracted and empty, due to a previous endarteritis or thrombosis. Third, pneumonic areas and evidences of chronic bronchitis are seen. Fourth, some thickening of the pleura is constant. This may be merely an acutely inflamed area rubbing against a corresponding area on the parietal pleura or it may be tightly adherent to it. Not infrequently perforation causes a pyopneumothorax. Fifth, enlarged bronchial glands are discovered which are caseous and often pigmented. Lastly, the bronchi are thickened and the lumina of the smaller ones frequently obliterated. The larger tubes show caseous deposits in the submucous and fibrous coats. (2) In fibroid phthisis the organ is permeated with interstitial overgrowth. In some cases the interstitial change is most prominent; in others the tuberculous process is slightly more marked. The unaffected portions of the lung are largely emphysematous and pigmentation is considerable. The right ventricle and sometimes the whole heart are hypertrophied to a considerable degree.

*Tuberculosis of the Lymphatic Glands.*—(1) Location, most frequent in the cervical chain. (2) Extension opposite that of the lymphatic stream. (Treves.) I. *Chronic Form.*—(1) Hard. (2) Non-adherent. (3) Yellowish white in color. (4) Little tendency to break down and suppurate. (5) Tendency to be localized. (6) Overgrowth of connective tissue considerable. In *tabes Virchow* compared them to a sectioned potato. II. *Less Chronic Form.*—(1) Not as dense. (2) Tendency to become adherent. (3) Gray or grayish white in color. (4) Tendency to liquefy and suppurate. (5) Connective tissue less in amount. (6) Tubercle bacilli more abundant. When tuberculous lymphatic glands are associated with phthisis, they are sometimes found to have opened into a bronchus and caused the disease. This is particularly common in children, and especially when the middle and lower lobes are involved.



*Tuberculosis of the Mammary Gland.*—(a) Female sex. (b) Strumous temperament. (c) Age from the fortieth to the sixtieth year. The seat of predilection is the gland duct. (1) Induration is at first small and very slowly increases in size. (2) The nipple may be retracted. (3) The skin over the gland becomes riddled with sinuses with indurated edges. (4) Associated with lymphatic enlargement, tuberculosis of bone, or other tubercular involvement near the gland.

*Tuberculosis of the Peritoneum.*—I. *Miliary Form.*—(1) On opening the abdominal cavity the serous membranes seem to be covered to a greater or less extent with miliary tubercles, which are present in the mesentery and the omentum also. Frequently the gray nodules follow the distribution of the blood-vessels. (2) In many cases there is little or no inflammatory exudate, although petechial hemorrhages are common. (3) The peritoneum, however, has not its normal shining surface, but is usually pale, somewhat sticky, and lustreless. (4) In many cases there is an effusion of straw-colored or bloody fluid which may amount to a litre or more. It contains a considerable amount of albumin and some cells. The exudate is rarely purulent. II. *Chronic Diffuse Form.*—(1) The abdominal viscera and peritoneum are bound together by tough, firm, membranous bands of organized exudate and the peritoneal cavity is obliterated. (2) The intestinal coils are shortened and contracted, while the mesenteries and omentum are enormously thickened. (3) The capsules of the liver and spleen undergo extreme thickening, varying from a few millimetres to several centimetres. The organs are rough and irregular in outline. III. *Ulcerative Form.*—(1) There is a formation of caseous masses that vary in size from a pea to a marble, and which tend to run together and break down, forming more or less extensive ulcerating surfaces. (2) Adhesions are formed of a serofibrinous or seropurulent character. (3) The new tissues are apt to become pigmented and of a gray or almost black color. (4) The intestinal walls are very friable. (5) Fistulae, opening at various points, are not infrequent.

*Tuberculosis of Serous Membranes.*—There are three groups of cases: (1) Acute miliary tuberculosis, which may develop very rapidly and is accompanied by more or less serous but turbid exudate. (2) A chronic form characterized by exudation, the formation of cheesy masses, and a tendency to suppuration. (3) Cases in which

the tubercles are hard and fibroid, the membranes much thickened, but with little or no fluid exudate. In these cases there may be no visceral tubercles.

*Tuberculosis of the Skin.*—Anatomical warts are small papillary outgrowths frequently seen on the hands of those who make many autopsies. The process is chronic, and, as in the case of one of my helpers in the post-mortem room at Blockley, may give rise to general tuberculosis. The bacilli are few, and are best demonstrated by inoculation of some of the secretion into a guinea-pig. The animal lives for a longer period of time than is usual when it is inoculated with tuberculous material taken from other sources. *Lupus vulgaris* is a cutaneous form of tuberculosis, characterized by the formation of nodules, which tend to break down, producing more or less ulceration. The tubercle bacillus is found in very few numbers. (1) The lesion begins as a small nodule, reddish brown in color and of soft consistence. These nodules vary in size from a pin-head to a cherry and quickly break down and ulcerate. The ulcers are more or less rounded and have a red base covered with granulations. The intervening tissues show diffuse infiltration and fibrous hyperplasia. Warty excrescences may develop in the epidermis or in the floor of the ulcers. The face is the most common seat of the disease. (2) In lupus of the larynx the lesion is surrounded by hyperæmic, œdematous tissue. In the course of time smooth, hard nodules appear, causing great deformity of the parts. Softening and ulceration give the larynx a worm-eaten appearance. The disease follows the lymphatic channels.

*TYPHOID FEVER.*—The intestinal lesions are: First week, intense catarrhal inflammation of the mucous membrane of the intestines and in the first few days only moderate swelling of the follicles. Towards the end of this week, however, there is more decided medullary swelling. Second week, the medullary swelling goes on to resolution or formation of eschar or, third week, ulcer formation. In the fourth week there is beginning cicatrization. The lesions are most marked in the lower ileum, but they also exist in the cæcum and large intestines, rarely in the jejunum. Hyperplasia of the mesenteric lymphatic glands and the spleen develops early in the disease. Cloudy swelling and fatty degeneration of the heart, liver, and kidneys may be present. Waxy degeneration and bleeding in the voluntary muscles should be looked for. Other lesions are lymphoma of the liver, acute nephritis, bleeding of the skin, hypostatic or catarrhal pneumonia,

purulent bronchitis, perforation, and peritonitis. The Widal test and the diazzo-reaction may be determined *post mortem*. Paratyphoid or paracolon infections are more common than was formerly supposed, and furnish most interesting cases for thorough study.

**YELLOW FEVER.**—The chief lesions are: (1) Bleeding from the mucous membranes. (2) Tarry blood. (3) High-grade fatty degeneration of the liver. (4) Acute hemorrhagic inflammation of the stomach and intestinal mucous membrane. (5) Icterus. The interesting work done by Reed, Carroll, and Agramonte in Havana, in showing that this disease is dependent on the *Stegomyia*, a variety of mosquito, is one of the most important contributions to medical literature of the past decade. The *Bacillus X* of Sternberg and the *Bacillus icteroides* of Sanarelli are by some supposed to be identical, by others not to be the cause of yellow fever. There is an interesting illustrated article on this subject in the *New Orleans Medical and Surgical Journal* for January, 1902.

#### PARASITES.

**PEDICULI.**—(a) *Pediculus capitis*.—The female louse measures from one and eight-tenths millimetres to two millimetres in length, the male being somewhat smaller. The darker the skin of the person infested the darker is the color of the parasites. So marked is this peculiarity that some writers are of the opinion that different species affect different races. The ova are grayish glistening specks enclosed in a membrane firmly adherent to the shaft of a hair not far from its root, and coming off at an acute angle, with the opening away from the scalp after the exit of this parasite. Considerable irritation is caused by these animals, and when this is severe the hair on the back of the head may be found matted with soft yellow crusts. The scalp is covered with moist red granulations. The cervical lymphatic glands posteriorly are enlarged. This condition is most frequently seen in children. (b) *Pediculus pubis*.—It differs slightly from the above in that it is smaller and infests regions, as the axillary, the pubic, and the pericocular, where the hair is short. (c) *Pediculus corporis* is the largest form of the parasite. It lives in the clothing, when not in search of food on the body. By its constant irritation it causes dermatitis, and if present for a long time, pigmentation and thickening of the skin. (d) *Cimex lectularius* (common bedbug). (e) *Pulex irritans* (the common flea). (f) *Pulex penetrans* (sand-flea, jigger). The latter



is common in tropical and subtropical countries. It is smaller than the common flea. It burrows under the skin and produces a pustular swelling. (g) *Sarcoptes (Acarus) scabiei*.—The female itch-mite is .45 of a millimetre long and .35 of a millimetre broad; the male is about one-half the size. Its color is pearly white. The burrow in the skin, wherein may be found the excrement and the eggs of the parasites, is about one centimetre in length, and is present where the skin is moist, as in the webs of the fingers and toes. Cutaneous lesions result from the scratching instigated by the irritation caused by the parasite.

CESTODES.—*Intestinal Cestodes*.—(a) *Tænia solium* in the mature form may reach to twelve feet or even more in length. It is composed of numerous segments about one-third of an inch long and averaging a fourth of an inch wide. The head is very minute, being no larger than the head of a pin. In front is a rostellum and at the base of this is a fringe of hooklets. It has four suckers. The worm is hermaphroditic. When mature thousands of ova are passed by the rectum. The embryo has six hooklets. It penetrates the walls of the stomach and burrows into the tissues of the animal that has swallowed it. (b) *Tænia saginata* is larger, longer, and of more frequent occurrence than the preceding. The head is nearly square and measures more than two millimetres in breadth, but has no hooklets. The segments are larger than those of the *Tænia solium*. The reproductive organs are on the ventral aspects of the segments in the median lines. (c) The *Bothriocephalus latus* is larger and longer than any of the flat worms. In the mature state it is twenty-five feet or more in length. It has no hooklets, but is furnished with slit-like fossæ on the head, which act like suckers. The larvæ develop in the peritoneum of fish. (d) *Tænia flavopunctata* is very rare. It is about sixteen centimetres long. (e) The *Cysticercus cellulosæ* is the larval form of the *Tænia solium*. It is found in the muscles, brain, cord, peritoneum, or almost any other tissue of the affected animal. The surrounding capsule is frequently calcified. In the making of many autopsies it is surprising how few tænia are found in the intestinal tract. My experience is limited to but two cases. One of these was that of a man who committed suicide with opium. Two *Tænia saginata* were found, the head of the first one being firmly attached beneath a fold of one of the valvulæ conniventes high up in the jejunum and the other five or six feet farther down the intestine, the segments of both worms then continuing on down to near the ileocæcal valve.

NEMATODES. — (a) *Ascaris Lumbricoides*. — It is a cylindrical worm with both ends pointed. The female is from ten to sixteen inches in length, the male considerably smaller. It is brownish yellow, reddish, or white in color. The head ends in three lips. (b) The *Oxyurus vermicularis* (seat-worm) is a very small round worm, about ten millimetres long. (c) The *Trichina spiralis* in the mature state lives in the intestine; in the immature state in the muscles. The embryo is surrounded by a capsule, which quickly calcifies. Under the microscope the embryo can be seen coiled up in its capsule; it is less than a millimetre in length. (d) The *Anchylostomum duodenale* lives in the upper part of the intestine. The female is the larger, and varies from ten to sixteen millimetres in length. At the anterior portion of its head are hooklets, with which it attaches itself to the intestinal walls. It is frequently associated with Egyptian chlorosis. Stiles and Harris have recently called attention to the wide distribution of uncinariasis in the South; the disease may be readily recognized by finding the ova in the fæces. (e) The embryo of the *Filaria sanguinis hominis* is a round worm one-seventy-fifth to one-one-hundredth of an inch long. It is enclosed in a delicate sac. It circulates freely in the blood, but only at night. The adult parasite is located in the lymphatic vessels and is three or four inches in length. According to Manson, it is introduced into the body by the mosquito.

DISTOMIASIS. — (a) Liver-flukes. (b) Blood-flukes. These worms are lanceolate in shape, quite flat, and possess a distinct head and neck. They are three-fourths of an inch long and about half an inch broad. The color is dull brown. The female blood-fluke has a grooved channel posteriorly for the reception of the male. They have two suckers, one near the mouth and the other near the ventral portion of the body. The liver-fluke infests the upper intestine and the bile-ducts. It causes the "liver-rot" in sheep. The blood-fluke is found chiefly in the portal system and the veins of the bladder. The ova may be seen in the urine as elongated ovoid bodies, sharply pointed at one extremity, and containing black pigment. They can easily be seen with a low power of the microscope. Parasitic hæmoptysis now occurs in America as well as in Asia, and is due to the *Paragonimus Westermanii*. The eggs are found in the sputum, the fluke measuring from eight to sixteen millimetres long by four to eight millimetres across.

MYIASIS.—By this term is meant a condition in which a diseased

part becomes "living," as it is called. It is caused by the larvæ of certain flesh-flies, of common house-flies, or of the bot-flies of oxen or sheep. The ova of these flies may be deposited in the nostrils, ears, conjunctiva, open wounds, or even in the vagina during the puerperium.

**ECHINOCOCCUS DISEASE.**—A parasitic disease, found most frequently in those countries, as Iceland and Australia, where the dog lives in intimate association with man; it is characterized by the formation of endogenous or exogenous multilocular cysts in various portions of the body. The *Tania echinococcus* is a very small, thread-like tapeworm (length from three to six millimetres), having only three segments. The head has four suckers, a rostellum, and a double row of hooklets. The adult worm is found in the dog. The embryos (scolices) are found in the ox, hog, sheep, horse, and man. *Distribution in Man.*—(a) Liver (most common). (b) Lung and pleura. (c) Intestinal tract. (d) Kidney, brain, etc. The embryo, freed from the cyst by digestion in the stomach, burrows through the intestinal wall and is carried to the various organs; it then loses its hooklets and is gradually converted into a cyst (hyatid) having two walls, external laminated, internal granular or parenchymatous, containing blood-vessels and muscle-fibres. The interior is filled with a clear non-albuminous fluid, specific gravity 1005-1009, usually containing sugar and hooklets. From irritation of surrounding tissues a fibrous capsule generally develops on the outside. The cysts vary in size from that of a small pea to that of a child's head. From the inner (parenchymatous) layer may develop brood capsules, which in their turn produce numerous scolices. The cysts grow slowly; when the embryo dies, the whole becomes calcified. Sometimes the cysts suppurate; occasionally they rupture into adjacent structures.

#### HÆMATOZOA.

**MALARIA.**—This widely distributed and much-studied disease is due to a true hæmatozoon, transmitted to man by the bite of the *anopheles* mosquitoes. Three varieties have been described: (a) Tertian. (b) Quartan. (c) Æstivo-autumnal. *Classification.*—(a) Acute malarial fever, which may be quotidian, tertian, or quartan. (b) Pernicious malaria. (c) Chronic malarial cachexia. In the blood of the cadaver the plasmodium is seldom visible, but it may be found in sections of the brain, liver, and spleen. (1) Cases of simple malarial fever



are rarely fatal. The blood shows disintegration of red corpuscles and an accumulation of pigment is thereby formed. The spleen is enlarged, dark in color, and may show pigmentary deposits. (2) In pernicious malaria the blood contains enormous numbers of the parasites. The red corpuscles are in all stages of destruction and the serum is tinged with hæmoglobin. The spleen is moderately enlarged. The pulp is soft, chocolate-colored, and turbid; it contains large numbers of red corpuscles and parasites and the amount of pigment is greatly increased. The liver is swollen and presents areas of focal necrosis and capillary thrombosis. Pigmentary deposits are also common. The kidneys present more or less parenchymatous change with only moderate pigmentation. (3) In malarial cachexia the blood presents all the characteristics of an advanced anæmia, often distinguishable from pernicious anæmia only by the presence of the parasite and icterus. The spleen is greatly enlarged: it may weigh from seven to ten pounds. The organ is firm and resistant to the knife. The capsule is thickened and the parenchyma brownish or slate-colored, with areas of pigmentation. The kidneys are enlarged and of a grayish-red color. The peritoneum is thickened, opaque, and of a deep slate-color; the gastric and intestinal mucous membrane may have the same hue. The gray matter of the brain is of a deep reddish-gray color or in very chronic cases a chocolate-brown. The meninges are congested. (4) Among accidental and late lesions is cirrhosis of the liver. Very extensive pigmentation may occur. Pneumonia is believed to be common; moderate albuminuria is frequent; acute nephritis is not uncommon; chronic nephritis may follow long-continued or repeated infection. Rupture of the capsule of the spleen may occur, followed by bleeding into the peritoneum and even peritonitis. In pernicious malaria the brain may show thrombosis, due to the parasites, with secondary softening of the surrounding tissue. The same thing may be found in the gastro-intestinal mucosa and be followed by superficial ulceration. There may be advanced fatty degeneration of the heart.

PSOROSPERMIASIS.—A condition produced by the presence of oval, transparent bodies belonging to the coccidia, to which class the malarial organism also belongs. I. (1) In the majority of cases of the internal form the psorosperms have been found in the liver. (2) Whitish growths resembling tubercles and containing the coccidia have been found upon the peritoneum, omentum, and pericardium. (3) Similar masses are sometimes seen in the ileum, liver, spleen, and

is common in tropical and subtropical countries. It is smaller than the common flea. It burrows under the skin and produces a pustular swelling. (g) *Sarcoptes (Acarus) scabiei*.—The female itch-mite is .45 of a millimetre long and .35 of a millimetre broad; the male is about one-half the size. Its color is pearly white. The burrow in the skin, wherein may be found the excrement and the eggs of the parasite, is about one centimetre in length, and is present where the skin is moist, as in the webs of the fingers and toes. Cutaneous lesions result from the scratching instigated by the irritation caused by the parasite.

**Cestodes.**—*Intestinal Cestodes.*—(a) *Tænia solium* in the mature form may reach to twelve feet or even more in length. It is composed of numerous segments about one-third of an inch long and averaging a fourth of an inch wide. The head is very minute, being no larger than the head of a pin. In front is a rostellum and at the base of this is a fringe of hooklets. It has four suckers. The worm is hermaphroditic. When mature thousands of ova are passed by the rectum. The embryo has six hooklets. It penetrates the walls of the stomach and burrows into the tissues of the animal that has swallowed it. (b) *Tænia saginata* is larger, longer, and of more frequent occurrence than the preceding. The head is nearly square and measures more than two millimetres in breadth, but has no hooklets. The segments are larger than those of the *Tænia solium*. The reproductive organs are on the ventral aspects of the segments in the median lines. (c) The *Botriocephalus latus* is larger and longer than any of the fat worms. In the mature state it is five feet in length. It has no hooklets, but is furnished with suckers like the head. The larva has the head of fish. (d) *Tænia dioxyli* is very rare. It is about one inch long. (e) The *Cysticercus cellulosus* is found in the muscles, and any other tissue of the affected animal. In the making of mummies frequently calcified. In the making of mummies few tenia are found in the intestines. One of these was noted in but two cases. Two *Tænia saginata* were found in a mummy with opium. Two *Tænia saginata* were found in a mummy, the first one being firmly attached beneath the jejunum and the second one high up in the jejunum. The segments of the first one were found down the intestine, the segments of the second one down to near the ileocecal valve.



NEMATODES. — (a) *Ascaris Lumbricoides*. — It is a cylindrical worm with both ends pointed. The female is from ten to sixteen inches in length, the male considerably smaller. It is brownish yellow, reddish, or white in color. The head ends in three lips. (b) The *Oxyurus vermicularis* (seat-worm) is a very small round worm, about ten millimetres long. (c) The *Trichina spiralis* in the mature state lives in the intestine; in the immature state in the muscles. The embryo is surrounded by a capsule, which quickly calcifies. Under the microscope the embryo can be seen coiled up in its capsule; it is less than a millimetre in length. (d) The *Ancylostomum duodenale* lives in the upper part of the intestine. The female is the larger, and varies from ten to sixteen millimetres in length. At the anterior portion of its head are hooklets, with which it attaches itself to the intestinal walls. It is frequently associated with Egyptian chlorosis. Stiles and Harris have recently called attention to the wide distribution of uncinariasis in the South; the disease may be readily recognized by finding the ova in the fæces. (e) The embryo of the *Filaria sanguinis hominis* is a round worm one-seventy-fifth to one-one-hundredth of an inch long. It is enclosed in a delicate sac. It circulates freely in the blood, but only at night. The adult parasite is located in the lymphatic vessels and is three or four inches in length. According to Manson, it is introduced into the body by the mosquito.

DISTOMIASIS. — (a) Liver-flukes. (b) Blood-flukes. These worms are lanceolate in shape, quite flat, and possess a distinct head and neck. They are three-fourths of an inch long and about half an inch broad. The color is dull brown. The female blood-fluke has a grooved channel posteriorly for the reception of the male. They have two suckers, one near the mouth and the other near the ventral portion of the body. The liver-fluke infests the upper intestine and the bile-ducts. It causes the "liver-rot" in sheep. The blood-fluke is found only in the portal system and the veins of the bladder. The ova are seen in the urine as elongated ovoid bodies, sharply pointed at one extremity, and containing black pigment. They can easily be seen at a low power of the microscope. Parasitic hæmoptysis now prevails in America as well as in Asia, and is due to the *Paragonimus westermani*. The eggs are found in the sputum, the fluke measuring from six to sixteen millimetres long by four to eight millimetres wide.

S.—By this term is meant a condition in which a diseased



kidneys. The liver may be enlarged and contain caseous foci which are surrounded by areas of congestion. (4) The spleen may be similarly affected. II. (1) In cutaneous affections the lesions closely resemble those of tuberculosis of the skin. They occur in Paget's disease of the nipple and by some are believed to be its cause. (2) A case has been reported in which at autopsy nodules were found in the lungs, adrenals, testicle, spleen, on the surface of the liver, and on the pleuræ. Great numbers of psorozoa were found in the lesions. (3) Successful inoculations were made into rabbits and dogs.

TRYPANOSOMA.—Four animal diseases are caused by varieties of trypanosomes, nagana, surra, *mal de caderas*, and dourine. Recently Nepveu, Dutton, and others have found them in man, and the sleeping sickness and the so-called trypanosoma fever (probably different stages of the same disease) are due to the entrance into the blood and cerebrospinal fluids of *Trypanosoma gambiense*. These organisms are transmitted to human beings by tsetse flies (*Glossina palpalis*). Monkeys, dogs, cows, horses, and rats are also susceptible to certain species. *Trypanosoma Lewisii* infests the rat and is transmitted by flea and lice. *Trypanosoma Evansi*, *Brucei*, and *Equinum equiperdun* attack horses and the *Trypanosoma disci* infests birds. The tsetse fly carries the human parasite and the *T. Brucei*. These parasites are flagellated protozoa, fusiform in shape, several times larger than a blood-corpuscle. On one side is an undulating membrane, which extends from the centrosome along the margin to the anterior end of the body, where it becomes a true flagellum. The parasite is non-sexual, reproducing by longitudinal division. It lives in the blood serum and attacks the red cells. The tsetse fly can carry the infection from sick to healthy up to forty-eight hours after having fed. Lately artificial cultivation of the *Trypanosoma Lewisii* from the rat to the hundredth generation has been made by McNeal and Novy.<sup>1</sup> Thus, the first strictly pure cultures of a pathogenic animal parasite have been obtained. The agglutination reaction occurs under proper conditions. Coplin<sup>2</sup> showed these organisms from Blockley rats, and also a slide of the lungs showing pneumonic changes. Ehrlich and Shiga, experimenting with the organism, found that the stain trypan red was able to destroy the trypanosomes in mice and to protect them

<sup>1</sup> VAUGHAN'S *Dedication Volume*.

<sup>2</sup> *Phila. Path. Soc.*, December 10, 1903.

against relapses. This stain they found equally effective when given by way of the stomach. The stain is the combination of one molecule of tetrazotized benzidin monosulphate and two molecules of sodium naphthylamindisulphate. The skin is well reddened in eight minutes, reaches a maximum color in twelve hours, and tint remains for from six to eight weeks, longer in the internal organs. This stain has little or no effect on rats, guinea-pigs, or dogs.<sup>1</sup> The Leishmann-Donovan bodies were first found in the spleen. The parasites are by no means uncommon in the tropics, and are situated intracellularly in large mononuclear macrophages. Rogers has recently announced that trypanosomes develop upon culture of these bodies.

---

<sup>1</sup> *Berl. klin. Wchnschr.*, 1904, March 28, p. 329, and April 4, p. 362.

## CHAPTER XXII

### THE PRESERVATION OF TISSUES FOR MICROSCOPIC AND MACROSCOPIC PURPOSES<sup>1</sup>

WHEN tissues are to be preserved for microscopic study, the method of fixing and hardening them should be decided upon at the time of their removal from the body. The object to be obtained by fixation and hardening is permanently to solidify the structural elements of a part as nearly as possible in their original form and situation. All our present methods, however, fail to give an accurate picture of the living cell, and not enough attention is now paid to the microscopic examination of unstained fresh scrapings removed during the performance of the autopsy. The use of as perfectly fresh tissues as possible is essential, for many structural details disappear on molecular death. Fortunately, this does not occur until several hours after molar death, so that it is often possible to obtain tissues to all intents and purposes still living. Special attention should be paid to those tissues which have been stained or hardened by reagents during the lifetime of the body under examination, as the experimental staining of tissue during life affords a most inviting field of original investigation.

The method of wrapping tissues in paper or cloth and transporting them to a distance is only to be regarded as a last resort. When this is done, pieces of sufficient size to insure preservation of their interior intact are enveloped in an abundant supply of clean cotton (antiseptic gauze causes markings on them), moistened very slightly with a bichlorid-tablet solution, and thoroughly protected from pressure; these segments are cut down to a proper size before they are put into

---

<sup>1</sup> Based on the works of LEE, *The Microtome's Vade Mecum*; MALLORY AND WRIGHT, *Pathological Technique*; APÁTHY, *Die Mikrotechnik der thierischen Morphologie*; FISCHER, *Fixirung, Färbung, und Bau des Protoplasmas*; SZYMONOWICZ, *Lehrbuch der Histologie*; STÖHR, *Text-book of Histology*; BÖHM AND VON DAVIDOFF, *Text-Book of Histology*; *Lehrbuch der Klinischen Untersuchungsmethoden*, and the *Encyklopädie der mikroskopischen Technik*, 1903. It is to be hoped that the ultramicroscope which will reveal particles with a linear diameter of .000001 millimetre will render valuable service. Working with this instrument, RAEHLMANN (*Deut. med. Wchnsch.*, March 24, 1904) found actively motile bacteria in the aqueous and vitreous humors of an eye enucleated for sympathetic ophthalmia. Zeiss has just put on the market the ultraviolet microscope.



the fixing agent in the laboratory. I sometimes cut the pieces in different shapes, each piece representing an organ or part, and always use the same shape for the same part.

Bottles containing the more common fixatives should be ready, and as soon as the tissues are exposed and described—before the part becomes distorted, fluids escape, or surfaces dry—they should be cut with a clean, sharp knife into pieces about two centimetres in length and breadth and one centimetre thick. Sections of organs should include their characteristic structures,—cortex, capsule, hilum, endocardium, etc. Sections of tumors should be taken from the centre, where degenerative changes are most marked, and from the growing peripheral margin, if possible including some normal tissue; this is of especial importance in the case of malignant tumors. Mucous and serous membranes are pinned out on cork, or wood that will give no stain when soaked in the preservative fluid to be used, with their secreting surfaces uppermost. Muscle-fibres are best preserved by being tightly stretched upon and tied at the ends of a piece of wood. The segments of tissue, without being touched by either fingers or forceps, are lifted on the blade of the scalpel and dropped immediately into a bottle containing an amount of fixing fluid far in excess of their bulk. Of energetic fixatives, such as Flemming's or Hermann's, about fifteen times the volume of the object introduced will suffice, while of milder fluids, like the bichromate of potassium or picric acid solutions, fifty times such volume should be employed.

If the different tissues are distinguishable macroscopically, they may be placed in the same jar; if not, separate bottles are better. Tags may be attached, the writing being done with a lead-pencil, so as not to be acted upon by the usual preservatives. The jars are labelled with the date, the number or name of the autopsy, and the fixative used. It is often of importance to add the exact locality from which the pieces have been removed and the plane on which they are to be cut when placed in the microtome.

The fluid should always be changed after it becomes turbid; or in the case of alcohol or formalin, preferably after three hours, whether it is turbid or not. If the specimens are to be sent away, they should not go until the fluid remains clear; if the time necessary for transportation exceeds that of the proper action of the fixative, they should be worked on up to 80 per cent. alcohol and shipped in that fluid, firmly packed in absorbent cotton.

The choice of a fixing agent is determined by the nature of the object to be preserved and the purpose for which the investigation is undertaken. The characteristics of different pathologic conditions are better brought out in some fixatives than in others. Thus, fatty degenerations are well preserved by an osmic acid, bichromate, or formalin solution; œdematous and parenchymatous changes, by corrosive sublimate; fibrin and hemorrhagic conditions, by absolute alcohol, etc. Moreover, different tissues require different treatment; the fixation of a lymph-node is quite a different matter from that of a retina. Then the purpose for which the examination is made will largely influence the choice. If it be simply a question of general diagnosis, Orth's fluid and alcohol will answer every purpose; by the use of alcohol we can preserve the specific staining properties of micro-organisms and hæmoglobin and various important chemical reactions, and by the use of Orth's fluid colloid and mucoid material retain their transparency, fat is preserved, etc. If we undertake the investigation of pathologic processes and the comparison of abnormal with normal cellular anatomy, then special fixatives must be used.

The advantages and disadvantages of the fixing solutions most in use will first be given, next a list of pathologic conditions and the solutions best calculated to preserve their characteristics, and finally a list of staining solutions requiring certain fixatives for their use.

**FIXATIVES; INSOLUBILITY.**—To preserve soluble cell contents they must first be rendered insoluble, and the transformation must be equable throughout. The colloid or fluid material must harden homogeneously and enclose the more solid structures without loss of former relationship: there must be no shrinkage, no condensation, no expansion; but everything should be precisely as it was when manifesting vital activities, except this change into a compound that will remain undissolved and persist through subsequent necessary manipulations. This insolubility is supposed to be due in some cases to a sort of clotting process; and if the coagulating property be stronger in absolute alcohol than its dehydrating power and less in alcohol of lower percentage, this fact explains why more shrinking is caused by 96 per cent. than by absolute alcohol, and why the shrinking increases with the lowering of the alcoholic strength. Other fixing agents, such as osmic acid, chromic acid, potassium bichromate, and corrosive sublimate solutions, seem to form a chemical union with the cell contents and so produce an extremely durable insolubility. Others, such as picric acid and nitric



acid, harden well, but form such unstable compounds that the fixation is easily removed by washing in water and must be preserved by placing the specimens in alcohol. It is evident that any solvent action by the reagent—*e.g.*, the action of alcohol on fat and that of acetic acid on protoplasm—lessens their practical value.

**OPTICAL DIFFERENTIATION.**—Some agents in producing insolubility effect another change which is equally valuable and which is known as optical differentiation. The various cell structures respond differently to the fixative. Their indices of refraction are altered; some are raised, some lowered, and marked contrasts in refractive properties are developed throughout the cell. In this way structures become visible that were before unseen. Bichromate of potassium stiffens very equably, with neither shrinkage nor expansion, but has no power of optical differentiation; while osmic acid possesses this in a high degree. Since observation with the microscope is directly dependent upon differences in refraction, it is evident that this is a most valuable property of a fixative.

**PENETRATION.**—The ability to reach all points of the tissue at the same time is another important characteristic of a fixing agent and one clearly connected with securing optical differentiation. Osmic acid has but little penetration. If pieces placed in its solutions are too thick or remain therein too long, the superficial layers become over-exposed, the indices of refraction are all equally raised, and differentiation disappears. This is true not only of cells in mass, but also of intracellular structures. Prompt and uniform action, the sharp fixation of tissues at the precise moment, insures good optical differentiation; slow, unequal action results in loss of definition.

**FIXING FLUIDS.**—All acids apparently possess fixing properties, and every fixing fluid should be acid, with possibly the exception of alcohol. Of the organic acids acetic and formic are those most used; of the inorganic, nitric, sulphuric, picric, hydrochloric, osmic, and chromic.

**Acetic Acid.**—By this term is always meant glacial acetic acid, which has very great penetrating power and aids in optical differentiation. It causes swelling and solution of protoplasm, and hence is not used alone, but with fixatives such as osmic acid to aid in penetration and prevent excessive blackening, with alcohol and corrosive sublimate to prevent shrinkage, and with chrome salts to aid in optical differentiation. It is usually added to these various in



strengths varying from 0.5 to 5 per cent. All liquids containing a large percentage of acetic acid should be allowed to act **only** for a short time. Acetic acid should not be used for connective tissue.

*Alcohol* (95 per cent. or absolute; 2-24 hours; 5 mm. thick).—Alcohol has certain important advantages. It can be readily procured, does not have to be made up, tissues are hardened as well as fixed by it, and, since it represents one of the last stages preparatory to embedding, its use saves much time and trouble, and the material for a general diagnosis is easily and promptly prepared, which is often a great convenience. It penetrates well, preserves the specific staining properties of micro-organisms and various important chemical reactions, permits the use of most stains and is demanded by others,—*e.g.*, Nissl's, Lenhossék's, Weigert's, Ribbert's phosphomolybdic hæmatoxylin, Unna's orcein, etc. It is especially good for glands, skin, and blood-vessels, mastcells, plasma cells, fibrin, and hyperæmic conditions, since it preserves the color-reactions of hæmoglobin. On the other hand, it sometimes causes shrinkage and exerts a bad solvent action, so that the cells come out lean and empty, with foamy, vacuolated protoplasm and with distortion or loss of original structure.

Tissues should not remain too long in absolute alcohol, as they sometimes stain very poorly after as short a time as twenty-four hours. Alcohol is not a good fixative for van Gieson's stain. Alcohol of lower percentage than 95 causes excessive shrinkage.

The shrinkage of alcohol is corrected by the use of acetic acid.

Carnoy's fluids (for nuclear structures) :

- |                             |          |
|-----------------------------|----------|
| 1. Glacial acetic acid..... | 1 part.  |
| Absolute alcohol.....       | 3 parts. |
| 2. Glacial acetic acid..... | 1 part.  |
| Absolute alcohol.....       | 6 parts. |
| Chloroform .....            | 3 parts. |

Leave pieces in for from fifteen to thirty minutes; wash out in alcohol. Avoid aqueous liquids.

(For acetic alcohol with sublimate see "Gilson's solution" and "Ohlmacher's solution" under *Corrosive Sublimate*.)

After the use of alcohol as a fixing agent, tissues must either be embedded in celloidin or paraffin as soon as hardened or left in cedar oil, or put through 95 per cent. alcohol and finally preserved in 80 per cent.

*Chromic Acid*.—Chromic acid is a powerful and rapid coagulating

agent, but, on account of its lack of penetration and tendency to cause shrinkage and make tissues brittle, it is seldom used alone. Its defects are remedied by adding acetic, formic, osmic, or nitric acid to its solutions. All tissues fixed by chromic acid solutions are to be washed in running water and hardened in graded alcohols in the dark.

Chromo-acetic acid (Rabl) :

Acetic acid, 0.1 per cent. in water.....	1 part.
Chromic acid, from 0.2 to 0.25 per cent.....	1 part.

Chromo-formic acid (Rabl) :

Chromic acid, 0.33 per cent.....	200 cc.
Formic acid, concentrated.....	from 4 to 5 drops.

Use at once, fix for from twelve to twenty-four hours.

Chromo-nitric acid (Perenyi) (4-5 hours) :

Nitric acid, 10 per cent.....	4 parts.
Alcohol .....	3 parts.
Chromic acid, 0.5 per cent.....	3 parts.

Transfer directly to 70 per cent. alcohol for twenty-four hours, to 95 per cent. for some days, and to absolute alcohol from four to five days.

Chromo-osmic acid has been superseded by

Chromo-aceto-osmic acid (Flemming) :

Chromic acid, 1 per cent.....	45 cc.
Osmic acid, 2 per cent.....	12 cc.
Glacial acetic acid.....	3 cc.

Objects may stay in this solution for hours or even several days. The pieces should be perfectly fresh and not thicker than 4 mm.

It should be made up shortly before using. When all the conditions are fulfilled, it is unequalled as a fixative and in producing optical differentiation. The most delicate structural details are brilliantly shown. Especially used for mitotic figures.

*Bichromate of Potassium*.—The simple aqueous solution is used in gradually increasing strengths from 2 to 5 per cent. for hardening purposes, for which it is excellent, but, on account of its lack of penetration and tendency to cause the chromatin to swell, it is not suitable for a nuclear fixing agent without being reinforced. The addition of glacial acetic acid gives a fluid which acts nearly as well as Zenker's and is much more convenient to use. The excess of bichromate is to be well washed out in running water and the tissues hardened in alcohols in the dark.

## Acetic bichromate (Tellyesniczky) (1-2 days) :

Bichromate of potassium.....	3 grammes.
Glacial acetic acid.....	5 cc.
Water .....	100 cc.

Begin hardening with 15 per cent. alcohol.

## Osmic, bichromate, and platinum chlorid (2 hours) (Dr. Lindsay Johnson) :

Potassium bichromate, 2.5 per cent.....	70 parts.
Osmic acid, 2 per cent.....	10 parts.
Platinic chlorid, 1 per cent.....	15 parts.
Acetic or formic acid (just before using).....	5 parts.

A fine fixative for delicate objects, such as a retina. Leave objects in for two hours. Wash in running water. Harden in alcohol.

The slow, mixed, and rapid methods of Golgi stain the cells with their prolongations, the nerve-fibres with their terminal ramifications, and the neuroglia cells.

Golgi's slow method: Harden pieces of tissue in a 2 per cent. solution of bichromate of potassium from two to six weeks. Keep in the dark and change often. Transfer to a 0.75 per cent. aqueous solution of silver nitrate.

Golgi's mixed method: Harden small pieces of tissue for from three to five days, or longer, in a 2 per cent. solution of potassium bichromate at 25° C. in the dark. Place in the following solution for from three to eight days.

Osmic acid, 1 per cent.....	2 parts.
Bichromate of potassium.....	8 parts.

Then into a 0.75 per cent. silver nitrate solution.

Golgi's quick method: Tissues should be absolutely fresh, and the pieces not more than three millimetres thick.

Osmic acid, 1 per cent.....	1 part.
Bichromate of potassium, 3.5 per cent.....	4 parts.

Leave pieces of neuroglia in the solution for two or three days, nerve-cells from three to five days, nerve-fibres and collaterals from five to seven days. Then place in 0.75 per cent. silver nitrate solution.

## Müller's fluid (6-8 weeks) :

Bichromate of potassium.....	2.5 grammes.
Sulphate of sodium.....	1. gramme.
Water .....	100. cc.



This fluid, once so universally used, is now largely replaced by better fixatives. It has all the faults of the plain bichromate solution and the same need of being reinforced. (For acetic acid and sublimate additions see "Zenker's fluid" under *Corrosive Sublimate*; for formalin see "Orth's fluid" under *Formalin*.) It hardens evenly without shrinkage and gives very good consistency to tissues, but it is in no way a nuclear fixative. As a hardening agent for nervous tissue it has been almost entirely replaced by formalin.

Pieces of tissue not larger than two-centimetres are hardened in from six to eight weeks. Change daily for seven days, then once a week. Wash in running water twenty-four hours. Nervous tissue is placed directly in alcohol.

*Erlick's Solution.*—

Potassium bichromate.....	2.5 grammes.
Copper sulphate .....	0.5 to 1. gramme.
Water .....	100. cc.

This is an extremely good agent for hardening voluminous objects. Its action is much more rapid than that of Müller's fluid. For microscopic work, however, it gives precipitates likely to be misleading and difficult to remove. It is used as a fixative for Freud's gold stain for nerve-fibres.

*Chlorid of Iron (Mallory) (3-5 days).*—For peripheral nerve-fibres.

Chlorid of iron.....	1 part.
Distilled water.....	4 parts.

Wash out thoroughly in water. Transfer to a saturated solution of dinitroresorcin in 75 per cent. alcohol for several weeks. Wash, dehydrate, etc.

This stain may be used after Flemming or Müller.

*Corrosive Sublimate (Bichlorid of Mercury).*—This is a very active penetrating and hardening agent, and since tissues are sufficiently affected by it in from three minutes to two hours and are then placed directly into alcohol, the process is a quick and convenient one. Carmin and van Gieson stains are particularly brilliant after it. The Heidenhain-Biondi triple stain requires its use. It is an especially good fixative for the alimentary tract; for cedematous tissues and albuminous degenerations, since it coagulates nearly as well as boiling water; it is used for connective-tissue fibrillæ with M&L.

lory's anilin-blue stain. Its disadvantages are that it causes shrinkage and the formation of precipitates which must be removed. If tissues are too long exposed to its action they become brittle, and if kept too long in alcohol they are very difficult to cut. Unless corrected by the addition of some other agent, poor optical differentiation is obtained, so that corrosive sublimate should be used only for general and not for cytological work. Pieces of tissue should not be larger than five millimetres, and must be removed as soon as they become thoroughly opaque, otherwise they will be too brittle. All solutions containing this salt act much better when freshly made, as they deteriorate by standing.

Sodium chlorid and bichlorid of mercury (Heidenhain's solution): A saturated solution of bichlorid of mercury in 0.5 per cent. solution of sodium chlorid.

Acetic sublimate: A saturated solution of corrosive sublimate in 5 per cent. glacial acetic acid.

Gilson's solution:

Absolute alcohol.....	1 part.
Glacial acetic acid.....	1 part.
Chloroform .....	1 part.
Sublimate to saturation.	

This liquid is one of the most penetrating and rapidly acting of any, if not *the* most. Wash out with alcohol containing tincture of iodin.

Ohlmacher's solution (15-30 minutes):

Absolute alcohol.....	80 parts.
Chloroform .....	15 parts.
Glacial acetic acid.....	5 parts.
Sublimate to saturation (about 20 per cent.).	

A cerebral hemisphere sectioned by Meynert's method is hardened in from eighteen to twenty-four hours.

Zenker's fluid:

Corrosive sublimate.....	5 grammes.
Glacial acetic acid.....	5 cc.
Müller's fluid.....	95 cc.

Add the sublimate and acetic acid just before using. Leave tissues in from twenty-four to forty-eight hours.

This fluid is comparable to that of Flemming in perfect fixation. It has better penetration, over-fixation is not so likely to occur, it gives better staining results, and is much cheaper. It is altogether most

satisfactory. Eosin stains are especially brilliant after its use. Its one disadvantage is that the sublimate must be removed by placing sections in 70 per cent. alcohol containing enough tincture of iodine to give it the color of a dark sherry wine; but this is true of all sublimate solutions.

Bensley's solution ( $\frac{1}{2}$ -2 hours):

Potassium bichromate, 1 to 2 per cent. solution in water...	1 part.
Corrosive sublimate, saturated solution in alcohol.....	1 part.

Mix the two solutions just before use. Leave tissues in from one-half hour to two hours. Wash well in water.

This solution is especially useful for the gastro-intestinal tract.

As sublamin,<sup>1</sup> the ethyldiaminsulphate of mercury, forms no precipitates, tissues are easily stained after its use, and preserve to a certain extent their natural color.

*Formalin.*—This agent acts very rapidly; it causes little shrinkage. Cytoplasm and nuclei are well preserved. Mitotic figures are fixed. Hæmoglobin and micro-organisms retain their specific staining reactions. Fat is not dissolved; mucin is not precipitated, but remains transparent. Formalin is an especially valuable fixative for nervous tissues: an entire brain may be hardened in a 10 per cent. solution in from a week to ten days. It gives great toughness and elasticity to tissues, and is required for many methods of staining nerve-fibres. Pieces of nerve tissue ten millimetres thick may first be fixed in formalin and then subjected to the action of any mordant desired.<sup>2</sup>

It is used in a standard solution of ten cubic centimetres of formalin to ninety cubic centimetres of distilled water. Change after three hours. Tissues are fixed in from one to two days, but may remain in the fluid indefinitely if the percentage of formalin is maintained.

Orth's fluid (1-2 days):

Potassium bichromate.....	2.5 parts.
Sodium sulphate.....	1. part.
Water .....	100. cc.
Formalin .....	10. cc.

Add the formalin just before using.

<sup>1</sup> KLINGMÜLLER AND VEIEL, *Centralb. f. allg. Path.*, 1903, vol. xiv, no. 20, p. 842.

<sup>2</sup> For the effect of formalin on tissues see *Ir. Am. Med. Assoc.*, March 12, 1904, p. 734 and September 3, 1904, p. 685.



This is Müller's fluid with 10 per cent. formalin. It is one of the best general fixatives in use.

*Nitric Acid* (3 per cent.; 6 hours; 70 per cent. alcohol).—It gives toughness to tissues and is especially suitable for organs rich in connective tissue. Bichromate of potassium may be used after fixation in nitric acid.

*Osmic Acid*.—This is one of the finest fixatives known, especially for cytoplasm. It has great power of rendering cell constituents insoluble and of developing optical differentiation, thus bringing to view structures previously unknown. As it has very little penetration, superficial cells may be overfixed and homogeneous. Carmin stains badly after its use, but hæmatoxylin is not affected. It is seldom used alone except for fixation by vapors. Very delicate objects are pinned out on the well-fitting cork of a wide-mouthed bottle and exposed to the vapors of a small quantity of a 1 per cent. solution poured into the bottle. A retina needs an exposure of some hours and is more equally fixed than when placed in the solution. Osmic acid solutions do not keep well and must be carefully protected from dust. Lee recommends a 2 per cent. solution in 1 per cent. chromic acid. This serves for vapor fixation and Flemming's solution. It may also be kept as a 1 per cent. solution in distilled water. (For Flemming's solution see "Chromo-aceto-osmic acid.") In making osmic acid solutions the capsule containing this acid is broken within the bottle containing the solution. Tellyesniczky<sup>1</sup> suggests as the best substitute for osmic acid the following:

Potassium bichromate .....	3 grammes.
Acidi aceti .....	5 cc.
Aquæ .....	100 grammes.

Platinico-acetico-osmic-acid solution (Hermann's solution; 1-8 days): This celebrated reagent is Flemming's solution with platinic chlorid instead of chromic acid.

Platinic chlorid, 1 per cent.....	15 parts.
Glacial acetic acid.....	1 part.
Osmic acid, 2 per cent.....	2 to 4 parts.

Its action is comparable to that of Flemming's solution. The most delicate structures are faithfully preserved and well shown.

---

<sup>1</sup> *Arch. f. mikrosk. Anat.*, 1898, vol. lii, p. 202.

## Pianese's solution (36 hours):

Chlorid of platinum and sodium, 1 per cent. aqueous solution .....	15 cc.
Chromic acid, 0.25 per cent. aqueous solution.....	5 cc.
Osmic acid, 2 per cent. aqueous solution.....	5 cc.
Formic acid, C. P.....	1 drop.

For karyokinesis and the so-called cancer bodies. Pieces of tissue must not be more than two millimetres thick. It gives very interesting results histologically.

*Picric Acid* (2-24 hours).—Picric acid is an extremely penetrating and delicate fixative. It hardens very slightly, and the insolubility caused by its action may be easily removed by washing in water; hence its preparations should always be placed in alcohol. It is used as a saturated aqueous solution and in large quantity,—about one hundred times the bulk of the object. It is an excellent fixative for delicate serous membranes, which may be floated in it without retraction or distortion. The omentum and peritoneum are well fixed in it.

Picro-acetic acid: A saturated solution of picric acid in one per cent. acetic acid; a very good fixative.

Picro-sulphuric acid (Kleinenberg): Add 1 cc. of concentrated sulphuric acid to 100 cc. of a saturated aqueous picric acid solution. Let stand for nearly four hours; filter; add double its volume of distilled water. This is an excellent fixative for delicate embryos.

Picro-nitric and picro-hydrochloric acid solutions are also used, but their action is essentially the same as that of picro-sulphuric.

The advantages of picric acid solutions are that they give a very delicate fixative with excellent cutting qualities, and delicate membranes are not thickened excessively as with stronger reagents.

**HARDENING.**—To give to tissues a proper cutting consistency they are gradually hardened by being passed through a series of graded alcohols. For general diagnosis tissues may go from water into 70 per cent. alcohol, then 95 per cent., and finally absolute alcohol, usually remaining twenty-four hours in each grade. Corrosive sublimate and Golgi tissues are to be placed for only a few hours in 95 per cent. and absolute alcohols, without passing through the lower grades. For finer work begin with 30 per cent. or even 15 per cent. alcohol, then use 50, 70, 80, 95, and absolute. When the tissues are passed from a lower to a higher grade of alcohol, surplus moisture should be removed with blotting-paper to avoid lowering the percentage of the next grade.

**PRESERVATION.**—After being fixed and hardened, tissues are usually preserved in 80 per cent. alcohol. Those fixed by formalin may remain in a 10 per cent. solution thereof. Golgi preparations keep indefinitely in the silver nitrate solution. Corrosive sublimate tissues will not cut well if kept too long in any kind of alcohol; they had better be kept in cedar oil. A sugar formation takes place in liver preserved in alcohol, and certain peculiar changes may arise in the nervous tissue, especially if acted upon by sunlight.

*Pathologic Conditions suggesting Certain Fixatives.*—Acute infectious processes: Alcohol.

Acute inflammatory exudates: The fibrin, leucocytes, and red blood-corpuscles of hemorrhagic conditions are preserved especially well in Zenker's fluid.

Albuminous degenerations: Corrosive sublimate, Zenker, or boiling water.

*Amœbæ coli:* Stain especially well with Mallory's chlorid of iron hæmatoxylin; any fixative may be used except perhaps formalin. *Amœbæ coli* may be studied either in the fæces or in the tissues. Collect the fæces in a perfectly clean dry vessel, warmed in cold weather, and keep them at the temperature of the room. Add a drop of a weak solution of toluidin blue to a particle of the fæces, make a cover-slip preparation, and preserve in Farrant's medium. For the tissues fix in Heidenhain's or Bensley's solution, stain with iron hæmatoxylin or with a weak solution of toluidin blue. If a contrast stain is desired, stain first with eosin or benzo-purpurin, then for fifteen or twenty minutes with a weak solution of toluidin blue; differentiate with alcohol.

Amyloid degenerations: Corrosive sublimate, Zenker, alcohol.

Blood: Make thin films; stain with a 0.5 per cent. solution of the eosinate of methylene blue in absolute alcohol.

Bone: For infectious processes, alcohol; for histological purposes, Zenker, Orth. Bone must always be fixed before decalcifying.

Bone marrow: Make smears on cover-slips. Fix pieces of bone marrow in Zenker or formalin.

Cartilage: Alcohol, Zenker, Orth.

Central nervous system: A whole brain may be hardened in about three thousand cubic centimetres of Müller's fluid. Change every day for a week, then every week for four weeks, and every two weeks thereafter; it takes about three months to complete the hardening. Keep in a refrigerator if the weather be very warm. Erlicki's fluid



hardens better and its action is more rapid, hardening being accomplished in about four weeks.

In a 10 per cent. solution of formalin a whole brain may be hardened in from ten days to two weeks. Change the solution every day for three days, then every third day. Cerebral hemispheres may be sectioned by Meynert's method and hardened in twenty-four hours in Ohlmacher's solution. These methods are not recommended for fine work. Pieces not larger than one centimetre may be hardened in formalin and then subjected to any bichromate or osmic acid mordant, including Golgi's methods.

Ganglion cells: For Nissl's method fix in 96 per cent. alcohol. For Lenhossék fix in 90 per cent. alcohol (or 10 per cent. formalin) and follow with 96 per cent. alcohol. For Golgi methods use Golgi fixatives.

Myelin sheaths: For Weigert fix with 5 per cent. bichromate until "ripe,"—that is, until color contrasts between white and gray matter are well developed. For Marchi use Müller's fluid. Use formalin for Busch-Mallory, Weigert, Weigert-Pal, and Heller. For Exner use 1 per cent. osmic acid; change second day; leave pieces in for five or six days.

Neuroglia fibres: These are not well preserved by chromates. For Weigert methods fix in formalin. For Mallory fix in ten per cent. formalin in a saturated aqueous solution of picric acid.

Medulla, pons, and basal ganglia: They may be removed together *en masse* and hardened entire in formalin for from one to two weeks, then cut into parallel slices not over one centimetre thick, and mordanted by Weigert's quick method or Mallory's or in any way desired. Golgi stains are not very applicable to the medulla.

Axis-cylinders and their terminal processes: For Freud's or Stroebe's gold stain fix in Erlicki or Müller. For Gerlach's method harden in 0.5 per cent. solution of bichromate of ammonium for from one to three weeks. (For Golgi see "Golgi methods" under *Bichromate of Potassium*.)

Degenerated nerve-fibres: Harden in Müller or Erlicki for Marchi or Algeri methods, or harden in 10 per cent. formalin followed by Müller and Erlicki.

Peripheral nerve-fibres: Fix in chlorid of iron.

Retina: The retina may be fixed in a 10 per cent. solution of formalin; in Zenker's, Orth's, or Lindsay Johnson's solution, as given

under *Bichromate of Potassium*; in equal parts of glacial acetic acid and osmic acid (2 per cent.); in equal parts of chromic acid and platinic chlorid (each 1.4 per cent.); or it may be pinned out on a cork and exposed to the vapor of a 1 per cent. solution of osmic acid.

Colloid material: Formalin or Orth.

Connective tissue: For Ribbert's phosphomolybdic hæmatoxylin stain for fibrillæ fix in alcohol. For Mallory's anilin-blue stain fix in corrosive sublimate or Zenker.

Elastic fibres: For Unna's orcein method fix in alcohol. For Weigert fix in alcohol or formalin.

Fatty changes: Flemming, Orth, Müller, Erlicki, or formalin.

Fibrin: For eosin hæmatoxylin, methylene blue, and Mallory's anilin-blue stain fix in Zenker or corrosive sublimate. For infectious processes and Weigert's method fix in absolute alcohol.

Glands: Fix in absolute alcohol.

Granulation tissue: Fix in Zenker, Flemming, or Pianese for attendant degenerations.

Hyaline degenerations: Zenker, corrosive sublimate, Orth.

Liver: For pernicious anæmia and amyloid degenerations fix in alcohol. For bile capillaries use Golgi method.

Mastcells: For Ehrlich's or Unna's methods fix in alcohol.

Mucoid material: For Mallory's anilin-blue stain fix in Zenker or corrosive sublimate. For other stains use Orth or formalin.

Myxomas: Zenker or corrosive sublimate.

Œdematous conditions: Throw small pieces of tissue into boiling water for a minute or two, or fix in corrosive sublimate.

Ovaries: For follicular degenerations use Flemming or Hermann if tissues are fresh, if not use Zenker, Orth, Carnoy, or Ohlmacher.

Pancreas: For Altmann's granules fix in equal parts of a 5 per cent. solution of bichromate of potassium and a 2 per cent. solution of osmic acid.

Plasma cells: Zenker is especially favorable for showing eosinophiles.

Pus or purulent conditions: Orth, Zenker, or corrosive sublimate.

Skin is best fixed in alcohol.

Spleen: For Heidenhain Biondi triple stain fix in corrosive sublimate. For eosinophiles or Ehrlich's triacid use Zenker or alcohol.

Suprarenal: If fresh fix in Flemming or Hermann; if not, in Ohlmacher, Zenker, or Orth.

Thyroid: For colloid degeneration fix in Orth or 10 per cent. formalin.

*Fixatives.*—The following list gives the fixatives used for the various stains.

Alum hæmatoxylin: Stains very slowly after chromic solutions.

Anilin blue (Mallory): Succeeds best after Zenker or corrosive sublimate. It may be used after formalin.

Biondi Heidenhain (see "Heidenhain Biondi").

Eosin and methylene blue: Best after Zenker.

Freud's gold stain: For axis-cylinders and nerve terminals; used after Müller or Erlicki.

Gold stains: Freud's, Stroebe's, after Müller or Erlicki; Gerlach after 0.5 per cent. bichromate of ammonium for from one to three weeks.

Golgi chrome silver preparation: After Golgi fixing solutions.

Heidenhain Biondi triple stain: Only after corrosive sublimate.

Lenhossék: For ganglion cells 90 per cent. alcohol or 10 per cent. formalin, both followed by 96 per cent. alcohol.

Nissl: For ganglion cells 96 per cent. alcohol.

Orcein (see "Unna's orcein stain").

Phosphomolybdic acid hæmatoxylin: Best after alcohol.

Phosphotungstic acid hæmatoxylin: After 10 per cent. formalin.

Thionin (Lenhossék's ganglion-cell stain): 90 per cent. alcohol followed by 96 per cent. or formalin 10 per cent.

Triple staining: Heidenhain Biondi only after corrosive sublimate.

Unna's alkaline methylene blue: Alcohol.

Unna's orcein stain: For elastic fibres, alcohol.

Weigert's stain: For fibrin and elastic fibres, absolute alcohol.

*Macroscopic Specimens.*—If a microscopic examination of the organ to be preserved is desirable, portions of tissue therefor should be removed before anything is done towards preparing it as a gross specimen.

If for any reason it be desirable to keep the specimen for a short time, it should be kept moist by being wrapped in cloths wet with 10 per cent. formalin solution. Parenchymatous organs of slaughtered animals will keep for a week packed in this way and, when sectioned, the tissues appear fresh. The organs of deceased animals do not keep as well. If the specimen is to serve for a bacteriologic investigation



and for inoculations, it should not be wrapped in any disinfecting agent, but simply packed in parchment-paper or rubber cloth.

By a percentage solution of formalin is meant such a dilution of the commercial 40 per cent. (which is sold as formalin) as will reduce it to the desired strength. For instance, ten cubic centimetres of commercial formalin added to ninety cubic centimetres of water produce a 10 per cent. solution of formalin or a 4 per cent. solution of formaldehyde. The percentage of formalin must be maintained, as it is quickly exhausted; when there is no odor of formalin, the fluid should be renewed.

It is not always necessary to save the entire organ to be examined, but enough should be preserved to show its relationship to the lesion.

**GENERAL CONSIDERATIONS.—*Washing.***—If alcohol be used as the preserving solution, blood and other impurities may be removed by a thorough washing with water. In other cases the parts should be carefully sponged with the preservative to be employed.

Cavities should be distended with tow or absorbent cotton. The lungs should be placed in a jar and the jar filled by pouring the fluid through the trachea. Mucous and serous membranes should be protected from the distortion caused by shrinkage by being pinned out on cork or on wood which will impart no color in soaking. A more elegant method is to sew the membranes over the edges of frames made of glass rods. The secreting surfaces of these membranes should always be uppermost.

Compression of any part of the specimen should be avoided by the use of a soft cushion of absorbent cotton placed in the bottom of the jar. Jars made especially for museum preparations are preferable, but if necessary they may be replaced by such as are used by grocers and druggists for candy, etc.

***Preserving Fluids.***—Alcohol is a convenient and efficient agent. It preserves form relationships very well, as in tumors, typhoid ulcers, invagination of the intestine, etc.; but it destroys all contrasts in a pathologic organ, such as a diseased lung or kidney, and makes recognition of the lesion very difficult. It bleaches the tissues and causes much shrinkage, so that natural appearances are not retained. The specimen is to be washed in water, then immersed in 60 per cent. alcohol (which is changed every day until it remains clear), and finally kept in 80 per cent. alcohol. To preserve the natural appearance of tissues, formalin followed by alcohol is used, and the specimen is

finally placed in glycerin solution containing some salt of acetic acid, usually potassium. Formalin converts the hæmoglobin into methæmoglobin and a brown color is developed; alcohol changes the methæmoglobin into a red pigment, so that the flesh-color is restored. The tissues are so thoroughly hardened that they may be kept in the glycerin solution without being thereby softened. The principles involved are simple, but their application requires experience and ingenuity. All tissues do not respond equally to the treatment, and to retain some color peculiar to a certain pathologic condition—such as prevails in icterus, for example—requires careful management. There are various formulæ and different methods of applying them, but the two following are perhaps as simple and useful as any. It must always be remembered that if the tissues are placed in too strong formalin, or remain too long even in a weak solution, the alcohol will fail to transform the brown or gray pigment back into red.

1. Place the fresh organ or a segment as large as the hand for from twenty-four to forty-eight hours in one of the following solutions.

Kaiserling fluid:

Formalin .....	200 cc.
Water .....	1000 cc.
Potassium nitrate.....	15 grammes.
Potassium acetate.....	30 grammes.

Melnikow-Raswedenkow:

Formalin .....	10. parts.
Sodium acetate.....	3. parts.
Potassium chlorate.....	0.5 part.
Distilled water.....	100. parts.

It is well to wrap the specimen in wadding and pour the fluid over it. The wadding protects the organ from distortion due to compression. If the organs are very thick, incise them or inject the blood-vessels, ureters, etc., with the fluid. This should be done very gently, in order not to wash out the blood. As formalin is injurious to the respiratory tract and the skin, it is well when using it to wear rubber gloves and to keep the jars covered.

2. After two days place the specimen in 60 per cent. alcohol, first removing the wadding. Two or three days later change to 80 per cent. alcohol, then to 90 or 93 per cent.

## 3. The specimen is finally placed in the preserving fluid :

Glycerin .....	400 grammes.
Potassium acetate.....	200 grammes.
Water .....	2000 grammes.

The solutions may be used several times, but a fresh preserving fluid is better, and it is even advisable to change it occasionally.

Pick adds at once to the formalin solution 5 per cent. of Carlsbad salts, which prevents the formation of acid hæmatin, while Marpmann uses fluorsodium both in the formalin solution and in the glycerin. The use of ten parts of an 0.8 per cent. salt solution with one part of the 40 volume strength formalin is also recommended.

Another method of preserving the natural color of specimens is as follows: <sup>1</sup> One-half of the capacity of a metal box is filled with a concentrated solution of ammonium sulphate, an excess of the crystals being left at the bottom of the tank. Above the crystals is arranged a grating upon which the specimens to be acted upon are placed. At the bottom of the box is a small opening through which carbon dioxid or ordinary illuminating gas is constantly passed, thus permitting it to bubble up through the fluid in the box. Another tube at the top of the box is fitted with a burner so as to burn off the escaping illuminating gas. The specimens which remain in the solution under the action of the carbonic-acid gas and ammonium sulphate for from forty-eight to seventy-two hours retain their color for a long while, if preserved in this solution.

INJECTED SPECIMENS.—Most beautiful and permanent specimens may be made by injecting various colored materials, such preparations giving especial opportunity for the study of the arterial and venous circulation. Thus, in the case of the liver, if the cystic duct, portal vein, hepatic artery, and hepatic veins be injected with four different colored solutions, the distribution of the various vessels may be shown to perfection. The microscopic study of these cases may be made later on. The writer has a fine specimen in his cabinet of the *Trichina spiralis* in the tongue of a cat in which the arterial circulation has been injected with carmin. Entwining capillaries surrounding the capsule are well brought out. If a warm injecting fluid be desired, that of Robin may be recommended. It consists of gelatin one part and seven to ten parts of water, heated on a water-bath, to which two

<sup>1</sup> CLAUDIUS, *Virchow's Arch.*, 1903, vol. clxxiv, no. 1, p. 193.



per cent. of chloral hydrate is added to prevent the formation of mould. Any dye may be used to color this solution.

Richardson blue may be prepared as follows:

1. Sulphate of iron.....	0.62 part.
Distilled water.....	30.00 parts.
2. Red potassium ferrocyanid.....	2.00 parts.
Distilled water.....	30.00 parts.

Slowly mix and shake, then when an opalescent blue, add

3. Distilled water.....	60.00 parts.
Glycerin.....	30.00 parts.
Alcohol.....	30.00 parts.

Beale's Prussian blue may be used, and is prepared as follows:

Glycerin.....	32.0 parts.
Alcohol (50 per cent.).....	32.0 parts.
Potassium ferrocyanid.....	0.75 part.
Tinct. of perchlorid of iron.....	4.0 parts.
Distilled water.....	128.0 parts.

To decalcify bone tissue the following formula may be employed:

Sodium chlorid.....	100 parts.
Distilled water.....	100 parts.
Hydrochloric acid.....	4 parts.

Strong nitric acid, two parts; chromic acid, one part; and water, two hundred parts, may also be used.

Littlejohn<sup>1</sup> recommends that fresh specimens or those preserved by any well-known method be kept in glass jars made air-tight by sealing their covers with gold size and putty. The one objection to this method is the vapor which collects in the jars. To avoid this the preparations are soaked for several weeks in glycerin and water and afterwards placed on wool to which some formalin glycerin is added. Perfectly washed stomachs from cases of poisoning, such as carbolic acid and the corrosive acids, require no preservative whatever, and when thus prepared retain their natural coloring for years.

The August 13, 1904, number of the *Journal of the American Medical Association* contains two excellent articles on the permanent preservation of specimens, one being by Coplin and the other by Herring.

<sup>1</sup> *Journal of Pathology and Bacteriology*, September, 1902, p. 369.

## CHAPTER XXIII

### BACTERIOLOGIC INVESTIGATIONS

ALTHOUGH it is well known that a bacteriologic investigation is often a most important factor in the ultimate value of a post-mortem examination, such an investigation is frequently neglected because of the lack of facilities or of knowledge of the technic.<sup>1</sup> This ought not so to be. In the first place, the cost of equipment, as in post-mortem sets, is very largely determined by the *conveniences*, rather than by the *necessities*. The outfit mentioned on page 35 can be kept always in readiness, while the culture-tubes may be obtained quickly and at reasonable rates from the larger pharmacal manufacturing companies and their agencies. In the second place, the technic is not so complicated as to require more skill, except in the finer manipulations and diagnoses, than should be expected from an educated physician. As time goes on, the general practitioner who is not within easy reach of a pathologic laboratory or of a board of health will be more and more expected to be sufficiently equipped with apparatus and adequately trained to make cultures and even inoculations for diagnostic purposes. Of course, it is impossible under such circumstances to do the work of well-endowed laboratories and skilled bacteriologists, but the material may at least be studied until the time arrives for placing it in the hands of those devoting their especial attention to the technic of bacteriologic investigations.

**COLLECTION OF MATERIAL FOR MICROSCOPIC OBSERVATIONS AND FOR CULTURE PURPOSES.**—The important factor in the technic of a bacteriologic examination is that all instruments shall be scrupulously clean and absolutely sterile, and all sources of contamination carefully guarded against in every possible manner. The fluid contents and accumulations in abscess and serous cavities, especially meningeal, pericardial, peritoneal, and pleural, the blood, endocardial vegetations, ulcerated areas, and the cut surfaces of solid organs may present foci of bacterial invasion which are at once examined by "smear prepara-

---

<sup>1</sup> SIMMONDS (*Virchow's Arch.*, vol. clxxv, no. 3, p. 418) believes, after making routine bacteriologic examinations in 1200 cases, that this procedure may sometimes give the only definite knowledge concerning the cause of death.

tions," and later on by cultures and by animal inoculations, should such be deemed necessary. It is important to obtain material as fresh as possible and in sufficient amount to permit of a thorough examination. Canon<sup>1</sup> states that there is danger of the migration of organisms within the cadaver during the first thirty-six hours, even if it be properly cared for. Should an early examination be wanted, one of the veins of an arm may be exposed shortly after death and the blood thus obtained. The method so frequently employed of taking up a small quantity of the blood with the platinum loop often gives negative results, especially for culture purposes. Bulbs blown in ordinary glass tubing furnish one of the most satisfactory means for the securing of fluids during an autopsy. They may be purchased in supply houses or prepared as follows: A piece of thick tubing is chosen measuring about nine inches in length. The lower portion of the tube is drawn to a point and sealed. About three inches from this end, the glass is heated to a white heat, the tube being turned all the time and not allowed to bend. The open end is now blown into until a bulb about one inch in diameter is produced in the heated portion of the glass. The upper part is now closed by heat, or the opening preferably is filled with a small plug of cotton, and then the whole is sterilized by dry heat. When the bulb is wanted for use, the capillary end is broken off, and, after aspiration of the fluid, immediately sealed by drawing it to a point again. The end previously filled with cotton is also melted until it closes. In these hermetically sealed bulbs the material may be kept securely until the autopsy is completed, and then be taken to a suitable place for such further examination as may be necessary. An ordinary 5 to 10 c.c. pipette sterilized and securely wrapped in cotton may also be used for this purpose. The end of the pipette is placed in the fluid and suction is made through a clean piece of rubber tubing, or the pipette is fitted with a suction cap or bulb. Both ends are then sealed with the flame. By either of these methods sufficient material may be obtained for making differential staining tests and also for the inoculation of cultures or of animals. Solid material may be removed from the interior of an organ by means of a small spear made for this purpose, which has an eye in which some tissue is retained as the spear is withdrawn.

**SMEAR PREPARATIONS.**—Smears which are to be examined during

<sup>1</sup> *Deutsche Zeitschrift f. Chirurgie*, 1901, vol. lxi, nos. 1 and 2, p. 93.



the autopsy may be easily and quickly made, as there is not the same risk of contamination that there is in obtaining fluids and solids to be used for inoculation. They are prepared in the following manner: A number of carefully cleaned and dried cover-slips and slides are placed in readiness.<sup>1</sup> A platinum loop for fluids, or the spear-headed spatula for solids, is then sterilized by heating to a red glow in an alcohol flame or in the upper (hottest) part of a Bunsen burner. If the liquid to be examined is of considerable consistency, like pus, blood, and exudates, a drop of it is placed by the aid of the sterilized loop upon a clean cover-slip. The cover-slip is then dropped upon a slide one-third of its length from one end, and, after the drop has spread, the cover-slip is drawn gently by means of forceps across the remaining two-thirds of the slide. The slide is much easier to manipulate than two cover-slips prepared by drawing one over the other, is not so easily broken, and gives a larger field for future study. Fluids may also be spread zigzag upon the slide or cover-slip with the platinum loop or with a small pipette, the latter being preferable whenever the fluid is very thin, making large amounts necessary. Should the material not be sufficiently fluid to make a satisfactory smear preparation, a little distilled water or physiologic salt solution may be put on the glass before performing the above manipulations. If preferred, a solid organ may be incised with a scalpel sterilized by heat, and the cover-slip or slide applied directly to the freshly cut surface. The material thus collected may then be smeared over the glass with the platinum loop. The "smear" being dried with *very little* heat,—or, better, with none,—now requires only "fixing" on the glass. This is done by the routine method of passing it three times through a flame, with the smeared surface upward to avoid burning the material. If a cover-glass is used, the passage through the flame is made more quickly than when the thicker glass slide is employed. In "fixing," very great care must be used to avoid the application of too high a temperature,—shown by a brownish coloration,—which would seriously distort the bacteria, especially if the film had not been thoroughly dried previously. By the heat applied in this way, the albuminous organic matter is dried or coagulated, and the bacteria and cellular elements are thus caused to

<sup>1</sup> It is well to use new cover-slips which have been cleansed in strong nitric acid, washed in distilled water, and kept in alcohol to which a few drops of ammonia have been added. When wanted for use, they should be wiped dry between the fingers with Chinese tissue paper or with a clean cloth.

adhere so firmly to the glass surface that they will not be washed off by future manipulations. Such preparations may be kept for a considerable length of time before being stained, and can be safely and easily protected by gumming the clean surface to a piece of card-board cut to the size of the ordinary glass slide, on which also may be written all necessary data. The cards may then be packed in a slide-box or in an ordinary pill-box, care being exercised that the films do not come in contact with anything that will be liable to rub or scratch them. Another way to keep two slides apart face downward is to lay short pieces of match-sticks across their ends and bind the slides firmly together with a gum elastic band placed around them lengthwise. The value of a negative finding in a slide from a suspected syphilitic sore may be considerable and is not sufficiently appreciated.

SELECTION OF CULTURE-MEDIA.—A diagnosis made from the study of smear preparations must often be corroborated by cultures, though the previous study of the smear has frequently offered valuable suggestions as to the particular kind of culture-media to employ in the case under observation. For instance, if a diplococcus be found as the prevailing organism, a special medium will be necessary, as the three most common varieties of the diplococcus—viz., the *Pneumococcus*, the *Gonococcus*, and the *Diplococcus intracellularis*—grow poorly and in many cases not at all upon ordinary media. Or the microbe may be one that can be most easily isolated by immediate inoculation into an animal, as in the case of the tubercle bacterium and the *Pneumococcus*. Again, existing conditions of the organs and tissues may point to infection by an anaërobic organism, as the gas bacterium, and in such cases a medium and a method suitable for anaërobic growth—namely, the exclusion of oxygen—must be employed. Wallis<sup>1</sup> drops a little of the melted media upon a cover-glass, places a hair suspected of containing the parasite upon it, puts the cover in a moist chamber, and incubates at the room temperature. The slide is examined in the usual manner.

The following list, prepared chiefly according to the nomenclature adopted in the third German edition of Lehmann and Neumann's *Bakteriologische Diagnostik*, gives the best media for the isolation of the pathogenic micro-organisms most commonly found *post mortem*, those having spores being called bacilli while those which have none are designated bacteria. (Plate V.)

---

<sup>1</sup> *Jr. Amer. Med. Assoc.*, August 20, 1904, p. 531.


- Bacillus anthracis*. All ordinary media, especially agar. White or house mouse. The *Bacillus subtilis* is motile, while this organism is not, though very similar morphologically. (Plate V, no. 10.)
- Bacillus œdematis maligni*. Anaërobic culture methods. Ordinary media with the addition of glucose.
- Bacillus tetani*. Anaërobic culture methods. Ordinary media with the addition of glucose. Spores flagellated.
- Bacterium aërogenes capsulatum*. Anaërobic culture methods. Ordinary media with addition of glucose. (Plate V, no. 11.)
- Bacterium coli commune*. Glucose agar-agar shows gas formation, thus distinguishing it from the typhoid germ. It also contains fewer flagella than the latter. Agglutinates.
- Bacterium dysenteriae* (Shiga). Plain agar-agar. A flagellated organism not stained by Gram's method. Agglutinates. (Plate V, no. 8.)
- Bacterium enteritidis*. The bacterium of hog cholera may produce gastric disturbances in man after eating contaminated meat. It is probably a modified form of the *Bacterium coli*.
- Bacterium influenzae*. Agar-agar smeared with human blood or glycerin agar.
- Bacterium paratyphoid*. Grows on ordinary media. The organism stands between the typhoid and the colon bacterium.
- Bacterium pestis*. All ordinary media.
- Bacterium pneumoniae*. A facultative organism growing on all ordinary media.
- Bacterium pyocyaneum*. Plain agar-agar.
- Bacterium rhinoscleromatis*. Probably the same as the *B. pneumoniae*.
- Bacterium septicæmiæ hæmorrhagicus*. The bacillus of chicken cholera. Should be studied with wine, milk, and glucose media.
- Bacterium typhi murium*. The bacillus of mouse septicæmia is similar to the bacterium of hog cholera and the paratyphoid.
- Bacterium typhosum*. Hiss's gelatin-agar medium. Can grow on all media. Not stained by Gram's method. Flagellated organism. Cultures best obtained from blood, spleen, and urine. Agglutinates. (Plate V, no. 7.)
- Bacterium vulgare* or *protium*. Non-motile; stains with Gram. It is an anaërobic, sugar-decomposing, and agglutinative organism.
- Corynebacterium diphtheria*. Blood-serum bouillon. A guinea-pig may be inoculated to see if organism is virulent, thus distinguishing it from the so-called pseudodiphtheria bacillus. The *Bacillus xerosis* is, in all probability, this organism. (Plate V, no. 6.)
- Corynebacterium mallei*. Optional anaërobic. Guinea-pig inoculated in the testicle. Not stained by Gram's method.
- Corynebacterium xerosis*. See *Corynebacterium diphtheria*.
- Gonococcus* or *Micrococcus gonorrhæa*. Ascites-glycerin agar and hydrocele agar give fairly good results. Gonorrhœal ophthalmia may be produced in rabbits by inoculating the mucosa of the eye. Does not stain by Gram. (Plate V, no. 1.)
- Mycobacterium lepræ*. It is innocuous for animals. It is grown upon glycerin agar-agar with great difficulty. Acid resisting.
- Mycobacterium tuberculosis*. Glycerin agar-agar is the best medium. Subcutaneous inoculation of guinea-pig. Probably agglutinates under proper conditions. von Schrön has recently announced, through his assistant Galbo (*Riforma medica*, 1904, vol. xx, no. 29, p. 800), the discovery of the phthisiogenic micro-organism in the caseous masses of tuberculous lungs. (Plate V, no. 4.)





PLATE V.—BACTERIOLOGIC CHART.

No. 1. *Gonococcus*: smear preparation from urethral pus; stain, methylene blue. No. 2. *Pneumococcus*: smear preparation from sputum; Welch's acetic acid stain. No. 3. *Streptococcus pyogenes*: smear preparation from pus; stain, methylene blue. No. 4. *Mycobacterium tuberculosis*: smear preparation from sputum; stain, Ziehl's method. No. 5. *Vibrio cholerae*: stain, carbol fuchsin. No. 6. *Corynebacterium diphtheriae*: stain, Löffler's method; lower portion of figure shows the polar staining of Neisser's granules. No. 7. *Bacterium typhosum*, showing flagella: stain, van Ermenghem's method. No. 8. *Bacterium dysenteriae*: stain, methylene blue. No. 9. *Achorion Schönleini* (favus fungus), with conidia and mycelia: stain, Bismarck brown. No. 10. *Bacillus anthracis*: smear preparation from spleen of a mouse; stain, gentian violet; to bring out the spores stain with Ziehl's solution. No. 11. *Bacillus aerogenes capsulatus*: smear preparation from spleen; stain, gentian violet. No. 12. Yeast cells with buds and ascospores, starch cell in lower right-hand corner: stain, weak Lugol's (Gram's) solution.





■

1

- Micrococcus* or *Streptococcus intracellularis*. Glycerin agar or agar smeared with blood gives the best results. Löffler's blood serum and potato may be used. It is found in the pus of the brain, cord, and ear, the nasal mucus, the sputum, and the urine of those affected with the disease. A large amount of the exudate should be used, for many of the bacteria are dead.
- Micrococcus melitensis*. Glycerin agar or potato. Grows as a bacterium. Material to be taken from the cervical or inguinal glands and from the lungs. Agglutinates.
- Micrococcus* or *Staphylococcus pyogenes aureus*. Grows luxuriantly upon all media in general use. The *M. citreus* and the *M. albus* are but varieties of this genus.
- Pneumococcus* or *Streptococcus lanceolatus*. Ascites agar and ascites-glycerin agar are the best media. The best method is to inoculate a mouse or rabbit subcutaneously with the rusty sputum or with pus. A mouse dies in from twelve to twenty-four hours, and its blood contains large numbers of the *Diplococcus pneumoniae*, showing the capsule most strikingly; a rabbit dies in from two to five days. (Plate V, no. 2.)
- Sarcina*. Best cultivated in bouillon or hay decoction, displaying fine shades of various colors.
- Spirochæta Obermeieri*. No culture method known. Monkeys show disease after inoculation.
- Streptococcus pyogenes*. This organism is most easily recognized by smear preparations and grows well upon all media in common use. It produces erysipelas as well as pus. (Plate V, no. 3.)
- Streptothrix actinomyces*. Grows well upon agar-agar and blood serum. Does not stain with Gram's solution.
- Vibrio cholerae*. Glycerin agar. All ordinary media, especially gelatin at 22° C. Usually a single flagellum. Agglutinates. (Plate V, no. 5.)

The following diseases are due to specific organisms or protozoa, but the etiologic factor as such, though described, has not yet been accepted as the cause of the affection. Acrodynia, or epidemic erythema; anterior poliomyelitis; beriberi (possibly an arsenical neuritis); cancer; chancroid; chicken-pox; chorea; cow-pox; foot-and-mouth disease; measles (bacillus of Canon and of Czajkowski); miliary fever; mumps; pellagra; rabies; rheumatism; rose rash; scarlet fever; smallpox; syphilis (bacillus of Lustgarten and of van Niessen; baboons, apes, and monkeys show the disease upon inoculation); typhus; whooping-cough (*Bacillus pertussis*, *B. minutissimus sputi*, and *B. tussis convulsivæ*); yellow fever (*Bacillus X* and *Bacillus icteroides*).

**INOCULATING CULTURE-MEDIA.**—Test-tubes containing any of the solid or liquid media may be inoculated at the place where the autopsy is performed when it is not so far from the laboratory as to endanger the growth of the culture by exposure to extremes of temperature.

Sufficient heat is secured, however, by placing the tubes **after inoculation, securely wrapped**, in an inside coat-pocket.

The first step in inoculation of the tube containing the medium is to "flame" the cotton plug in its mouth, for the purpose of **killing any bacteria** that may have fallen upon the cotton-wool. During inoculation the tube is held as nearly in the horizontal position as is **consistent** with safety to its contents, so as to diminish the risk of **contamination** of the medium by the falling into the tube of the bacteria from the air. Thus, should the tube contain a solid medium, such as blood-serum or agar, it may even be inverted before inoculation. The platinum wire, held in the right hand, is now sterilized by heat and cooled, while the cotton plug is removed from the test-tube by a corkscrew motion and held, inner part outward, between the index and middle fingers of the left hand in such a manner that it does not come in contact with any portion of the hand or other extraneous object. With the tip of the platinum wire a small portion of the substance to be inoculated is now placed on the surface of the medium; if this surface is slanting, the fluid is rubbed gently over it, or drawn in a line across it, thus making a "smear" and "stroke" culture, while the needle is thrust deep down into the medium if a "stab" culture is to be made. Theöse is then withdrawn, the cotton plug reinserted, the needle sterilized, and the tube labelled and put in a warm spot until it can be placed in the incubator.

If the culture is to be made from the surface of a solid organ, the method is the same, except that the organ is first seared with a hot knife and next incised with a sharp, sterile knife. Should the same knife be used for both purposes, it is wise, as a precautionary measure, to sterilize the instrument again before plunging it deep into the tissue. In some instances, in which many of the bacteria are dead, much larger amounts of the infectious material must be used for inoculation, and then the entire contents of a bulb or pipette may be introduced into a flask of bouillon or other medium.

**CULTIVATION OF THE INOCULATED TUBES.**—The pathogenic bacteria all grow best at the temperature of the body, and so all cultures, except gelatin plates, are placed in an incubator during growth. If anaërobic methods are indicated, the tubes must be prepared for the exclusion of oxygen. Wright has devised a simple and satisfactory method, which is a modification of Buchner's two-tube method and does not require any special apparatus. Its principle is based upon the



absorption of the oxygen in the air by pyrogallie acid and caustic soda, leaving an atmosphere of nitrogen. A single tube is inoculated in the usual way. The cotton plugs are then cut off even with the mouth of the tube and pushed in to the distance of three or four centimetres. One cubic centimetre of a ten per cent. aqueous solution of pyrogallie acid is then dropped upon the cotton plug and followed by the same amount of a decinormal solution of caustic soda. The mouth of the tube is then quickly closed with a close-fitting rubber stopper and the culture-tube is placed in the incubator. Hirshberg<sup>1</sup> has slightly modified this method by mixing the pyrogallie acid and sodium nitrate directly with a second tube of melted agar and then pouring this over a deep stab culture of the inoculated agar tube. Hesse's method of turning sterile oil into a test-tube in which a deep stab culture has been inoculated is often convenient.

**PREPARATION OF THE INOCULUM.**—The materials used for inoculation are either the pathologic products—namely, the secretions and excretions and the solid tissues, which have been collected in sterile bulbs or pipettes from the different cavities of the body and from the organs—or cultures which have been obtained by the inoculation of these products upon culture-media. If the material is in a sterile bulb, break off the end of the tube with sterile forceps and expel the contents, by the application of heat to the bulb,—the warmth of the hand is usually sufficient,—into a sterile capsule (a small Petri dish with a ground-glass tightly-fitting cover). The injecting syringe is then filled from the capsule. If the fluid is too thick, it may be diluted with a sterile salt solution. Cultures of bacteria may be prepared in the same manner, and solid tissues may be emulsified in a salt solution.

**INOCULATION OF ANIMALS.**—The animal to be used for inoculation should be carefully selected from those commonly employed for experimentation with reference to its own health and to the species required, as indicated by the microscopic examination of the smear preparation. The animals most used for this purpose are the rabbit, guinea-pig, white mouse, rat, pigeon, and domestic fowl. These differ in their susceptibility to the different pathogenic bacteria infesting man, and negative results are sometimes seen because animals are used which are unsuited to the case under observation. Thus, in cases of suspected syphilis the baboon, ape, or monkey must be employed, and in

<sup>1</sup> *Journ. Amer. Med. Assoc.*, May 21, 1904, p. 1355.

typhoid inoculations the animal must be specially medicated so as to render it susceptible to the bacterium causing the disease.

The animal should be weighed and the rectal temperature (Fig. 169) taken before inoculation, and both weight and temperature should be recorded at a certain time each day during the period of observation. The same rules apply as to asepsis in the inoculation and to avoidance of contamination of the inoculum as are required in a surgical operation. The instruments, similar to those used by surgeons, are sterilized by boiling, owing to the difficulty of removing chemical disinfectants which even in traces might inhibit the growth of bacteria and so vitiate the experiment. An ordinary hypodermic syringe may be employed, but one should be selected that can easily be cleansed and disinfected. Koch's inoculation syringe is much used; many, however, prefer Roux's or some modification of it. The glass and metal portions of the syringes may be sterilized by boiling, but the washers are injured by frequent boiling and should be disinfected by a 5 per cent. solution of carbolic acid followed by *careful* washing in sterile water.

The site of inoculation varies with the different animals and also with the different varieties of the pathogenic bacteria. The usual methods are the subcutaneous, the intraperitoneal, and the intravenous. In small animals like the mouse the last two methods are rarely used, although a very small dose, one or two minims, of fluid may be injected into the peritoneal cavity. Subcutaneous inoculation is commonly practised in the mouse. A fold of skin is pinched up between the thumb and forefinger of the left hand, the hypodermic needle attached to the barrel of its syringe, filled with the material to be introduced, is thrust into the ridge of skin until it enters the subcutaneous tissue, when the fluid is slowly injected; or a piece of skin is snipped with a pair of sharp-pointed scissors, a probe is pushed into the subcutaneous tissue, making a small pocket, in which a portion of the solid inoculum is deposited. (Fig. 170.) The wound may then be dressed with gauze and sealed with collodion.

The intraperitoneal inoculation is made as follows: The animal is held by an assistant or secured to a table. A broad area over the abdomen is shaven, care being taken not to injure the nipple, and the skin is thoroughly disinfected with a two per cent. lysol solution, which is washed off with alcohol. The entire thickness of the abdominal parietes is then pinched up into a triangular fold, the peritoneal surfaces are slipped one over the other to ascertain that no coil of intestine





FIG. 169.—Method of determining the rectal temperature of a guinea-pig.



FIG. 170.—Animal-holder employed in securing a mouse in position while being inoculated.





FIG. 171.—Method of performing peritoneal injection in a rabbit.



FIG. 172.—Ear method of inoculating a rabbit.



FIG. 173.—Post-mortem examination of guinea-pig, made in Ravel pan. Near the four corners, but not shown in the illustration, are hooks upon which the chains are fastened in order to hold the animal in position.



FIG. 174.—Post-mortem examination of a rabbit.



is included in the fold, and the hypodermic needle is passed through the fold near its base. The fold is then released, but the syringe is held steady. The parietes now flatten out, leaving the needle free in the peritoneal cavity. The fluid is then slowly injected. (Fig. 171.)

Intravenous inoculation is not usually practised in animals smaller than a rabbit. In this animal the posterior auricular vein is the one selected for the operation. (Fig. 172.) If a guinea-pig should be chosen, the jugular vein is selected and an anæsthetic is used, the A. C. E. mixture being preferred for general and cocaine for local anæsthesia. The animal is held by an assistant or securely wrapped in a towel fastened with pins, the selected ear is grasped by the root and stretched forward towards the operator, the dorsum of the ear having been previously shaven and cleansed. The syringe is held as one holds a pen, and the needle is thrust through the skin and into the vein itself, being pointed in the direction of the blood stream. The inoculum is then slowly injected and the needle withdrawn.

In the selection of a site for inoculation the general rule is to inject each variety of micro-organism into the kind of tissue which it most often infests in man. For instance, if the lymph-glands are the primary seat of invasion, the animal should be inoculated in that portion of its body which will afford the quickest means of carrying the infectious material to the lymphatics; as, in the test for glanders the bacillary matter is introduced into the testicle, which contains an abundance of these vessels and but a scant blood supply. Or, if the brain and spinal cord are most affected by the disease, as in rabies, an intracranial inoculation is preferred. Thus, the lesions found during the autopsy and the bacteria seen in the microscopic specimens will give a clue regarding the most desirable seat for the inoculation.

POST-MORTEM EXAMINATION.<sup>1</sup>—The post-mortem examination of animals dying from disease produced by experimental inoculation should always be made as soon as possible after their death, and not later than twelve hours. The animal is fastened on a board, ventral surface upward, by nails driven through the extremities, or by being tied with string to special contrivances placed at the corners. Its fur is then wet with a weak antiseptic liquid, such as a 2 per cent. lysol or

---

<sup>1</sup> The reader is referred to the chapter on comparative postmortems for a fuller description of the technic, especially in necropsies on the larger animals, as the cat, dog, sheep, cow, and horse. The methods in vogue for performing a bacteriologic postmortem are seen in Fig. 173 (guinea-pig) and Fig. 174 (rabbit).

a 5 per cent. carbolic solution, to avoid contamination, to prevent the flying off of hairs, and to kill any vermin which are so often present.

The tray of the sterilizer containing the necessary instruments, such as scissors, scalpels, forceps, etc., and the slides, cover-slips, and culture-media, are placed where they may be easily reached. Then, with sterile forceps and scalpel, the skin of the animal in the middle line is incised from the top of the sternum to the pubes, or, if scissors be used for this purpose, the reverse order is followed. Two other incisions are made at right angles to this line through the axillæ and groins. The skin is next reflected in flaps, and may be tacked to the board for the sake of security. The seat of inoculation is then inspected, and, if any lesions are visible, the surface is seared and material removed for cultures and smears. The surface of the thorax is seared and the ribs are divided on each side of the sternum, the chest-plate being removed in the usual way. The pericardial sac is burnt through with the searing iron or incised with a scalpel. The right ventricle of the heart is seared, and into it the pipette is passed and filled with blood for the preparation of smears and inoculation of tubes. Through a seared tract in the middle line of the abdominal wall an incision is made and a specimen of the peritoneal fluid collected. A specimen of the urine is now saved in the same manner as described for collecting the blood. The spleen is excised and placed in a sterile capsule, and through its seared surface the spear-headed spatula is plunged and twisted around so that the eye is filled with material for microscopic and cultural purposes. This process may need to be repeated several times. The other organs, as the lungs, liver, kidneys, lymphatic glands, etc., are removed in a like manner, and all the cavities of the body carefully examined. Specimens of the various tissues may be incised into cubes and placed in fixing fluids for future sectioning. Eyre suggests that a different knife or separate sterilization is needed for cutting each organ. After use the instruments are sterilized and disinfected by boiling. The animal is wrapped in a cloth moistened with an antiseptic solution, as pure formalin, and cremated. Every precaution should be taken to prevent dispersion of the pathogenic bacteria,—as, *e.g.*, by the dropping of cover-glasses, which on becoming broken might cause infection later on. Thus, von Székely found some dried-up gelatin cultures of anthrax and of malignant œdema, eighteen and one-half years old, to be still virulent to white mice.

## CHAPTER XXIV

### WEIGHTS AND MEASURES

It is customary in this country and in England to give the weights of the organs in avoirdupois ounces, their dimensions in inches, and their capacity in cubic inches, though the metric system has more to commend it and is fast gaining favor in English-speaking countries. Troy weight is sometimes used and may give rise to much confusion.

The grain is the same in both Troy and avoirdupois weights. The ounce avoirdupois is 437.5 grains, or 28.34 grammes. The ounce Troy is 480 grains, or 31.1 grammes. To convert grammes into avoirdupois ounces divide by 28.34, into Troy ounces divide by 31.1. To convert grammes into grains divide by 0.065. Conversely, to convert ounces avoirdupois into grammes multiply by 28.34; Troy ounces multiply by 31.1. To convert grains into grammes multiply by 0.065.

A kilogramme equals one thousand grammes, or 2.2 pounds. A gramme equals one thousand milligrammes, or 15.433 grains. A metre equals one thousand millimetres, or 39.37 inches. A litre equals one thousand cubic centimetres, or 61.027 cubic inches, and is equivalent to 2.113 American pints or 1.76 English pints.

An organ should be measured before it is weighed, with the exception of the intestine, which is preferably measured after it has been opened in its entirety, cleansed, and weighed. Measurements should be made as nearly as possible under similar conditions. In the healthy body size is dependent upon sex, age, height, and weight of the subject, while in morbid states it differs according to the above conditions and the disease present. Some parts are preferably weighed before opening, others after opening, and still others both before and after the incisions are completed. Letulle gives the following table:

Weighed before opening.	Weighed after opening.	Both before and after opening.
Pineal gland.	Alimentary tract.	Brain.
Pituitary body.	Aorta.	Liver.
Spinal cord.	Bladder.	Lungs.
Spleen.	Glands, including salivary.	Kidneys.
Suprarenals.	Heart.	Uterus.
Thymus and thyroid.	Pancreas.	



## I. Average height (European standard) :

Adult male .....	172 centimetres, or 5 feet 7.7 inches.
Adult female .....	160 centimetres, or 5 feet 3 inches.
New-born male .....	47.4 centimetres, or 18.66 inches.
New-born female.....	46.75 centimetres, or 18.4 inches.

When a child is two years old, it is about one-half as tall as it will be when fully grown. The rule that a child has usually attained double its birth weight at the fifth month and triple at from the twelfth to the fourteenth month is convenient and useful in estimating an infant's probable age. Keating's *Cyclopædia of the Diseases of Children* gives some valuable information on the physical conditions of childhood.

## II. Average weight (European standard) :

Adult male.....	65 kilogrammes, or 143 pounds (av.).
Adult female .....	55 kilogrammes, or 121 pounds.
New-born child.....	3250 grammes, or 7.15 pounds.

The American Insurance standard: <sup>1</sup>

A man of five feet and one inch should weigh.....	120 pounds.
A man of five feet and three inches should weigh.....	130 pounds.
A man of five feet and six inches should weigh.....	143 pounds.
A man of five feet and nine inches should weigh.....	155 pounds.
A man of five feet and eleven inches should weigh....	165 pounds.

A child may be born weighing less than a pound and live. The greatest recorded weight attained by man is some 1000 pounds. Gould and Pyle in their *Anomalies and Curiosities of Medical Literature* quote from the *Medical Press and Circular* an instance of a man born in North Carolina in 1798 who measured 7 feet 8 inches in height and who weighed over 1000 pounds.

According to Orth, the mean length of a full-term, sound child is between fifty and fifty-one centimetres, the male being slightly longer than the female. The average weight of a full-term boy at birth is thirty-six hundred grammes, that of a girl thirty-two hundred and fifty grammes. For the last five lunar months of fetal life, if the height expressed in centimetres be divided by five, the approximate age of the child in lunar months will be obtained. For example, if the child measures thirty-five centimetres, we divide this by five, and we have seven, which is the number of months which the child has passed

---

<sup>1</sup> From FINLAYSON'S *Clinical Manual*.

*in utero*. The fetal age of the child in the first five months about equals the square root of the height expressed in centimetres. For example, if the height is sixteen centimetres, the child is four lunar months old. In terms of the English system the length of the new-born child is twenty inches, which divided by two will give approximately the number of lunar months that the child has passed *in utero*.

Embryos<sup>1</sup> about one millimetre long are about twelve days old; 2.5 mm., fourteen days old; 4.5 mm., nineteen days old; seven mm., twenty-six days old; 11.5 mm., thirty-four days old; seventeen mm., forty-one days old. For all embryos from one to one hundred mm. long, multiply the length of the embryo from the vertex to the breech in millimetres by one hundred and extract the square root; the result will be the age in days. For embryos from one hundred to two hundred millimetres long, measure from vertex to breech; this length in millimetres will equal the age expressed in days.

Lambinon<sup>2</sup> gives the following figures, obtained at the Liège Maternity, as to the weight of the placenta in cases of miscarriage. The average weight of the placenta at six weeks was 20 grammes (about 5 drachms); at ninety days, 67 grammes ( $17\frac{1}{3}$  drachms); at one hundred and twenty days, 111 grammes ( $28\frac{2}{3}$  drachms); at one hundred and sixty-five days, 262 grammes ( $67\frac{2}{3}$  drachms); and at two hundred and thirty-five days, 330 grammes ( $85\frac{1}{4}$  drachms). The average weight of the placenta at term is a little over a pound (500 grammes). Under similar conditions to the above the length of the umbilical cord averages one-half a metre. Its weight is 27 grammes.

According to Hirst, the following are the dimensions of a full-term, healthy child: Length of hair, from two to three centimetres; anterior fontanel, from two to two and one-half centimetres; occipito-frontal circumference, thirty-four and one-half centimetres; occipito-frontal diameter, eleven and three-fourths centimetres; occipitomenal diameter, thirteen and one-half centimetres; bisacromial diameter, twelve centimetres; intertrochanteric diameter, nine or ten centimetres. The width of the large fontanel may be stated to be from two to two and a half centimetres.

<sup>1</sup> MALL, *Bull. Johns Hopkins Hosp.*, vol. xiv, no. 143, February, 1903, p. 29; abstracted in *Medicine*, vol. ix, no. 3, 1903, p. 240.

<sup>2</sup> *De la détermination de l'âge du fœtus d'après le poids du placenta dans les cas de fausse couche*. Paris, 1898.

John C. Cook<sup>1</sup> cites Fehling as giving the percentage of water in a very young foetus as 97.5 per cent.; after birth, 74.7. In the adult it is 58.5 per cent. Katz<sup>2</sup> has made some interesting chemical analyses of muscle taken from man and from the lower animals. The younger the animal the more water will it contain, the cardiac muscle containing the least water. Horse-flesh can be told by its high iodine index and specific blood-test.

### III. Table of approximate weight of the internal organs:

	Adult, grammes.	New-born, grammes.		Adult, grammes.	New-born, grammes.
Brain.....	1397	385	Both kidneys.....	299	23.6
Heart.....	304	24	Testicles.....	48	0.8
Lungs.....	1172	58 <sup>3</sup>	Ovary.....		0.5
Liver.....	1612	118	Muscles.....	29,880	625
Pancreas.....	201	11.1	Skeleton.....	11,560	445
Right kidney.....	141		Thymus.....		8.5
Left kidney.....	150		Spleen.....		8.5

### IV. The body weight by percentage:

	Adult, per cent.	New-born, per cent.		Adult, per cent.	New-born, per cent.
Heart.....	0.52	0.89	Liver.....	2.77	4.39
Lungs.....	2.01	2.16	Brain.....	2.37	14.34
Stomach and alimen- tary canal.....	2.34	2.53	Thymus gland.....	0.0086	0.54
Pancreas.....	0.346	0.41	Skeleton.....	15.35	16.70
			Muscles.....	43.09	23.40

In measuring an organ its length, breadth, and thickness may often be more quickly and accurately ascertained by thrusting the steel rule through it than in any other manner.

### THE SKULL AND ITS CONTENTS.

*Shape.*—Even in members of the same race the form of the skull is subject to marked variations, and these are still greater when individuals of different races are compared. The characteristic measurements of the cranium are its length, height, and breadth. The cephalic index is the ratio of its length (taken as one hundred units) to its breadth. The altitudinal index is the ratio of its length to its height.

<sup>1</sup> *Jr. Amer. Med. Assoc.*, June 6, 1903, p. 1548.

<sup>2</sup> *Arch. ges. Physiol.*, 1896, vol. lxiii, no. 1, p. 1.

<sup>3</sup> The right being about 5 grammes the heavier.



The accepted horizontal plane is that passing through the upper edges of the external auditory meatus and the lower orbital margin.

According to the variations of the cephalic index, we distinguish the *dolichocephalic* (index less than 75) and the *brachycephalic* (index more than 80) types. Intermediate forms are called *mesocephalic*. If the ratio of the breadth to the height is less than 70, the skull is *platycephalic*; if between 70 and 75, *orthocephalic*; if above 75, *hypsicephalic*. The character of the facial profile is indicated by the *facial angle* of Camper,—namely, the angle between a line on the level of the external auditory meatus and the floor of the nasal cavity and a line touching the middle of the forehead and the anterior portion of the alveolar process of the superior maxilla. If this angle be 80 degrees or more, the skull is called *orthognathous*; if it is between 80 degrees and 65 degrees, *prognathous* (Gegenbaur).

Pathologic types of skull are due in part to premature synostosis. Among them we distinguish the *hydrocephalic* type (from dropsy of the ventricles), the *cephalonic* (or big head), the *microcephalic* (or small head), the *dolichocephalic* (or long head), the *sphenocephalic* (or wedge-shaped head, due to compensatory development of the anterior fontanel), the *leptocephalic* (or narrow head), the *clinocephalic* (or saddle-shaped head), the *trigonocephalic* (or triangular head, due to narrowing of the frontal bone from fetal synostosis of the frontal suture), the *brachycephalic* (or short head), the *pachycephalic* (in which the bones of the cranium are thickened), the *oxycephalic* (or pointed head), the *platycephalic* (or flat head), the *trochocephalic* (or round head), and the *plagiocephalic* (or unsymmetrical oblique head).<sup>1</sup>

*Weight.*—The maximum weight of the adult male encephalon is about 2222 grammes, or 74 ounces, and the minimum is about 960 grammes, or 34 ounces. The average is about 1400 grammes, or 49.5 ounces. The maximum weight of the adult female encephalon is about 1585 grammes, or 56 ounces, and the minimum is 880 grammes, or 31 ounces. The average is from 1230 to 1245 grammes, or from 43½ to 44 ounces. Thus it will be seen that the adult male brain is on an average four or five ounces, or about nine per cent., heavier than that of the female. See also *American Medicine*, May 17, 1902, p. 830.

<sup>1</sup> ZIEGLER'S *Text-Book of Special Pathological Anatomy*, English Translation by MACALISTER and CATTELL, vol. i, pp. 206, 207.

Table showing in grammes the mean weights of the brain at different ages in the two sexes:

	Male.	Female.
Children stillborn at term .....	393	347
Children born alive at term .....	330	283
Under three months of age.....	493	451
From three to six months .....	602	560
From six to twelve months .....	776	727
From one to two years.....	941	843
From two to four years .....	1,095	990
From four to seven years.....	1,138	1,135
From seven to fourteen years....	1,301	1,154
From fourteen to twenty years.....	1,374	1,244
From twenty to thirty years.....	1,333	1,237
From thirty to forty years .....	1,364	1,220
From forty to fifty years.....	1,351	1,212
From fifty to sixty years.....	1,343	1,220
From sixty to seventy years .....	1,313	1,208
From seventy to eighty years.....	1,288	1,168
Over eighty years.....	1,283	1,125

By the above table it appears that the brain is relatively heavier between fourteen and twenty years of age than at any other period; but according to Broca, and also Peacock, the maximum is attained between the ages of twenty-five and thirty-five.

Orth quotes Meynert, whose results were obtained from the investigation of 157 cases in the Vienna insane asylum. He gives the mean weight of the brain, in men between the ages of twenty and sixty-nine years, as 1296 grammes; in women, 1169 grammes. He says the maximal weight is attained during the fourth decade in men and the fifth decade in women. The average weight of the cerebrum is 1018 grammes in men and 917 grammes in women; of the brain stem, 143 grammes in men and 129 grammes in women; of the cerebellum, 135 grammes in men and 123 grammes in women. Weisbach found that in sane German-Austrians the brain weighed 1314.5 grammes in men and 1179.52 grammes in women, while the cerebrum weighed 1154.97 grammes in men and 1038.90 grammes in women, the cerebellum 142.2 grammes in men and 125.56 grammes in women, and the pons 17.33 grammes in men and 15.06 grammes in women. Bischoff found the weight of the pia and arachnoid to be from 25 to 40 grammes. Nauwerck quotes Vierordt, who found the mean weight of the brain in men within the ages of twenty and eighty years to be 1359 grammes, in women 1235 grammes. The brain of the recently deceased Japanese

anatomist Taguchi weighed 1920 grammes, the body weight being 49 kilos. Tourgenieff's brain weighed 2120 grammes, while that of Rustan reached 2222 grammes (74 ounces).

The weight of the encephalon relative to that of the body is subject to great variation, but may approximately be put down as 1 to 36.5 in the adult male and 1 to 35.2 or 1 to 36.46 in the female. These figures are based on observations upon persons dying from more or less prolonged disease, but in the cases of a few individuals who died suddenly from disease or accident the average ratio was found to be 1 to 41. The proportion to body weight is much greater at birth than at any other period of extra-uterine life, being about 1 to 5.85 in the male and 1 to 6.5 in the female.

The weight of the human cerebrum also bears a somewhat definite relation to the stature of the individual. The weight in ounces may be obtained for a male by dividing the height in inches by 1.6, and for a female by multiplying the quotient thus obtained by  $\frac{30}{31}$ . The weight in grammes may be obtained by multiplying the height in centimetres by 7 for a male, and the product again by  $\frac{30}{31}$  for a female. Thus,

$$\begin{aligned}\text{Weight in ounces of the mean cerebrum} &= \frac{\text{height in inches}}{1.6} \\ \text{Weight in ounces of the mean female cerebrum} &= \frac{\text{height in inches}}{1.6} \times \frac{30}{31} \\ \text{Weight in grammes of the mean male cerebrum} &= \text{height in centimetres} \times 7 \\ \text{Weight in grammes of the mean female cerebrum} &= \text{height in centimetres} \times 7 \times \frac{30}{31}\end{aligned}$$

These proportions are slightly deficient for the higher and excessive for the lower statures.

*Dimensions.*—The mean cubic capacity of the male cranium is 1450 cubic centimetres; that of the female is 1300 cubic centimetres (Welcker). The length of the male brain is from 160 to 170 millimetres, or from  $6\frac{3}{8}$  to  $6\frac{1}{2}$  inches, and that of the female brain is from 150 to 160 millimetres, or from 6 to  $6\frac{3}{8}$  inches. The greatest transverse diameter is 140 millimetres, or  $5\frac{5}{8}$  inches, and the greatest vertical diameter is 125 millimetres, or 5 inches. The volume is about 1330 cubic centimetres, or 81 cubic inches.

The specific gravity of the brain is from 1035 to 1040.

Pituitary gland, length, .008; breadth, .012; thickness, .065; weight, 5 grammes (Zander).

Pineal gland measures .010  $\times$  .005  $\times$  .005 (Charpy) and weighs 0.20 gramme (Engel).



The length of the spinal cord in the adult is 0.448 metre and its weight, deprived of its nerves, 27 to 30 grammes. The transverse diameter of the cervical enlargement, 0.013; of the dorsal, 0.01; and of the lumbar, 0.012. The anteroposterior diameter of the cervical enlargement, 0.009; of the dorsal, 0.008; of the lumbar, 0.009.

#### THE HEART.

*Weight.*—The mean weight of the heart in the adult male is about 310 grammes, or 11 ounces; its proportion to the body weight is 1 to 169. That of the adult female is about 255 grammes, or 9 ounces; proportion to body weight, 1 to 149. According to Krause, the proportion of the heart weight to the body weight is as 1 to 169 in men and as 1 to 162 in women. The increase in weight from the fifteenth to the seventieth year is about 60 grammes in the male and 25 grammes in the female.

The weight of the heart increases with the body weight, but in a gradually decreasing ratio, until the seventieth year, when it begins to diminish. At birth it is about 24 grammes; proportion, 1 to 130 (Quain).

*Dimensions.*—The determination of the exact measurements of the heart is most difficult, as the muscular fibres contract and expand under such diverse circumstances and the positions of the parts are so different. Constantin Paul has called attention to the fixed manner in which the inferior vena cavity enters into the heart on a line with the tip of the right auricular appendix. The heart is generally of about the same size as the right fist of the cadaver. Its extreme length is about 125 millimetres, or 5 inches; width, 87 millimetres, or 3 inches; thickness, 62 millimetres, or  $2\frac{1}{2}$  inches, slightly less in the female than in the male. The thickness of the wall of the right ventricle is from 2 to 5 millimetres, or  $\frac{1}{8}$  to  $\frac{1}{4}$  of an inch; of the left ventricle, from 7 to 12 millimetres, or  $\frac{1}{4}$  to  $\frac{1}{2}$  of an inch; ventricular septum, 15 millimetres. Pathologically these measurements may be increased threefold or more.

Nauwerck and Orth quote Bizot as follows: The weight of the heart is 300 grammes in men and 250 grammes in women. The length in men is from 85 to 90 millimetres, in women from 80 to 85 millimetres; the breadth in men is from 92 to 105 millimetres, in women from 85 to 92 millimetres; the thickness in men is from 35 to 36 millimetres, in women from 30 to 35 millimetres. The thickness of the

right ventricle without the trabeculæ is from 2 to 3 millimetres in men and slightly less in women; the left ventricle is from 7 to 10 millimetres thick. The heart of the elephant Jumbo is reported to weigh 36½ pounds after having soaked several years in alcohol.

The dimensions of the orifices of the heart are shown in the following tabular statement.

Orifices.	Diameter.	Circumference.		Area.	
		Male.	Female.	Male.	Female.
Aortic.....	24 to 25 mm., or 0.9 to 1 in.	81 mm.	76 mm.	530 sq. mm.	452 sq. mm.
Mitral .....	30 to 35 mm., or 1.2 to 1.4 in.	103 mm.	101 mm.	855 sq. mm.	804 sq. mm.
Pulmonary ....	27 to 30 mm., or 1.1 to 1.2 in.	91 mm.	89 mm.	660 sq. mm.	615 sq. mm.
Tricuspid.....	37 to 45 mm., or 1.5 to 1.8 in.	122 mm.	115 mm.	1194 sq. mm.	1017 sq. mm.

Bizot's figures for the circumference are, as a rule, slightly less.

*Volume.*—In the new-born this is about 22 cubic centimetres, which is increased to 250 centimetres at twenty years and about 280 centimetres at fifty years, after which it gradually decreases. Up to the age of puberty it is about the same in both sexes, but after that it is from twenty-five to thirty centimetres larger in the male. Because of obvious difficulties, these figures can only be regarded as approximate.

Thickness of the aorta, 1½ to 2 millimetres (Orth). Circumference of the thoracic aorta, 4 to 6 centimetres; of the abdominal aorta, 35 to 45 centimetres; weight, 35 to 45 grammes. The transverse sections of the aorta will about admit the thumb. The length of the inferior vena cava is 0.22 to 0.25 metre; of the superior, .06 to .08; of the great azygos, .20 to .25; and of the portal veins, .05 to .12.

The length of the thoracic duct is .30 to .34 metre.

#### THE LUNGS.

*Weight.*—Obviously the lungs are subject to great variation in weight, depending upon the amount of blood or other liquid and of air in their cavities. Their combined weight ranges from 850 to 1370 grammes, or from 30 to 48 ounces, the average being from 1020 to 1190 grammes, or from 36 to 42 ounces (1300 grammes in the male and 1023 grammes in the female.—Krause). The right is generally 2 ounces heavier than the left. The weight of the right lung is from

## THE THYMUS GLAND.

*Weight.*—At birth this gland weighs about half an ounce. In twenty adult cases it was found to average 5 grammes (Quain). Friedleben says that the thymus weighs at birth 14 grammes and at nine months of age 20 grammes. Up to the second year it weighs a little more than 26 grammes, and from the third to the fourteenth year a little less than this.

*Dimensions.*—At birth the length is about 60 millimetres, or 2 inches; width, 37 millimetres, or  $1\frac{1}{2}$  inches; and thickness, from 6 to 8 millimetres, or  $\frac{1}{4}$  to  $\frac{1}{3}$  of an inch. From birth to the second month the length is 5.2 centimetres; from the ninth month to the second year, 6.96 centimetres, and from the third to the fourteenth year, 8.44 centimetres. The breadth across the middle is from 2.7 to 4.1 centimetres; above and below, from 0.7 to 0.9 centimetre (Friedleben).

## THE THYROID GLAND.

*Weight.*—From 28 to 56 grammes, or 1 to 2 ounces, being larger in the female. Orth gives the weight as from 30 to 60 grammes.

*Dimensions.*—Each lateral lobe is about 50 millimetres, or 2 inches, in length; from 18 to 30 millimetres, or  $\frac{3}{4}$  inch to  $1\frac{1}{4}$  inches, in breadth; and from 18 to 25 millimetres, or  $\frac{3}{4}$  to 1 inch, in thickness. The right lobe is usually the larger. The isthmus is nearly 12 millimetres, or  $\frac{1}{2}$  inch, in breadth, and from 6 to 18 millimetres, or  $\frac{1}{4}$  to  $\frac{3}{4}$  inch, in depth. According to Orth, each lateral lobe is from 5 to 7 centimetres long, from 3 to 4 centimetres broad, and from 1.5 to 2.5 centimetres thick.

## THE TESTES.

*Weight.*—Each testicle with its epididymis weighs from 18 to 25 grammes, or 6 to 8 drachms, the left being slightly the heavier. Orth gives 18 to 26 grammes as the weight; Nauwerck says the testicle and epididymis weigh from 15 to 24.5 grammes.

*Dimensions.*—Length, about 37 millimetres, or  $1\frac{1}{2}$  inches; breadth, anteroposterior, 30 millimetres, or  $1\frac{1}{4}$  inches; thickness, from side to side, 24 millimetres, or 1 inch.

## THE OVARIES.

*Weight.*—From 4 to 8 grammes, or 1 to 2 drachms. Orth gives 5 to 7 for the weight, and Nauwerck quotes Puech, who puts the mean weight at 7.0 (from 5 to 10) grammes.



*Dimensions.*—Length, usually about 37 millimetres, or  $1\frac{1}{2}$  inches; breadth, 18 millimetres, or  $\frac{3}{4}$  inch; thickness, 12 millimetres, or  $\frac{1}{2}$  inch. The right is usually a little larger than the left. According to Orth, the ovary is from 2.5 to 5 centimetres long, from 2 to 3 centimetres broad, and from 7 to 12 millimetres thick. Nauwerck gives the following dimensions, quoted from Puech.

Length, maidens .....	from 4.1 to 5.2 centimetres.
Length, women .....	from 2.7 to 4.1 centimetres.
Breadth, maidens .....	from 2.0 to 2.7 centimetres.
Breadth, women .....	from 1.4 to 1.6 centimetres.
Thickness, maidens .....	from 1.0 to 1.1 centimetres.
Thickness, women .....	from 0.7 to 0.9 centimetre.

#### THE UTERUS AND BREASTS.

*Weight.*—Generally from 28 to 42 grammes, or 1 to  $1\frac{1}{2}$  ounces. Orth quotes Huschke, who gives from 33 to 41 grammes as the weight of the uterus in virgins and 105 to 120 grammes as the weight in multiparæ. Nauwerck gives 33 to 41 grammes as the weight in virgins, and 102 to 117 grammes as the weight in multiparæ.

*Dimensions.*—Length, about 75 millimetres, or 3 inches; breadth, 50 millimetres, or 2 inches; thickness, nearly 25 millimetres, or 1 inch. The virgin uterus is from 5.5 to 8 centimetres long, from 3.5 to 4 centimetres broad, and from 2 to 2.5 centimetres thick; in multiparæ the womb is from 9 to 9.5 centimetres long, from 5.5 to 6 centimetres broad, and from 3 to 3.5 centimetres thick. The walls of the virgin uterus are from 1 to 1.5 centimetres thick; of the cervix, from 0.7 to 0.8 centimetre thick. In multiparæ the uterine walls may be as thick as 2 centimetres, and the cervix is from 0.8 to 0.9 centimetre thick. (Orth.)

The length of the virgin or nulliparous uterus, from the fundus to the external os, is from 7.8 to 8.1 centimetres and the breadth of the fundus is from 3.4 to 4.5 centimetres; the thickness below the fundus is from 1.8 to 2.7 centimetres; the length of the cervix is from 2.9 to 3.4 centimetres; the breadth of the cervix is 2.5 centimetres; the thickness of the cervix is from 1.6 to 2 centimetres. In multiparæ the length of the uterus is from 8.7 to 9.4 centimetres, the breadth 5.4 to 6.1, and the thickness 3.2 to 3.6 centimetres. The length of the uterine cavity in virgins is 5.2 centimetres, after the menopause 5.6 centimetres; in multiparæ 5.7 centimetres, after the menopause 6.2 centimetres. (Nauwerck.)

Breasts at birth, 0.30 to 0.60 gramme; of adult, 150 to 200 grammes; during lactation, 400 to 900 grammes.

#### THE PROSTATE AND SEMINAL VESICLES.

*Weight.*—Average, from 18 to 20 grammes, or  $4\frac{1}{2}$  to  $4\frac{3}{4}$  drachms. Orth gives 17 to 18.5 grammes as the weight; and Nauwerck quotes Krause and Bischoff, who give 19 to 20.5 grammes as the weight.

*Dimensions.*—Transverse diameter, about 37 millimetres, or  $1\frac{1}{2}$  inches; vertical, 30 millimetres, or  $1\frac{1}{4}$  inches; anteroposterior, 18 millimetres, or  $\frac{3}{4}$  inch. These measurements are subject to great variation, according to the fulness of the rectum and bladder. According to Orth, the prostate measures from 32 to 45 millimetres in its transverse diameter, 14 to 22 millimetres in thickness, and 25 to 35 millimetres from apex to base. Nauwerck gives the following dimensions: Transverse diameter (breadth), from 3.2 to 4.7 centimetres (mean, 4.5 centimetres); sagittal diameter (thickness), from 1.4 to 2.3 centimetres (mean, 2 centimetres); from apex to base (height), from 2.3 to 3.4 (mean, 2.7 centimetres).

The seminal vesicles measure 4.2 by 17 by 0.9 centimetres.

## CHAPTER XXV

### COMPARATIVE POSTMORTEMS <sup>1</sup>

THE great number, importance, and variety of diseases which human beings may contract from the lower animals are more and more coming to be recognized. Our domestic animals suffer from nearly all the contagious maladies found in man, and impart to him various disorders from which he would otherwise be exempt, such as glanders, actinomycosis, anthrax, hydrophobia, foot-and-mouth disease, echinococcus cysts, trypanosomiasis, etc. The rat disseminates bubonic plague, the mosquito malaria, yellow fever, and dengue, and the pig trichinosis, and were it not for the rat, the mosquito, and the pig these diseases would probably cease to exist.

The skin, extremities, joints, excessive functionation of the mammary gland, and the frequency of parasitic lesions in the muscular tissue are so often subject to pathological conditions that they present a rich field in post-mortem examinations of lower animals. Malformations are also quite common.

There has recently taken place an interesting discussion as to Koch's statements that human tuberculosis differs from bovine and cannot be transmitted to cattle, and that man does not, except in the rarest instances, contract tuberculosis from the cow. Both sides admit, however, that there is a great difference between the virulence of various forms of the tubercle bacilli. As we go to press, Koch promises the publication of experiments carried on while he was in Africa which will completely prove his original assertions, while the British Royal Commission report that they have reached positive conclusions which refute the claims of the celebrated German bacteriologist. In Switzerland and in this country the writer has been struck with the freedom from tuberculosis of districts in which cow's milk is not used.

Many of the suggestions made in the previous chapters apply with equal force to the performance of necropsies upon the lower animals. Such comparative examinations are of two distinct classes,—veterinary

---

<sup>1</sup> Much of the material and all the illustrations in this chapter are taken from KITT's excellent work entitled *Lehrbuch der pathologischen Anatomie der Haustiere*, 1900, vol. ii, pp. 1-54.



postmortems and laboratory postmortems. For laboratory study small animals, such as the guinea-pig, rabbit, mouse, and rat, are generally chosen, while in veterinary investigation the subject is usually a dog, a horse, a cow, or a cat. So intense is the interest now taken in comparative pathology that all classes of animals come to section, even reptiles (especially snakes) receiving no small amount of attention.

**INSTRUMENTS.**—In post-mortem examinations of the large domestic animals (cow, horse, mule, etc.) the instruments used must necessarily be larger than those employed in human autopsies. The following is a partial list. (1) Large butcher's knife, to expose the thorax and abdomen and remove the skin; (2) large cleaver; (3) large butcher's saw, to open the thoracic and cranial cavities, expose the nasal septum, etc.; (4) large chisel, to remove the cord; (5) hammer, for the same purpose; (6) bone-forceps (costotome); (7) enterotome; (8) scissors; (9) brain-knife; (10) dissecting forceps; (11) large needle; (12) strong twine, etc.

**UTENSILS.**—Buckets, pitchers, large and small enamelled plates, sponges, soap, towels, and disinfectants, and green soap or lysol are especially useful.

**CLOTHING.**—An operator's apron may be drawn over the clothes or an ordinary rain-coat worn, but a special suit for operating is better.

**GENERAL SUGGESTIONS.**—In many cases the necropsy must be made at the place where death occurred, be this in the fields, stable, slaughter-house, or veterinary morgue. The procedure will vary with the conditions and conveniences, but the end in view should be carefully considered and certain general rules observed. If the animal is alive, the method of killing to prepare for the desired investigation should be one that will not injure the organs involved. In cerebral trouble the animal should not be killed by a blow upon the head, but by poison or chloroform; in inflammatory conditions all loss of blood should be avoided; if the trouble is in the digestive system, no poison should be used; and in pulmonary affections the animal must not be shot through the heart (Csokor).

**OPERATIVE TECHNIC.**—In opening the cadaver the normal position of the intestines should be retained as far as possible, and they should be carefully examined to see that they are uninjured and are sufficiently exposed. Horses, large and small ruminants, and the larger swine are usually placed upon the left side of the body so that the right side may be opened. A dorsal position may be chosen for dogs and

cats, and even for swine or larger animals if sufficient assistance be present, as it gives a better view of the abdominal cavity.

The postmortem is begun by removing the hide, which has a market value and must not be injured. As scalpels and straight-edged knives are apt to button-hole the skin, a butcher-knife with rough cutting edge is to be preferred. Beginning at the angle of the chin a longitudinal incision is made down the median line the whole length of the body, avoiding the udder, prepuce, and scrotum, and the navel in the case of young animals. A transverse incision is made perpendicular to the first along the median surface of the foreleg and the skin is drawn back from the edges up over the dorsal surface. A similar cut is made upon the median surface of the thigh and leg down to the tuberosity of the os calcis. On both the limbs and the body the hair-seams will serve as a useful guide for the knife. A circular incision is made around the head from angle to angle at the lips. If the head is to be preserved, as in the case of a deer, the circular incision is made at the manubrium. The skin may be detached either with the hands or with the handle of a chisel.

**REMOVAL OF THE EXTREMITIES.**—After the animal has been skinned, it is placed on its side, and the uppermost limbs are removed in order to secure more room for subsequent manipulation. First the foreleg is held up by an assistant and the shoulder-joint disarticulated. The musculature of the part is cut through in the median portion by a butcher-knife grasped firmly by the whole hand. During the exsection the extremity should be constantly raised by an assistant and the blade of the knife should be held somewhat towards the thorax so as to cut obliquely to the ribs.

To remove the posterior extremity make a deep circular incision through the hip muscles, beginning with the broad crural fascia and above the large trochanter, passing up over and through the musculature of the croup and downward and outward into the ischiatic fossa, but not behind the tuberosity of the ischium; raise the foot; cut through the adductors in a line with the acetabulum, open its capsular ligament, and section the round ligament. The incision of the capsular ligament is accompanied by a snapping sound, due to the entrance of air into the joint. The limb can now be drawn backward, the remaining fascia and muscles sectioned, and the whole removed.

**EXPOSURE OF THE ABDOMINAL CAVITY.**—Before opening the abdominal cavity of a filly the udder should be entirely removed from the

abdominal wall, and in geldings and stallions the scrotum and the penis should be isolated and thrown back. It should be remembered that in herbivora meteorism occurs soon after death, so that the intestines are pressed up closely against the abdominal wall and may easily be injured.

The operator should stand in the space between the remaining extremities with his face towards the breast of the animal. An incision is made through the median line of the body, beginning with the ensiform cartilage of the sternum, extending as far as the pubic region, cutting through the muscles and fascia only and not injuring the peritoneum. This will not be difficult if the blade of the knife be held flat and the ball of the thumb placed near the edge and close to the point. As the peritoneum is carefully torn through with the fingers, the exit of gases, liquids, or abnormal contents of the abdominal cavity should be noted, as well as the position of the intestines. The index and middle finger are then separated so as to form a V-shaped space, in which the knife is placed and its point thrust through the abdominal wall along the line of the linea alba, the fingers following. At the posterior end of the longitudinal incision a second incision is made, perpendicular to the first, extending from the pubic region to the lumbar. The right upper half of the abdominal wall is held up by its edges with the left hand. The assistant pulls on the lower ribs in order to make the abdomen tense, and its covering is cut through with sawing strokes of the knife as far as the costal processes. The knife is so held by the whole hand that the point is shoved away from the operator towards the lumbar region and the lower part of the blade is used instead of the point.

We have now a large anterior and a small posterior segment of the abdominal wall. They may easily be drawn back and a view of the abdominal organs obtained. The ribs of the horse extend so low down that a sufficiently extensive view for pathologic purposes cannot be obtained; therefore, before removing the abdominal contents the thoracic cavity is exposed. Then, by thrusting the hand well up under the lower ribs, we notice whether the diaphragm is tightly vaulted forward or is more or less relaxed.

**EXPOSURE OF THE THORACIC CAVITY.**—A small incision is made between two of the true ribs and note is taken whether or not air enters the thoracic cavity and the diaphragm becomes relaxed. If the abdominal examination showed the diaphragm drawn down posteriorly,



the incision should receive special attention; instead of air entering, there may be an exit of gas from the pleural cavity, indicating some essentially pathologic condition.

The direction for cutting the ribs is through the costal angles following the course of the iliocostal muscle. An incision is made




FIG. 175.—Equine viscera, the animal resting on its right side, the anterior and posterior left limbs having been removed, and the abdominal, thoracic, oral, and pharyngeal cavities opened. The double lines show the places in the intestines which are to be tied previous to being cut.

between the true ribs and the blade of the saw introduced, an assistant making the breast tense while the sawing is done; very little pressure should be used or the bone will splinter. When the ribs have been sawed through, they are turned over towards the median line and removed by severing the costal cartilages. The situation of the organs and the pathologic contents should be carefully noted. (Fig. 175.)

**REMOVAL OF THE ABDOMINAL CONTENTS IN THE RIGHT LATERAL POSITION.**—After exposing the abdominal cavity by the longitudinal and transverse incisions, pull the two left coils of the colon either up over the thorax or out across the body on the right side, so that the sigmoid flexure looks towards the head or lies on the ground and the body and tip of the cæcum come into view. Spread the mesorectum out over the left flank and pelvic region. Stroke back the fæces, doubly




FIG. 176.—Further dissection of animal seen in Fig. 175. Appearance of the parts after removal of the rectum, ileum, and jejunum.

ligate the rectum at its entrance into the pelvis, and section. Cut away the mesorectum up to its origin at the rectoduodenal ligament, doubly ligate the rectum, section, and remove.

The ileum is easily recognized by its thicker walls and its entrance into the cæcum. Apply a double ligature, section, and, holding the intestine in the hand, cut away all the mesentery from the whole of the small intestine as far as the rectoduodenal ligament, divide this, doubly ligate the duodenum, and section. The junction of the colon

with the rectum is now exposed,—the so-called stomach-like or gastroid dilatation,—under which lies the anterior root of the mesentery. Grasping the dilatation with the left hand (Fig. 176), pull it towards the cæcum, and with the right hand work loose or cut partly away the connections between the gastroid dilatation and cæcum and the omental sac, kidney, and pancreas. In this way better access to the portal vein and anterior root of the mesentery is obtained. With the fingers work through the cellular tissue surrounding the root of the mesentery, grasp it with the hand, and together with the portal vein cut it away close to the intestine, leaving as much of it as possible with the aorta. The colon and cæcum are now drawn out of the cavity, all the remaining sections being easily torn or cut away, while the right branch of the pancreas which lies upon the cæcum and the root of the mesentery must be carefully dissected away. Grasp the spleen, section the suspensory (gastrosplenic) ligament and the gastrosplenic omentum, and free the spleen from the stomach. Separate the branches of the pancreas from the larger blood-vessels and the kidneys, so that it hangs only by its body from the liver, and leave it in this position or, after examining its excretory duct, cut it away. Next remove the stomach and duodenum by cutting along the sigmoid curvature and the smaller curvature of the stomach and by sectioning the duodenorenal ligament, the hepatic and pancreatic ducts, the diaphragmatic and gastrohepatic ligaments, and the œsophagus, after pulling the latter down as far as possible from the diaphragm. Excision of the liver is an easy matter: section first the left lateral ligament, then the coronary and suspensory ligaments, the vena cava on the anterior surface of the liver, the right lateral portion of the coronary ligament, and the right hepatic and renal hepatic ligaments.

**REMOVAL OF THE ABDOMINAL CONTENTS IN THE LEFT LATERAL POSITION.**—The rectum is sectioned at its entrance into the pelvis after pressing back the fæces with the fingers, applying a double ligature, and cutting between them. Seize the colon at its anterior curvature and pull it carefully out of the abdominal cavity as far as possible. The left folds of the colon will fall out with very little assistance. (Fig. 177.)

In the region of the kidney will be seen the arch of the duodenum lying between the anterior and posterior roots of the mesentery and covered by the ribs. Cut through this arch and its mesentery, after applying a double ligature, and remove. The cellular tissue lying



between the cæcum and psoas muscle and the right kidney should be carefully worked loose and the pancreas separated from the cæcum and the colon; this is done by tearing or cutting through the peritoneum covering the intestine and pancreas, getting the hand in under the pancreas, and working it loose. Beginning posteriorly, cut away the mesorectum from behind forward and any connections that may remain between the cæcum and colon and the region of the kidney,




FIG. 177.—Further dissection of animal seen in Figs. 175 and 176. Appearance of the parts after removal of the large intestine.

grasp as much as possible of the attachment of the mesentery, pull the intestine back away from the kidney, and section the root of the mesentery in front of the left hand, as far from the aorta as possible. With the exception of a small portion of the duodenum and the pelvic portion of the rectum the large and small intestines can be drawn out from the abdominal cavity by cutting or tearing away any attachments which may remain; the operator stands alternately at the back and in front of the cadaver while removing these portions.

**REMOVAL OF THE KIDNEYS, STOMACH, LIVER, AND SPLEEN.**—The removal of the kidneys leaves a freer field for the stomach, spleen, and liver. With the hand and fingers separate first the right and then the left kidney and the suprarenal capsules from the surrounding cellular

tissue. If the ureters and kidneys are intact the kidneys may at once be cut away together with their vessels. In case of any abnormalities they should be left hanging or a sufficient length of the ureters removed with them, together with the surrounding tissues, or they may remain attached to their ureters and placed in the pelvic region.

The pancreas, spleen, and stomach are freed from the mesentery and sectioned. The assistant pulls on the right side of the diaphragm, and the inferior vena cava between it and the liver is cut through together with the œsophagus and the right hepatic ligament. The stomach is turned backward. The left hepatic ligament is sectioned and all the three organs removed together in a mass.

If the kidneys are left in place, the exenteration of the stomach, pancreas, and liver is more difficult and demands more caution, especially if the animal has not been bled, because the field is obscured by blood and other impurities. Dissect away carefully the attachments of the right kidney to the suprarenal capsule and left branch of the pancreas, which lies deep down, covered by the branches of the mesenteric arteries; next the adrenals, then the fundus of the stomach from the cruræ of the diaphragm, the suspensory ligament of the spleen, the splenorenal ligament, the right coronary and lateral ligaments of the liver, the hepatic renal ligament, the vena cava, with the falciform ligament, the œsophagus, and the left lateral and coronary ligaments of the liver.

All these organs may be removed with the diaphragm, and, when there are adhesions to its posterior surface, this is the preferable method. The right lobe of the liver is first separated from the kidney; the pancreas, spleen, and stomach are worked loose from the spinal column; the posterior vena cava, the œsophagus, and the pulmonary attachments to the diaphragm are sectioned; the diaphragm is freed from the thoracic wall by a circular excision, and the whole mass removed together. Finally, the aorta and the venæ cavæ with their branches are dissected off the spinal column from the diaphragm to the pelvis.

**EXENTERATION IN THE DORSAL POSITION.**—The body may be kept on its back by tying the feet to rings in the wall or to posts or poles. The extremities remain attached to the body, of course, and the broad muscles of the chest are only to be sufficiently incised to permit the anterior extremities to spring out a little and give access to the chest. If during the postmortem the extremities are released too

much, the body will fall to one side and make the exenteration more difficult.

A longitudinal median incision is first made, then a bilateral transverse incision just posterior to the last ribs. The two left folds of the colon are drawn up over the right side of the body. The rectum is pulled out and spread over the left thigh and left ventral wall and the small intestine spread out over the region of the lower ribs. The ileum is found at its insertion into the cæcum; it is thicker than the rest of the small intestine. It is tied off and sectioned, remaining in the hand after its mesentery is severed close up to the intestine. In this way the whole of the right lateral small intestine is removed from the abdominal cavity and its mesentery left hanging by its root. When it passes into the duodenum between the two roots of the mesentery, doubly ligate and section. Doubly ligate and section the rectum at its entrance into the pelvis and again at its junction with the colon.

The pancreas and first part of the duodenum are dissected away from the colon as in the first method. The roots of the mesentery and both the branches going to the large intestine are sectioned close up and the large intestine is removed.

The stomach, spleen, etc., are removed as in the first method. Many operators prefer to excise the spleen and open the stomach along its greater curvature and the duodenum on its inferior surface, whereupon the pathway of the bile-ducts may be determined and then the empty organs cut away.

In the dorsal position the thoracic organs may be ablated by drawing them down towards the abdominal cavity. An incision is made between the rings of the trachea, two fingers are inserted, the trachea is grasped firmly, and the larger vessels are sectioned at the thoracic inlet; the aorta is dissected away from the vertebræ and the posterior vena cava and œsophagus are sectioned. If it be desired to remove the thoracic viscera together with the trachea and cervical organs, the first rib is sawed through and excised; the cervical organs are then ablated according to the method to be described later.

**VIENNA METHOD OF EXENTERATION IN THE LEFT LATERAL POSITION.**—Csokor's quick method for removing the thoracic and abdominal contents is as follows: The extremities are removed and the abdominal cavity is exposed by a longitudinal and a transverse incision as in the first method; then the muscles of the back are cleared away and the sectioned abdominal wall is drawn up by a hook. With a



hatchet each rib is cut away from the spinal column and then from the breast-bone. The whole right wall of the thorax and abdomen is now drawn up over the head of the animal and the contents of both cavities are exposed. The right kidney is next removed and then the thoracic contents. After their ablation the cardiac end of the stomach is freed from the diaphragm and the duodenum is detached from the liver and its surroundings and excised together with the stomach and spleen. The abdominal aorta is separated from the spinal column, the rectum sectioned, and all the intestines removed. The remaining organs are extirpated as in the other methods. This modification permits a very rapid necropsy, but the removal of the stomach and spleen is somewhat difficult.

DISCUSSION OF THE ABDOMINAL CONTENTS.—To ascertain the macroscopic conditions of the abdominal contents it is necessary to make a few special incisions. The aorta is first examined and its dorsal wall slit up with the shears to expose the entrances into its branches, which are then cut open. On account of its great frequency, close search is to be made for an aneurism in the root of the mesentery. It is usually felt externally as a thick, cystic expansion. The branches to the small intestine—the duodenal, jejunal, and iliac arteries—are first given off from the short trunk of the artery lying in the root of the mesentery (the anterior mesenteric artery); next a large vessel, the ileocolic artery, which gives off a large branch, the inferior colic, and the ileocaecal artery with its three branches. The superior colic comes off above the root of the mesentery on a level with the anterior rectal artery. After examining these branches slit the inferior and superior colic arteries in the mesocolon from their origin to the sigmoid flexure. If it seems necessary, examine the arteries of the small intestine in the same way and observe the mesenteric lymph-nodes. The bowel is opened with the shears along the line of the attachment of the mesentery so as to get a good view of Peyer's patches; keep the intestine lying flat, for if held up the contents run down into the lower portions, which is a nuisance.

If the stomach is sufficiently full, cut it open with a knife along its greater curvature. If the duodenal portion remains with the stomach and liver, open it with the shears on its inferior surface in such a way that the termini of the hepatic and pancreatic ducts will not be injured and their patency may be demonstrated. Press and push along the course of the ducts so as to force out their contents. If there is any

suspicion of abnormalities in these ducts, it is better to leave the stomach and duodenum in place and to open them before removal.

**REMOVAL OF THE THORACIC CONTENTS.**—First carefully examine for sharp points of bone and excise them with cutting forceps. The pericardium should then be examined and worked free with the hands. The posterior vena cava is tied off and divided between the ligature and the diaphragm; the attachments of the liver and heart to the diaphragm are sectioned and an incision is made obliquely through the aorta down to the vertebral column. Thrust the finger into the posterior aorta, pull it up, and cut along the spinal column in the line of the vena azygos and the attachment of the longus colli. Now make an oblique section through the œsophagus, trachea, anterior aorta, and anterior vena cava along the line of the first rib, so that the thoracic organs may be removed. This avoids cutting the large veins, which bleed so freely as greatly to obstruct the view of the parts under observation.

**SECTION OF THE ORAL CAVITY AND CERVICAL ORGANS.**—This is begun by removing the ramus of the lower jaw on one side. Cut the buccal parietes and the cheek at the angle of the lips up to the zygomatic arch, between the molar teeth and the space between the lower jaw and the large maxillary swelling, dividing the masseter and sawing through the bones. The ramus of the jaw may now be worked up and down, its muscular connections severed by a knife introduced along its median surface, and an incision made between the parotid gland and the posterior border of the bone. The temporal muscle is cut through above the coronoid process and the ligaments and capsule of the joint are sectioned, the jaw being moved up and down to find the joint. After examining the local conditions, sever the left connections of the tongue with the jaw and the soft palate; saw through both to the large branches of the hyoid bone. The larynx, trachea, and œsophagus are easily freed from their loose cellular tissue by cutting into the channel of the external jugular vein, between the longus colli muscle and the œsophagus, so that the thyroid gland is not injured.

**DISSECTION OF THE THORACIC AND CERVICAL ORGANS.**—In order more closely to inspect these organs, cut through the vault of the velum palati with the shears and continue down into the œsophagus, sectioning it dorsally. With the knife grasped firmly incise the larynx in the median dorsal line between the arytenoids. Pushing the œsophagus

aside, cut the posterior muscular ligament of the trachea with shears throughout its whole length and thrust the cartilages apart to get a good view of the interior. The lobes of the lungs are laid open with long, deep, bisecting strokes, and portions of each lobe are tested by throwing them into water to see whether they contain air and will float or will sink because of collapse or the presence of an exudate. The lymph-nodules around the roots of the bronchi should always be examined and sectioned.

If the heart is hacked into or improperly opened, the distinctive appearance of any abnormality that may be present is destroyed, and these anomalies are of great importance to the whole organism. First make an incision into the right ventricle along the septum, insert the shears, and cut up into the pulmonalis. Holding the heart by this flap, lengthen the incisions towards the apex and the flap so as to get a better view of the ventricle. In the same way incise the left ventricle close to the septum and on the anterior surface; insert a finger through the opening, find the entrance into the aorta, and with the shears cut down between the pulmonalis and the left auricle. It is true that in this way both semilunar valves are sectioned, but the auriculoventricular valves are spared and they are much more likely to present abnormalities than the semilunar. The size of the openings can be tested by inserting a finger, and the thickness of the walls measured, after which each auricle is cut through up into its vessels and a good view of their openings obtained.

EXENTERATION OF THE PELVIS.—The removal of the pelvic organs is preceded by the previously described excision of the kidneys and ureters and in males by the exposure of the testicles and the external genitalia. The scrotum and penis were then turned back, and now their dorsal suspensory ligament and surroundings are divided as far as the ischiatic notch and all the flesh lying ventrad to the ischiatic suture is carefully cleaned away. The scrotum and the right and left inguinal canals are split open and the testicles together with the spermatic vessels pulled up into the abdomen. It is especially necessary to cut the tendinous ligament which binds the corpora cavernosa to the ischium close to the bone, as well as the strong ischiopenile muscle. Two sections made by sawing will remove the right wall of the pelvis. The first one is made through the ischiopubic suture over the acetabulum to the iliac spinal column; the second, through the thin part of the iliac bone, after cutting away the flesh that lies over the acetabulum



on the iliac column. By cutting the bone loose from the pelvic cellular tissue, it is easily pulled away.

The lateral wall of the pelvis being removed and a good view of the organs obtained, divide the connective tissue between the rectum and the superior pelvic wall; free the uterus and ovaries, the neck of the bladder, the vagina, and the accessory sexual glands; cut through the strong rectococcygei and the skin between the tail and the anus; and make a circular incision around the anus and the vulva (or the region of the penis). Remove the whole mass and section the organs dorsally.

**EXENTERATION OF THE CRANIAL CAVITY.**—To remove the head from the trunk we may either cut around the joint as if the throat were being cut or puncture the capsule ventrally and amputate between the condyles and the atlas. It is best to remove the whole of the lower jaw and let the skull, wrapped in a cloth, rest on its base and the molar teeth; it may then be held much more steadily than if the inferior maxilla had been left in place. The cranial attachments of the cervical and temporal muscles are next cut away and the soft parts removed from the roof of the skull.

There are three lines for sectioning the cranium. The first lies transversely across the forehead about a thumb's breadth above the upper border of both superciliary ridges. The two other lines begin at the ends of the frontal incision, pass backward across the temples and petrous bones, and converge to the condyloid apophyses (Figs. 178, 179). The first section can be made continuously, but the second and third will have to be done in several portions on account of the convexity of the cranium.

The walls of the cranial vault are not equally thick, and care must be taken not to penetrate too deeply into the middle of the parietal bones and the squamous portion of the temporal bones. The frontal section passes through the frontal sinuses, so that there is very little danger here; and the same is true of the vertex and the pyramidal region above the condyles. The plates are not usually sawed clear through along the whole line, but the connections are broken with a chisel. Rest the palm of the hand upon the skull, grasp the chisel firmly near its edge so that it cannot enter too deeply, and tap gently with the hammer. When the bones are completely severed, pry the piece off by rocking the chisel backward and forward, first in the frontal and then in the condylar region. A sudden strong pull on the

pericranium, grasping it at the edge of the frontal section, will generally separate it from the other parts of the head; sometimes the



FIG. 178.—Lines to guide the saw in opening the cephalic cavities of a horse.

whole brain will come away at one jerk, together with the root of the skull.

If the dura is too closely held or is adherent to the inner table of



FIG. 179.—Lines of sawing for opening the cranial cavity of a horse.

the skull, with the shears incise it in the line of the section in such a manner that the dorsal portion will come away with the calvarium.

Next excise the longitudinal and transverse blood-vessels in the duras. That part of the dura lying over the hemispheres is held up with forceps and cut with scissors so that it may be thrown back on both sides. The tentorium is sectioned anteriorly and posteriorly and removed. The membranous transverse septum which is torn away from the falx is incised laterally and pulled out from the transverse fissure, due attention being paid to its vascularity.

**DISSECTION OF THE BRAIN.**—After examining the pia mater and the superficial surface of the brain, the hemispheres should be separated so as to expose the corpus callosum. The interior of the brain may either be examined now or after its removal. A horizontal incision is made immediately over the corpus callosum, starting at the median surface, and using preferably the so-called “brain-knife” or a long, flat scalpel. If the incision is not quite deep enough to enter the lateral ventricle, you will come first to the so-called “egg-shaped middle point” (*centrum semiovale Vieussenii*); press this gently with the finger and you will find a yielding point which, when incised, opens into the lateral ventricle. Follow the finger with the knife and slit open the roof anteriorly and posteriorly. Look for a collection of fluid, and examine the choroid plexus, corpora striata, horns of the ventricle, and median septum. This is seized in the middle, raised a little, sectioned transversely, and thrown back, the connections holding it to the peduncles being severed. Now carefully insert four fingers into the transverse fissure and raise the posterior lobes in order to expose the corpora quadrigemina, optic thalami, pineal gland, and middle choroid plexus. By separating the two thalami a little, you can divide the commissura mollis and see into the third ventricle.

To remove the brain, support the skull upon the incisors in such a position that the condyles look upward and the brain would fall out if it were free. Into the space thus obtained between the medulla and the base of the skull, insert a finger, the closed scissors, or the handle of a scalpel, and sever the nerves one by one as they appear. The olfactory bulbs, which are unusually large in comparison with those of man, are worked out from the ethmoidal depressions by a circular thrusting motion of the handle of the scalpel. When they are all separated, the brain will fall into the waiting hand, which must steady it constantly or the olfactory bulbs would be torn away by its falling out too soon.

After the brain is removed, the inferior surface is first exam-



then, turning the brain over, the cerebellum is cut into halves. Expose the fourth ventricle and incise the floor longitudinally. With a thin-bladed knife cut radially to the cortex and transversely to the cruræ, making numerous narrow incisions to detect the presence of any small hemorrhage or other lesion.

**REMOVAL OF THE SPINAL CORD.**—This requires much time and labor when properly done, but is managed in various ways. But little time is spent in routine work when you have a butcher to assist you. The animal is suspended and the vertebræ are split off from their bodies by a hatchet; when this is cleverly done, the line of cleavage being kept a little to one side, the cord is but slightly injured. It is better, however, to proceed as follows: Saw off the ribs at their angles, separate the ilium from the sacrum, and clean off all the flesh. Laying the spine upon the table, begin at the pelvis and chisel off the vertebral arches, remembering that two chisels are necessary, one for each side, as the two instruments have different curves (Fig. 37). If an ordinary chisel is used, the arches should be partially sawed through to make their division easier. The hand holding the chisel supports itself on the spine, and the chisel is held as flat as possible while an assistant grasps the spinous processes and springs the arch apart. You may also expose the spinal canal ventrally by sawing through the vertebral bodies and arches on one side only. Section the nerves at their points of exit laterally to the intervertebral ganglia and lift out the cord enclosed in its membranes. Cut open the dura with the scissors and section the cord transversely with a sharp, thin knife.

**EXPOSURE OF THE ACCESSORY SINUSES.**—To expose the nasal fossæ saw the head in two, after removing the brain, a little to one side of the median line so as not to injure the septum on either side. These fossæ may be sectioned transversely or their walls chiselled away to show the accessory sinuses. Csokor saws through the osseous structure of the nose transversely from the level of the malar or lachrymal bone to the roots of the molars; a section is then made horizontally beginning at the anterior nares and joining at the first section (Fig. 178). On raising this cap you have the maxillary, nasal, and frontal fossæ well exposed.

One or two long bones should be sawed through to judge of the condition of the bone marrow.

**POSTMORTEMS ON RUMINANTS.**—There are certain peculiarities of ruminants which must be remembered when exposing

the cranial cavity. It is only in very young animals that the cranial bones possess diploe, and in necropsies on hornless cattle the incisions are the same as for horses. On account of the prominent crests, which fall away very abruptly, and because a calf's head is somewhat rounder,



FIG. 180.—Lines used in sawing in order to expose the cranial and nasal cavities in a ruminant.

the sawing will have to be done in more segments, and great pains must be taken on account of the thinness of the bones. The older the animal the larger are the hollow places between the internal and external plates; the diploe disappears and only a few crusts and plates




FIG. 181.—Appearance of cranial cavity of a cow after removal of the bony vault.

of bone interrupt the hollow spaces. The lateral and posterior portions of the skull are very prominent because of two large crests. The transverse section is nearly coincident with the posterior border of the superciliary ridges. The lateral sections are made in two segments,



beginning at the ends of the transverse frontal incision and passing back over the temples to the foramen magnum. Clement has devised a better method (Figs. 180 and 181). First clear away all that part of the calvarium formed by the frontal eminence and the lateral depressions by sawing through the skull in a line passing from just in front of the horns obliquely backward and downward to the condyles or foramen magnum. After removing this plate of bone the whole of the posterior portion of the brain is exposed. Next make a transverse incision on a level with the superciliary ridges across the anterior end of the cranial cavity. Finally make two short longitudinal incisions, one on each side, about three centimetres from the median line; with mallet and chisel remove the oblong piece enclosed, and the whole brain is exposed. The curved horns of a sheep or a goat serve as convenient handles for removing the calvarium and may very well be left on, while the horns of neat cattle should be knocked off.

POSTMORTEMS ON SWINE.—With the body lying on its left side, the right extremities are removed, the abdomen is exposed by longitudinal and transverse incisions, the diaphragm observed, and the lateral thoracic wall divided by cutting with the bone-shears or sawing through the angles of the ribs and severing the cartilages close to the sternum. To remove the abdominal contents, first find where the duodenum is attached to the rectum; sever the duodenorectal ligament, separate the pancreas from the mesentery, and section the duodenum. The anterior root of the mesentery is loosened by working it free with the hand and pulling on it, then sectioned, the whole of the mesentery excised from before backward, and the rectum divided. Now cut away the spleen from the stomach, examine the opening of the bile-duct, section it and the œsophagus, and separate the stomach from the diaphragm, leaving the liver freed from its suspensory ligament. The thoracic and cervical organs are removed as with other animals.

In old quadrupeds the brain lies very deep, because of the immense air-spaces in the cranial bones which surround the brain on all sides except the temporal region. The transverse section is made a full thumb's breadth above the superciliary ridges (the eyes being first removed) and the lateral sections run back to the occipital foramen. Instead of a transverse section we may make two oblique ones, beginning at the posterior border of the frontal process and joining each other and the lateral incision in the frontal region.



POSTMORTEMS ON DOGS AND CATS.—The necropsy of a dog is easily and quickly made in either the dorsal or the left lateral position. The procedure is the same as for the horse, but it is not necessary to remove the extremities entirely or to take off the hide; simply cut through the muscles enough to allow the limbs to fall away a little and the body will be sufficiently steady. (Figs. 182, 183.) The thick-




FIG. 182.—Postmortem of the dog. Double lines show places at which the intestines are to be tied; the dotted line indicates the direction for incising the mesentery.

ening at the junction of the cartilages with the ribs is easily felt, the articulations are cut, and the sternum is pushed upward and forward after freeing the pericardium and the pleura. Section the tracheal

vessels and œsophagus at their entrance into the thorax and remove the thoracic organs.

The removal of the abdominal contents of a dog is easy. Divide the rectum at the pelvis and the two mesenteric roots, and the abdominal aorta and inferior vena cava behind the liver; thrust the hand in between the liver and the diaphragm, and with scissors section the suspensory ligament of the liver, the vena cava, and the œsophagus after it is pulled down from the diaphragm and tied off or compressed with the fingers. All the abdominal contents may now be removed together. Spread them out, examine each again, test the patency of the bile-ducts, and straighten out the bowels. It is, however, better first to remove the intestine, which is sectioned through the duodenum



FIG. 183.—The left ramus of the mandible has been removed and the tongue pulled outward and downward, thus exposing the oral and pharyngeal cavities in a dog.

at the pancreas and through the rectum at the pelvis. You may next either remove the liver with the stomach, or after inspecting the bile-ducts you may cut away the stomach from the œsophagus and duodenum and then remove the liver.

To expose the cranial cavity we have the same three lines as usual, the transverse section lying directly posterior to the rudimentary superciliary ridge, crossing the frontal sinuses and the anterior lobes of the brain. The anterior temporal and the parietal bones are not thick and contain diploe, so that the sawing must be carefully done. Since the petrous portion of the temporal bone has deep impressions upon its internal surface, in which convolutions of the cerebellum lie, and since

the bony processes project from the adjacent bones, great care must be taken not to tear the cerebellum. In small dogs with round heads the line for sectioning is more nearly a circular one.

The postmortem of a cat is made in the same way.

**POST-MORTEM EXAMINATION OF BIRDS.**—Plug up the nostrils, mouth, and vent with cotton; make an incision from the point of the breast-bone, or a little above, backward to and through the anterior portion of the anus, leaving the uropygium (pope's nose). Loosen each leg to the knee (above the femur) by tearing the soft parts with your thumb and fingers, then cut with a knife until they meet around the pelvis at the rump. With your thumb-nail work the wings loose, hold the skin firmly, and, pressing your nail towards the body, cut off the wings at the elbow. To get at the brain make a V-shaped slit with its apex towards the median line at the foramen magnum, running up towards the centre of the skull; the brain is then removed attached to the cord, and the skin is kept whole for taxidermic preservation. The "wish-bone" in birds is the joined clavicles.

**POST-MORTEM RECORDS.**—Kitt suggests the following scheme for the more intelligible recording of the findings in postmortems on the lower animals.

#### RECORD OF NECROPSY.

Species	Gender	Age	Color of hair	Owner
Clinical history	Treatment	Mode of death	Date of death	
Necropsy performed by		Where performed	Date	
Order of		Persons present		

#### A.—EXTERNAL EXAMINATION.

Position of the cadaver (on back, right or left side, hanging)  
 Nutritional condition (weight)  
 Removal or absence of parts  
 Rigor mortis  
 Condition of the skin and its appendages (the skin around the head, trunk, and extremities: the horns, claws, hoofs, ears, scrotum, prepuce, udder)  
 The natural body openings and visible mucous membranes (the discharge of foam, fluids, and excrementa; the color of the lips, nasal mucous membranes, conjunctivæ, anal and vaginal mucosæ)

#### B.—INTERNAL EXAMINATION.

Facts obtained in removing the hide  
 Condition of subcutaneous tissues, fat, lymph-nodules, vessels, extravasated blood, muscles, ligaments, tendons, fascia, joints, and bones  
 Abdominal and thoracic data  
 Condition of diaphragm, position of organs, appearance of peritoneum, mediastinal and costal pleuræ, and pericardium



The oral cavity, tongue, teeth, soft and hard palate, salivary glands, pharynx, Eustachian tubes, œsophagus, retropharyngeal and laryngeal lymph-nodules  
 The larynx, trachea, thyroid, and surroundings  
 The lungs, bronchi, bronchial lymph-nodes  
 The pericardial sac, heart, and thoracic vessels  
 The liver and bile-ducts, portal vein, and periportal lymph-nodules  
 The spleen (capsule, pulp, trabeculæ, Malpighian bodies, and vessels)  
 The stomach and crop  
 The pancreas; the large and small intestines  
 The mesentery, omentum, posterior aorta and its branches, and vena cava  
 The kidneys, adrenals, ureters, capsule and pelvis of the kidney, and its half section  
 The urinary bladder, urethra, and accessory sexual glands  
 The pelvic portion of the rectum  
 The genitalia: uterus, vagina (pregnancy, foetal membranes, embryo), and the male genitals  
 The cranial cavity and the brain: calvarium, sinuses, cavities at the base of the skull, dura, cerebral superficies, ventricles, gray and white matter  
 The eyes; the middle and internal ears  
 The fourth ventricle and the spinal cord with its membranes  
 The nasal fossæ and accessory sinuses  
 The udder and supramammary lymphatic nodules  
 The bone marrow  
 Microscopic and chemical report

**INSPECTION OF SPECIAL ORGANS.**—The essentials for diagnosis which are to be looked for and recorded are about as follows: 1. Name of the organ; from what animal; whether it died or was killed; whether the organ was entire or fragmented; whether parts, lobes, etc., have been amputated; and if there are any adhesions to adjacent parts. 2. Weight. 3. Length and breadth of the part. In the absence of a tape measure we may ascertain these dimensions approximately by comparison with the breadth of the hand and the length of the finger. Every person should know the length of his index-finger, which is usually about ten centimetres and may be used to measure organs, pathologic spots, streaks, canals, etc. 4. Surfaces: whether smooth, even, wavy, granular, corrugated, rough, transparent, or cloudy. Color of the surface: general and primary color, special deviations and shades. The external contour of the organ and any prominences, with especial reference to their size as compared with grains of sand, millet-seeds, lentils, peas, beans, hazel-nuts (or filberts), a pigeon's, a hen's, or a goose's egg, the fist, the thickness of a child's arm, a child's head, etc. (Plate II.) 5. The consistence as determined by palpation; soft, elastic (like the lungs), doughy, splenified, hepatized, tough, inelastic, carnified, indurated, leathery, like the kid-

neys and skin, as hard as wood, cartilage, bone, or stone. 6. Sectioning of special parts: through the compact, so-called parenchymatous organs (muscles, liver, kidneys, lungs) large dissecting incisions are made. Through the brain and heart sections must be made in a certain way in order properly to expose certain cavities. On sectioning notice the resistance of the tissue, whether it cuts easily or is tough and pulls, whether the knife creaks as it goes through, whether the tissue is so hard that a saw is necessary, and observe if any fluid follows the section or if there are any abnormal contents. The surfaces of the section must be noted, their color, thickness, consistence, fluidity, and vascularity, as well as any other peculiarities which may be present. The pathologic diagnosis is made by considering the details gained in this way, which lead to one conclusion and exclude another. A gross anatomic diagnosis is often only provisional and dependent upon microscopic and chemical confirmation.

## CHAPTER XXVI

### MEDICOLEGAL SUGGESTIONS

ALTHOUGH a physician is not expected to have a profound knowledge of legal matters pertaining to his profession, yet every doctor should be more or less familiar with the medical laws of the State or country in which he is practising. He should be well acquainted with the regulations of the board of health, of the coroner's office, of the criminal court, etc., and do all in his power to aid in their rigid enforcement. A synopsis of such laws and regulations is usually readily obtainable in book form, and nearly every physician has among his patients or friends a lawyer who is glad to discuss legal questions in return for medical information. Some of the salient points relating to medicolegal investigations and autopsies will here be briefly considered, though many references to these matters will be found elsewhere throughout this work, especially in Chapter XXVII.

OBLIGATIONS OF PHYSICIANS TO THEIR PATIENTS.—The obligation of a physician to society in the practice of medicine is in a certain sense voluntary. His is the right to refuse any and all cases that may apply to him for treatment or advice. Services once begun, however, he must, after giving notice of his intention to discontinue them, allow his patient reasonable time to fill his place, as otherwise he renders himself liable for damages. This obligation is equally binding in the case of charity patients. Contracts between a physician and a patient may be either express or implied. An express contract is where services are rendered in accordance with a definite agreement previously entered into between the parties. There is the promise of proper treatment in an implied contract just as there is the promise of payment. Generally, however, the contract is implied by the law from the fact of employment and consequent attendance. Both forms of contract are equally binding, and both are subject to public policy. Contracts making the payment contingent upon successful treatment are valid, but, should the patient fail to follow the doctor's directions or to give him sufficient opportunity for treatment, the Court would probably allow the latter reasonable compensation. If the physician fail to exercise ordinary skill, he renders himself liable for malpractice.



In law malpractice consists in the failure to possess or use such ordinary knowledge or skill as is generally possessed and used by physicians and surgeons in similar localities, whereby injury accrues to the patient. The term is also applied to acts which are expressly forbidden by statute. The average physician is not bound to possess the highest expert skill nor is his implied contract one binding him to effect a cure; he must simply treat the case with reasonable skill, diligence, and faithfulness. That done, he cannot be held accountable for results. Births, deaths (with their probable cause), and infectious diseases are to be reported to the proper authorities for registration. In hiring a wet-nurse for a syphilitic child the woman must be informed of the fact that the infant is specifically infected and of the risk that she runs in taking it to nurse. It is a criminal offence to practise medicine or surgery while intoxicated.

The obligations of a patient to his physician, in so far as they relate to the treatment given, are indefinite and more or less vague. They can hardly be considered to come within the province of exact definition. The patient should conform to the directions given him, but no legal liability arises if he does not: the risk is his. If in such circumstances the attending physician is of the opinion that his advice is being disregarded, to the detriment of his patient and perhaps of his own reputation, he is always at liberty to withdraw from the case and to require that another physician be called in.

As to compensation, it may be laid down as a rule that in the absence of a special contract the measure of the liability of the patient to his physician is the customary charge made by others of equal standing for similar services. The question of compensation is a broad one and its full treatment is beyond the scope of this work. It may be remarked, however, that the specialist should protect himself by an express contract. Physicians should also be careful to have their accounts so kept as to be able to use them as evidence if dispute should arise. Much trouble and loss are frequently caused by neglect in the entries made by physicians in their books. In the case of *Laffin v. Billington*, the Appellate Term of the Supreme Court of New York held that a contract by a physician to give expert testimony for the plaintiff in a personal-injury case and to receive as his pay therefor a percentage of any amount realized is invalid.

EXPERT TESTIMONY.—Applying these principles pertaining to medical practice to our subject, no Court can compel a physician to

give expert testimony, to make autopsies, or to conduct laboratory investigations without his consent, but any knowledge which the doctor may possess pertaining to an individual criminal case must be given to the Court in the same manner as if he were an ordinary witness. His scientific training is, however, his own personal property, the result of many years' study, careful research, and expenditure of money, and he is entitled to commensurate remuneration for the expert use of his knowledge. For the good of society, any facts pertaining to a given criminal case which are known to him should be freely and willingly given to the Court, though he thereby may be put to considerable loss of time and money. The expert should be cautious in expressing opinions before the case is called for trial. (Wormley.) Thus, in the Williams case, tried in Philadelphia in 1903, I was asked by the district attorney while on the witness stand if I had ever expressed to the attorney for the defence an opinion as to the cause of death, and, if so, what that opinion was.

Whether the Court will compel him to divulge professional secrets is a debatable question. In some States and countries such confidences of the patient are held sacred, as are the relations between counsel and client; while in other places such confidences (wrongly, we believe, in civil cases, but rightly in first-degree criminal cases) must be divulged to the Court should questions pertaining to the same be asked the physician while on the witness stand. The matter is one for the Court to decide, and such decision being given absolves the witness. Whether the prescription of a physician is a privileged communication is a matter of much debate, the weight of opinion, probably, being that it is not.

An expert is one who by reason of his peculiar experience, special study and training, or the performance of certain duties, is competent to ascertain particular facts of a technical nature or to form an opinion or judgment upon them, such as could not be expected from the judge or jury. No ordinary witness is permitted to express an opinion upon the facts as presented to him, as this is supposed to be the province of the members of the jury. Thus, in one of my cases, where infanticide was suspected, an iceman had found the dead body of the baby in an ash-barrel, and the judge would not permit the iceman to act as an expert in giving the approximate weight of the child, though it would seem that, on account of his frequent weighing of ice, he would be more fitted to give a correct estimate of the weight than an ordinary

person. The weight of the child (nine pounds) was desired in order to show that it was born at or near full term.

The jury is supposed to arrive at a decision as to the point at issue from the facts proved before it, and hence must necessarily form an opinion thereon. Where, however, there arise in the case technical issues, matters involving special knowledge, as to which persons in general are not qualified to reach a correct judgment, and where, therefore, the opinion of those versed in such knowledge is necessary to the formation of a proper verdict,—in such case experts are allowed to testify. In so doing they must frequently give their judgment as to the facts, often presented to them in the form of hypothetical questions. Such questions are presumed to be framed from the testimony already accepted. Great care should then be exercised in testifying, especially as sometimes the ingenuity of counsel is used so to formulate the hypothetical question as to confuse the issues in the mind of the jury or to extract from the witness evidence contrary to the true state of the case.

A medical man should refuse to testify as an expert unless he is thoroughly qualified. In no case should he go on the witness stand without being as fully informed as is possible on the subjects on which he is to be examined, nor should he allow himself to be questioned on subjects on which he is not prepared. He should be honest and candid with those securing his services before the trial, and, no matter what may be the consequences, his answers while on the witness stand must be made with absolute impartiality. Upon a suit for damages an expert may be held responsible for errors which he may have committed in the performance of his work. Thus, a chemist passing glucose as free from arsenic might have to pay damages, should beer be made out of the glucose and arsenical poisoning result therefrom.

The medical expert should at all times confine himself to purely medical topics and never become involved with matters that will place him in the light of an ordinary witness, of a detective, or of an attorney, and he should carefully avoid acting as a champion of the parties who are paying for his services or attempting to plead one side of the case.

His language should be as free as possible from technicalities and such as can readily be followed by the least educated of the twelve jurymen, many of whom are only too often, unfortunately, unfit for the performance of their duties. Some judges carry this plainness of lan-



guage to an extreme. Thus, while acting as an expert in a murder trial, the writer was once requested by the judge not to use the word "hemorrhage" in testifying, as this term was too technical for the jury to understand. I at once substituted "bleeding" for the objectionable word and proceeded with my testimony. When not absolutely certain of a point he should unhesitatingly acknowledge the fact; thus harm and the possible endangering of a human life will be avoided. But when sure of his ground he should undeviatingly adhere to it. At the close of his testimony, especially if long and exacting, an opportunity is almost always given him to correct any misstatements which he may inadvertently have made, and to make clear the meaning of any dubious points of his original testimony which may have been clouded by the cross-examination conducted by the opposing counsel.

Too much is often expected from the expert, as the following instance shows. While testifying as an expert in a country town on a case where the postmortem revealed beneath the left eye a small incision closed with two stitches, ecchymoses about the eyeball, and a fracture of the skull, the district attorney and the judge criticised me severely because I would only state that the man had died from hemorrhage of the brain due to fracture of the skull. They desired me to say that the man had been knocked down with the fist of a person who had a ring upon his ring-finger, and that in this manner the cut and the fracture had been produced. I was naturally willing to say that they could have been made in this way, but would not state, apparently much to their disappointment, that they were so caused.

MEDICOLEGAL POSTMORTEMS.—The objects of a medicolegal postmortem include the finding out of the cause and mode of death, the establishment of a *corpus delicti*, the determination as to whether a crime has been committed, and if so the discovery of a motive therefor and the exact nature of the process employed therein. In such legal investigations the pathologist should protect his reputation in every possible manner, and he ought to hesitate to make a postmortem without the presence of a witness, who should, if practicable, be a professional brother. The autopsic findings should be dictated at once to an amanuensis, and the record verified and signed upon its completion.

The obducent should have an opportunity to view and examine the body before it is taken away from the place or position in which it is found, and especially before the clothes are removed. He should prior to the autopsy be put in possession of information as to the

general, and also any special, circumstances of the case, and more particularly in regard to any injury or violence which the deceased may have received. He should also be advised as to any known disease or other condition which may have contributed to the death. No one ought to be allowed to witness the examination out of mere curiosity or unless specially authorized. The examination should not be commenced unless there is sufficient daylight in prospect to allow of the whole inspection being made without artificial light. All measurements should be accurately determined. Examine carefully the contents of the stomach to ascertain the length of time which has elapsed since the death occurred. Where the services of the obducent are called in by the law, he is freed from the necessity which exists in ordinary cases of obtaining the consent of the relatives of the decedent.

In general the medicolegal post-mortem examination does not differ materially from the pathologic, except that in the former greater precautions are necessary in order to avoid sources of error or confusion, and that the cranial contents are examined before opening the large blood-vessels, as signs of congestion disappear after the severance of the aorta and *venæ cavæ*. Doubly ligaturing the *œsophagus* at the left of the trachea is a good routine practice, and it should always be done in cases of suspected poisoning. The urine collected from a cadaver is practically always albuminous. The importance of examining the *vertebræ* in all autopsies cannot be too strongly insisted upon, as severe injuries thereof may exhibit no external signs of violence while there may be other lesions found on the body which might otherwise be erroneously assigned as the cause of death. In some cases after a most rigid and painstaking inspection no cause of death can be ascertained, but with care and systematic examination mistakes and inaccuracies will be reduced to a minimum.

In case of suspected poisoning the *primæ viæ* should be tied at each end and removed. Double ligatures should then be applied at the junction of the duodenum and the ileum and at the end of the small intestine, dividing the viscera into three portions. The contents of the stomach and those of the intestines should be emptied into separate jars. Many poisons are extremely volatile and without great care traces of them may be lost and justice defeated.

Each organ should be received in a separate receptacle, and each receptacle should be marked, sealed, dated, and deposited where tampering with it would be impossible. The mouth of the receptacle

should be so large that no injury will be done to the organ in its introduction. It is also well to remember that a tissue in its fresh state goes into a bottle more readily than it comes out after being hardened by the preservative fluid.

The form of report used by the writer in medicolegal cases is as follows: "I made a post-mortem examination of the body of Walter Foster on April 10, 1898, at St. Agnes Hospital, Philadelphia. The body was identified by George Bell, 636 Siegel Street, and Michael A. Bruder, 1847 Sartain Street, both of Philadelphia. I find that death was caused by shock and hemorrhage from stab-wound of the heart."

While acting as coroner's physician I rarely volunteered more than this, but waited for the district attorney to ask questions in regard to the nature of the wound and as to other facts of interest. By this method the jury is not confused by an enormous amount of irrelevant testimony, though the expert must be prepared to give, under cross-examination by counsel for the defence, the minutest details as to how the postmortem was performed.

It may be remarked that there is a growing tendency to perfect and render more scientific the proceedings incident to and growing out of violent deaths. Courts have even appointed a commission of physicians to examine into the condition of the plaintiff in personal injury cases. It has been suggested that in postmortems involving the question of crime the investigation should be carried on jointly by experts representing the State and the defendant; also that the question as to the cause of death shall be disassociated from that of the guilt or innocence of any particular individual and determined by a jury or commission of experts; but this brings us into the realm of speculation. We do not now confront such conditions.

**AUTOPSIES ON INFANTS.**—The first question to determine in the examination of a babe is, was it born alive? If so, was it a full-term or a premature birth? If born dead, how many months of uterine gestation caused it to reach its present development, and after attaining its maximum growth was it carried as a foreign body in the uterus?

**DETERMINATION OF THE VIABILITY OF A CHILD FROM THE POST-MORTEM APPEARANCES.**—The reader is advised carefully to read Paragraphs 23 and 24 of Virchow's regulations for the performance of medicolegal postmortems. (See Chapter XXVII.) To discover the ductus arteriosus remove the thymus gland, incise the right ventricle along its septum, and extend the incision into the pulmonary



artery along the middle portion of its anterior wall. The orifice is situated between and beyond the two openings of the right and left pulmonary branches. If the duct is open, a sound will readily pass into the aorta. It should be remembered that decomposition may produce bullæ in the lungs, that the lungs may be distended by the forcible introduction of air in the methods used for artificial resuscitation, and that air may get into the lungs of the child from emphysematous conditions affecting the uterus of the mother.

If in the hydrostatic test the lungs float on top of the water, they have been completely aerated, a strong proof of breathing at or after birth; if they float beneath the surface, aeration is incomplete; and if they sink, no respiration has occurred. Decomposition of the lung tissue may cause it to float. A very valuable sign of the viability of the child is the presence of uric acid crystals in the kidneys.

Rigor mortis does not prove, as has sometimes been asserted, that the infant was born alive in the legal acceptance of this phrase. The rigidity may be of the so-called ante-natal variety; it may even unduly prolong labor by interfering with delivery.<sup>1</sup>

The *Lancet* of April 26, 1902, raises the query whether the dead body does not possess properties akin to radio-activity, and alludes to the photographs taken by Vignon and exhibited by him, with the winding sheet preserved at Turin and traditionally said to be that of Christ, which seem to justify the belief that the human body is either radioactive or that it gives off vapors which exhibit a similar action to light upon sensitive surfaces. Peroxide of hydrogen may be the main factor concerned. *N*-rays of short wave length can be reflected, polarized, and refracted, and are transmittible by a metal wire. Ballet has found the emission of these rays to be decreased in diseases of the cord, as myopathies and neuritis, and increased in diseases of the brain, as hemiplegias and spastic paraplegias. Certain substances seem to store up these rays and emit them later. The question is whether there are not various rays of a similar nature. The conversion of radium into helium may also be of importance in this connection. Fluorescence and phosphorescence have also been much studied of late, and all these phenomena may have an important bearing on the future of pathology. An ion contains electrons. The *N*-rays increase the brightness of a spark, and phosphorescent bacteria may be used for detecting them.

---

<sup>1</sup> *Lancet*, February 14, 1903, p. 460.

Sight, taste, smell, and hearing become more acute under their influence. Chloroform applied to plants seems to hinder their production. The *Comptes-rendus hebdomadaires des séances de l'Académie des sciences*, 1903-4, contains most of the best literature on the N-rays.

PERIOD OF INTRA-UTERINE GESTATION.—In deciding the age or period of development of the infant the external evidences of value are: (1) Length and weight of the child (for tables of dimensions and weights of the new-born see page 358). (2) Conditions of the skin and its appendages. In the healthy babe at full term the skin is white and covers the body smoothly; woolly hairs are present in perceptible numbers only on the shoulders; the hair of the head is from two to three centimetres long; the nails are hard and horny, extending beyond the ends of the fingers, but not of the toes. (3) Condition of the umbilical cord, which at term is fifty centimetres in length and is inserted somewhat below the middle of the abdomen, falling off by inflammatory demarcation on the fifth or sixth day. (4) State of the cartilages of the nose and ear, being hard in the mature infant. (5) Presence or absence of the membrana pupillaris, which disappears after the eighth month. (6) Condition of the genitals in both sexes; as descent of the testes begins at the seventh month, those of the full-term male should be in the scrotum. The female labia are generally found closed. (7) The measurement of the fontanelles, of the cranium, and of the transverse diameter of the body at the shoulders and hips. (8) The size of the centre of ossification (Béclard's) in the lower epiphysis of the femur. To view this the leg is flexed on the thigh, a transverse incision is made below the patella, which is removed, and the femur is then exposed. Thin, transverse sections of the cartilage are made until the greatest diameter of the centre of ossification, if present, is reached. The centre is absent before the thirty-seventh week, and in the child at full term has a diameter of from two to three lines, though it may even then be absent. If the diameter is more than three lines, the child has very likely lived for a time since its birth. (Reese.) The osteochondral line is also to be examined for syphilitic changes. (9) With but very rare exceptions, a full-term child presents in the inferior maxilla eight alveolar compartments completely separated the one from the others. (Vibert.)

From the internal examination important evidence as to the age of the child, and especially as to respiration, is secured. Upon ex-

posing the abdominal cavity, which is to be done before opening the thorax or cranium, the position of the diaphragm in its relation to the ribs is immediately noted, as especially urged by Virchow. If the lungs do not contain air or are but partially distended, the diaphragm reaches to the fourth rib; when the lungs are fully distended, the diaphragm is at the fifth or sixth rib on the right and at the sixth rib or intercostal space on the left.

To facilitate the examination of the umbilical vessels, Nauwerck recommends a division of the usual abdominal incision, shortly before reaching the navel, into two diverging incisions extending to the pubes. The abdomen is opened, and the umbilical vein, made prominent by traction on the triangular flap, is traced along its course, ligated, opened with small scissors, and divided. Turning down the flap over the pubes exposes for examination the umbilical arteries to either side of the remains of the urachus. (Fig. 154.)

**CRIMINAL ABORTION.**—Formerly abortion was not legally a crime if performed with the consent of the mother prior to the viability of the fœtus. It was at one time not regarded as murder even to take the life of a child at any period of uterine gestation. The barbarousness and danger to society of this view were early recognized, both abroad and in this country, and various laws with different penalties attached thereto were enacted making it a criminal offence to practise abortion at any period of gestation, unless for the express purpose of saving life. (Witthaus and Becker.)

There is no other class of cases so trying to the patience, ingenuity, and skill of the pathologist as those of abortion, which is accomplished by numerous methods. Many respectable women expose themselves to cold, falls, and douches with the hope of relieving themselves of their offspring apparently by accident. Many pills and potions are sold to induce a resumption of the menstrual discharge, and one often finds them on sale in drug-stores of the first rank and openly advertised even in the religious press. These nostrums are sometimes composed of poisons that may cause the death of the mother. The use of instruments, especially the spiral douche advertised in so many papers, is a very common method of procedure. Indeed, the most successful criminal abortionists operate so that, unless through accident, no evidence of the operation is left. Usually all that can be found is evidence of a recent pregnancy. The five most frequent causes of spontaneous abortion are syphilis, alco-



holism, the infectious fevers, endometritis, and diseases of the placenta, as apoplexy. It must be remembered that, while there are generally accomplices in abortion, yet in many cases it is self-inflicted. The difficulty of distinguishing between the two is almost insuperable. All that the obducent can do is to use the utmost care in his examination and to note all the circumstances, with a view not only to the conviction of the guilty abettors, but also to clear, if possible, those unjustly accused.

When violence is done to the child, the nature of the injuries must be carefully noted. When violence is done to the uterus, some form of infection usually follows. In examination care must be taken to exclude the possibility of previous disease of the uterus or adnexa as a cause of the infection or possibly as a cause of abortion. In former days, when curettage was more used than it is now in the treatment of abortion, an additional factor was added, making it difficult and often impossible to distinguish dilatations of the os due to the instrument producing the abortion and to the passage of the foetus from those induced by the introduction of the curette and the subsequent packing with iodoform gauze. The vital history of the foetus should be compared with the physical condition of the mother, the history of the sexual life of the parents, specific disease, etc.

**USUAL CAUSES OF DEATH.**—In Chapter XXVIII will be found a list of all the recognized causes of death, and it is recommended that this classification be used by every one in order that uniformity of nomenclature may be secured throughout the world. Sudden death is usually due to failure of the circulatory apparatus, to cessation of respiration, to disturbance of the nervous system, to deficient nutrition, to poisons either produced within the body or introduced from without, or to violence by physical or chemical forces, heat or cold, electricity, wounds, missiles, etc.

Many conditions that have existed a long time may cause sudden death by breaking the balance of life. Thus, in chronic nephritis uræmia may develop suddenly and cause death after only a very slight acute illness. Again, an aneurism may rupture without sudden increase in the symptoms or any violence, simply by a natural slow progress of the lesion. All mortal diseases and many that by themselves do not end fatally may contribute to the causing of sudden death as well as to slower dissolution.

No disease causing severe disturbance of heart, kidney, lung,

nerve, or digestion can be ignored in estimating the factors that brought about the death of the patient. Certain maladies of common occurrence should be in our minds in making examinations, though never so prominently as to prevent a proper search for other conditions. Thus, in children think of pneumonia, enteritis, bronchitis, meningitis, congenital syphilis and other hereditary diseases, infectious fevers, malformations, etc.; in young adults, infections, local and general, violence, typhoid fever, and tuberculosis; in middle life, diseases of the lungs, kidneys, heart, and blood-vessels, hepatic and gastro-intestinal conditions, infections, violence, occupation neuroses, pneumonia, tuberculosis, cancer, etc.; in old age, nephritis, carcinoma, sarcoma, aneurism, cerebral hemorrhage, embolus, thrombosis, tumor or abscess, arteriosclerosis and obstruction of the coronary arteries, heart lesions, etc.

In coroner's cases death very commonly results from heart exhaustion, due, as the case may be, to intrinsic disease, to excitement, or to poisons. Care should be taken to determine the cause of this exhaustion, whether it was due wholly to heart disease, such as a valvular lesion, or to one of the exciting causes. In kidney congestions consider whether death was due to failing heart causing passive congestion, to poisons, or to inflammatory congestion, such as would be part of an acute nephritis. Ascertain if the œdema of the lungs is dependent upon cardiac, renal, or cephalic lesions or primarily upon a lung condition principally.

**VIOLENT DEATH.**—When there is doubt as to homicide, all the precautions necessary for such cases must be strictly observed. The sort of violence, its mode of application, and something of an estimate as to the amount, direction, and conditions of application of force can usually be made from post-mortem examination. In the inspection of wounds the condition of the tissues and the position and direction of all lesions discovered are to be very carefully noted, as sometimes the instrument with which they were inflicted may safely be inferred therefrom, and at times the findings will point to the circumstances under which the injuries were received. A minute description of the injuries is absolutely necessary, so that if called upon in court an exact account of them can be given. The amount of contusion, laceration, extravasation of fluids, and damage to any vessels must be carefully noted. It is important in gunshot wounds that the projectile should be found. About the wound of entrance look for

# PLATE VI




PLATE SHOWING COMPARATIVE EFFECTS OF "BLACK" AND "SMOKELESS" POWDER CARTRIDGES AT SHORT RANGE.

Figs. 1, 2, 3, and 4 are produced by a revolver (.38 inch) charged with ordinary black powder and held in hand. The firing distance is eight inches. The markings were made before the members of the Academy of Surgery of Philadelphia on the evening of May 5, 1902. In Fig. 1 the hammer is up; Fig. 2, to the left; Fig. 3, to the right; Fig. 4, down. Fig. 5, revolver immovably held by fixing in a vice; bullet-hole in centre of brand; black-powder cartridge. Fig. 6, revolver cartridge of "smokeless" powder, fired from hand, hammer up; showing circular powder mark, with little or no "brand"; Fig. 7, same, hammer down. (The illustrations are about one-half natural size.)



1

2

3

powder marks, singeing, and smudge. If the projectile struck a bone, a splinter may have been detached and caused injuries not along the line of the main wound. The effect of small-arms loaded with smokeless powder has recently been studied by Johnson.<sup>1</sup> Such markings are less distinct and more indefinite than in the case of black powder. Even at a distance of three inches or less the markings may be so indistinct as to be capable of being wiped away from the skin with ease, and when the part is covered with clothing no powder marks are found nor is the clothing scorched. Herold<sup>2</sup> reports a fatal case where a pistol fired at short range left no powder markings of any description. Plate VI shows the effects of black and smokeless powder at short range. (Brinton.<sup>3</sup>) Death is frequently due to shock, which may result from a blow that leaves no mark visible at the post-mortem. This is quite uncommon. Injuries to the head make it necessary to estimate the structural and tensile strength of the skull in each case. When a fracture of the skull is found or suspected, the skullcap must be cut away with the saw only, not using the chisel. Contrecoup must always be considered in hunting for fractures and lacerations of blood-vessels. Rawling<sup>4</sup> has recently published an interesting article on the mechanism of skull fractures and Wadsworth<sup>5</sup> takes up the question of injuries to the brain. Accidents of various sorts may produce most marked and varied deformities, and give rise to such interesting medicolegal questions as the following: What was the duration of life after the reception of the fatal injuries? Did an electric shock immediately kill the person, or, being only stunned, did he die from the effects of the fall? Did a woman whose decomposed body was found in water, with enough arsenic in her system to kill, die from the effects of the poison or from drowning? In a case where a man shot his wife and then committed suicide, which one died first?

SUICIDE.—Upton, formerly of the *Chicago Tribune* and an authority upon suicide in America, says that during the last thirteen years, 1891 to 1903 inclusive, 77,617 suicides (57,317 men; 535 physicians) were reported as suicides in the newspapers of this country. That the number of suicides, especially among children, is increasing

<sup>1</sup> *Annals of Surgery*, 1904, May, p. 798, and June, p. 1006.

<sup>2</sup> *Legal Medicine*, 1902.

<sup>3</sup> *Int. Clinics*, Twelfth Series, vol. iii, p. 148.

<sup>4</sup> *Lancet*, 1904, April 9, p. 973, and April 16, p. 1034.

<sup>5</sup> *Proceedings Phila. Path. Soc.*, February 28, 1901.

throughout the world is undoubted, this being more notable in times of financial trouble. Up to the time of the Japanese war, suicide had not increased in Russia. Prior to 1894 the larger number of suicides shot themselves; now such poisons as gas and carbolic acid are most frequently employed, arsenic not being used so often as formerly. In Chicago in 1902 there were 147 cases of self-destruction, 127 of which were by carbolic acid. In Philadelphia during the same year there were 196 cases, of which 42 were from gunshot wound, 33 from gas asphyxiation, 32 from strangulation, and 31 from carbolic acid. According to Vibert, the order of frequency of suicides in France is by hanging, drowning, shooting, illuminating gas, and poison. In Germany a favorite method of committing self-murder in the army is to explode a blank cartridge in a rifle, the barrel being previously filled with water.<sup>1</sup> In a recent suicide in Philadelphia an insane patient at Blockley held a razor in each hand and gashed his throat in both directions. I have seen cases where a man cut his wife's throat and then his own, thus affording a favorable opportunity of comparing the wounds inflicted. Great care must be used in making deductions from such examinations, as well as in saying from the inspection of an incision whether it was made by a right-handed or a left-handed person. To prevent detection ingenious ways of hiding the effects of poisons upon certain parts are often tried. Thus, potassium bichromate may be introduced into the stomach inside of figs, a device which will prevent injury to the upper intestinal tract.

During my term of service of nearly three years as Senior Coroner's Physician of the City of Philadelphia, I performed, according to the official records, 799 postmortems, of which 59 were homicidal. It should be remembered that there were two physicians for Philadelphia, and that the Coroner's jury were able in many cases to render a verdict without a necropsy being made, practically no accident cases, no suicides from known causes, and no subjects accompanied by letters from a physician stating the probable cause of death undergoing such examination. Of persons "found dead" the bodies were too much decomposed to permit of a proper diagnosis, some of them having been in the water for months or showing skeletal parts alone. Under the head of Bright's disease and uræmia are included some cases of alcoholism or of deaths incapable of diagnosis *post mortem* without

<sup>1</sup> *London Times*, July 30, 1903.



extended chemic or microscopic study, etc. All the victims of heat-stroke, with one exception, appeared to be alcoholics.

The chief causes of death, arranged alphabetically, were:

Abortion .....	33	Heart disease .....	76
Abscess of various parts of the body .....	5	Heat .....	9
Alcoholism .....	45	Hemorrhage from various causes .....	
Aneurism .....	24	other than extra-uterine, injuries, .....	
Apoplexy .....	28	etc. ....	16
Appendicitis .....	7	Inanition .....	14
Bright's disease and uræmia .....	84	Injuries, as kick of horse, blows, run .....	
Burns and scalds .....	5	over by wagon, etc. ....	16
Cancer .....	12	Meningitis, tuberculous, and spotted .....	
Childbirth .....	3	fever .....	6
Cholera infantum .....	9	Peritonitis from other causes than .....	
Cholera morbus .....	5	abortion and appendicitis .....	9
Concussion of brain .....	3	Poisoning .....	54
Convulsions .....	7	Scarlet fever .....	4
Croupous pneumonia .....	19	Stab wound .....	8
Cyanosis .....	8	Strangulated hernia .....	4
Diphtheria, including croup .....	11	Strangulation .....	7
Drowning .....	50	Stillborn .....	15
Electric shock .....	8	Suffocation .....	9
Erysipelas .....	2	Syphilis .....	3
Ether narcosis .....	2	Tetanus .....	2
Extra-uterine hemorrhage .....	4	Traumatic hemorrhage of the brain. ....	9
Found dead .....	9	Tuberculosis of the lungs .....	16
Fracture of skull .....	27	Typhoid .....	6
Fractures, other .....	3	Miscellaneous .....	68
Gunshot wounds .....	25	Total .....	799

The fifty-nine cases of homicide were made up as follows:

Burns .....	1	Gunshot .....	18
Drowning .....	2	Knife wound .....	10
Injuries, as hemorrhage from frac- .....		Poisoning (illuminating gas 2 and .....	
ture of the skull from a fall fol- .....		strychnine 1) .....	3
lowing a blow, etc. ....	24	Strangulation .....	1

The kind of poison used in the fifty-four cases was:

Aconite .....	1	Oil of merbane .....	1
Ammonia .....	1	Opium (acute) .....	10
Arsenic .....	5	Opium (chronic) .....	1
Carbolic acid .....	10	Oxalic acid .....	1
Chloroform .....	1	Phosphorus .....	1
Cresosote .....	1	Silver nitrate .....	1
Cyanide of potassium .....	1	Stramonium .....	1
Hydrocyanic acid .....	2	Strychnine .....	2
Illuminating gas .....	12	Sulphuric acid .....	1
Lead .....	1		

Cases of poisoning are almost daily being reported from new sources, by novel methods of procedure, and from a constantly increasing number of unexpected causes. Thus, in flash-light photography the magnesium oxide and the chlorate of potassium may produce on ignition chlorine gas sufficient to inflict bodily harm.<sup>1</sup> A dye, paraphenyldiamin, used in the preparation of certain furs, may give rise to bronchial asthma and skin eruption. The use of thread to remove particles of food from the teeth may cause arsenical poisoning and the picking of the teeth with splinters of matches may give rise to phosphorous necrosis. Children playing with "tin" toys and soldiers in whose flesh are embedded bullets may suffer from lead poisoning. Poisonous toxins are often produced in imperfectly cured fish. The spraying of fruit trees may occasion fatal poisoning by the material falling on vegetables below, such as salads and beet greens. Löffler calls attention to poisoning by milk from cows that have eaten poisonous plants. Petroleum spees, like the old-time chloroform parties, are increasing, boys sometimes stealing kerosene from the public streets for the purpose. Women smoke "tea" cigarettes and boys suck the cologne from automatic machines. The effects of poisons may be modified in many ways; thus, the presence of insoluble substances may inhibit to a certain extent their toxic action.

**BURNS AND SCALDS.**—Burns are produced by dry heat and show when fresh no maceration of the tissues. When inflicted by intense heat or by flame, there will be found scorching or singeing of clothing and hair, and possibly of flesh. When resulting from contact with a hot surface, note especially the shape of the burn, and, if the supposed hot object is to be obtained, a corresponding mark may be found upon it. In burning the hair often reddens. In burns from electricity the markings are apt to be branched. It must not be forgotten that burns and scalds, especially when preceded by an explosion, may cause considerable injury or even destruction of the parts. Scalds are produced by vapor, steam, or a liquid, and usually show some trace of the action of the fluid on the mucous membrane or skin. In plain scalds singeing is absent, but where fire has followed an explosion both scalds and burns may be found. In such cases the mucous membrane of the air-passages should always be examined. In cases of scalds and burns the extent of the injuries must be determined both in breadth and in

<sup>1</sup> GRAEFE, *Deut. med. Wchnschr.*, March 13, 1902, p. 191.

depth, with a careful observation of secondary changes, such as sepsis, internal congestions, and inflammations. There are probably produced by these means hæmolysins and hæmoagglutinins,<sup>1</sup> which products are poisonous to the organism, and act as in other forms of auto-intoxication.

DEATH BY ELECTRICITY.—There are no absolute and constant indications. In some cases the point of entrance or of exit can easily be made out by the change in tissues or in clothes. Frequently there is marked burning of the skin. In many instances the only evidence is an unnatural rigidity of the muscles, sometimes with distortion, due to a coagulation of the muscle substance by the current, which, if found in one part and not in another of the same body, may be of significance. There may be evidence of electrolytic action in the blood and organs, as in the brain and cord. There may be livid areas, even hemorrhages, though after sudden death they are not usual.

The face is sometimes distorted. The heart is usually flaccid, although the left side may be hard or tense. On the right side dark fluid blood is often found distending both auricle and ventricle. The same condition exists in the left auricle, but the ventricle is almost empty. The pupils are invariably widely dilated immediately after death. The blood is usually fluid, but clots have been found in the heart and large veins.

Jellinek<sup>2</sup> finds that the anatomic changes in the tissues resulting from the passage of a powerful electric current diminish the resistance of future currents. Mice are killed with a weak current, but pigs show the greatest resistance. Death by electricity occurs more quickly after administration of morphine or cocaine, but is retarded by chloroform anæsthesia. A dose of morphine might therefore be administered with benefit before an electrocution. Microscopically, degenerations are found in the gray matter of the spinal cord along with dilatation of the central canal and hemorrhages.

I know of no case where it has been alleged that death was due to the X-rays. The Galway case, tried in the Dublin courts in March, 1904, contains the best account on record of the legal liabilities, the case being decided for the defendants that the burn was not caused by

---

<sup>1</sup> EDITORIAL, *Jr. Amer. Med. Assoc.*, January 9, 1904, p. 103.

<sup>2</sup> *Wiener klin. Wchnschr.*, 1902, nos. 16, p. 405, and 17, p. 446. See also Bors, *Arch. d'électricité med.*, Bordeaux, 1903, xi, p. 608.



negligence. Schönberg has shown that all the male rabbits and guinea-pigs exposed to the Röntgen rays proved sterile afterwards.

**DEATH FROM HEAT OR COLD.**—After fatal heat-stroke the body is often very hot for hours and decomposition may be uncommonly rapid. There may be general internal congestion. It is usually necessary to know somewhat of the history of the case before a verdict can be rendered of heat-exhaustion, sunstroke, or thermic fever. In cases of death from cold we often find pallor or discoloration of the skin and congestion of the viscera with blood of rather bright color. No single characteristic lesion results from exposure to moderate excess of either heat or cold. When no pathologic lesions can be found, death is probably due to shock. Any chronic disease of viscera tends to reduce the power to resist severe temperature changes. There is no significance in the freezing of the body beyond showing that considerable time may have elapsed since death. The frozen flesh of the mastodon sometimes found in the Siberian plains is good eating, though it must be thousands of years old.

There are no characteristic changes in sunstroke. Rigor mortis comes on early. Lividity and putrefactive changes develop rapidly after and even before death. Venous engorgement is extreme, particularly in the cerebrum. The left ventricle of the heart is contracted; the right is dilated and may be full of blood imperfectly coagulated and deficient in oxygen. The blood is fluid, dark in color, acid in reaction, and probably contains, as in burns, a poisonous substance which acts on the more highly specialized cells of the body. Petechial patches may appear in the subcutaneous and subserous tissues. The elevation of temperature is often remarkable, and it is extremely disagreeable to make an autopsy in these cases, as I have done, soon after death, with a temperature of 106° F. In a case of mine of stramonium poisoning, with a temperature of nearly 110° F., the clinician had diagnosed sunstroke.

**INFANTICIDE.**—Many methods have been resorted to, as exposure to cold, smothering in various ways, strangulation either by the hands or by a ligature around the neck, and wounding with various instruments, sometimes accompanied by efforts to conceal the act. The child may be intentionally drowned in a vessel containing fluids discharged from the vagina at the time of birth. Gross violence or poisons may be employed.

**DEATH BY STARVATION.**—There is usually extreme emaciation, which is shown especially by a sinking of the eyes and an unfilled

condition of the skin. It is sometimes necessary to determine whether starvation resulted from disease or neglect, especially in cases of those children which have been reared in foundling homes and hospitals.

SUFFOCATION; STRANGULATION; HANGING; DROWNING.—All these produce death by asphyxia, or carbon-dioxid poisoning, combined with oxygen starvation, the signs of which are more or less marked. In death from asphyxia there are usually hemorrhages into the thymus gland, as well as Tardieu ecchymoses in the pleura and pericardium.

Plain suffocation may show no marks of violence. The dark fluid blood, possibly hemorrhages from increased blood pressure, general congestion of the lungs, frequently congestion of viscera, often blue nails and lips, occasionally suffusion of the face with dark venous blood, and an absence of other pathologic conditions, give a general type of finding that is not easily mistaken when clearly marked but is difficult to recognize when not conspicuous.

Strangulation adds the factor of mechanical arrest of respiration, and may result from the presence of food, some foreign substance, or a growth or swelling in the throat. When due to throttling the marks about the neck are of great importance. There may be compression of veins.

Hanging may cause death by injury to the spinal cord as well as by compression of the blood-vessels and air-passages. The parchment-like appearance of the skin on the sides of the neck and the rupture of the intima of the carotids afford valuable evidence.

Wachholz<sup>1</sup> has shown experimentally that in acute suffocation there may be found, along with the soft currant-jelly clots in the heart, solid white clots embedded in the meshes of the cardiac muscle. La Casagne and Martin have described a method, called *docimasie hépatique*, of diagnosing sudden death by a marked increase in the sugar contents of the liver of persons who have died suddenly. Wachholz finds from his experiments that no such relation exists.

Reuter, working with Kolisko,<sup>2</sup> from a study of twenty-two cases of throttling and two hundred cases of hanging, thinks that these two very similar modes of death may be differentiated from each other. In throttling there is (1) cyanosis of the face, with ecchymoses of the

<sup>1</sup> *Vrtljschr. f. gerichtl. Med.*, 1902, p. 34.

<sup>2</sup> *Zeitschr. f. Heilk.*, 1901, vol. xxii, p. 145.

eyelids and conjunctiva. (2) The scalp, the coverings of the brain, and its membranes are always rich in blood. (3) As a rule, hemorrhages in the soft tissues of the neck, especially in the muscles, occur. (4) There is marked injection of the upper air-passages, combined with numerous small hemorrhages. (5) Injuries to the larynx and hyoid are rare. (6) Rupture of the intima of the carotid is never noted; in only three cases were there suffusions into the adventitia. In hanging (1) cyanosis of the face is usually not noted; ecchymoses are seen in twenty per cent. of typical and in thirty per cent. of atypical strangulations. (2) The amount of blood contained in the organs in the skull varies, but usually consists only of that which was present in these parts at the time the circulation was interrupted. (3) Hemorrhages in the muscles are rare,—two per cent. in typical and fourteen per cent. in atypical cases. (4) Injuries to the laryngeal and hyoid structures are common,—sixty per cent. in typical and thirty per cent. in atypical cases. (5) Rupture of the intima of the carotids occurs in five per cent. of typical and four per cent. of atypical hangings. The external markings on the neck are also often different.

In a case of drowning water or foreign substances may be found in the openings of the body, in the respiratory organs, or in the stomach, or death may be due to spasmodic arrest of respiration. The froth from the air-passages is coarser than that seen in cases of œdema. Very soon after death we often find watery fluid in the pleura. The spongy condition of the lungs is found only where there has been inhalation of water, which does not always happen. After decomposition has set in, the evidence of drowning gradually disappears until it is impossible to make the diagnosis. In drowning the bleaching of the palmar and plantar skin surfaces occurs very early. Littlejohn<sup>1</sup> discusses the differences in appearance after drowning in salt and in fresh water. Of those drowned in sea water the soft parts are rapidly destroyed by crabs and fishes, in some cases the bones alone remaining after ten days, while the body undergoes putrefactive changes more slowly. He reports a case where calcium phosphate crystals studded the pleura. The place in which a person is drowned may sometimes be told by the character of the material found in the smaller bronchi. Revenstorff<sup>2</sup> determines the freezing-point of the

<sup>1</sup> *Edinburgh Med. Jr.*, February, 1903, p. 123.

<sup>2</sup> *Münch. med. Wchschr.*, 1902, no. 45, p. 1880.



blood from both sides of the heart, as more or less of the fluid in which an animal is drowned usually passes through the capillaries of the lungs and dilutes the venous blood. He concludes that the method, when positive,—*i.e.*, when it can be shown that the freezing-point of the blood from the right side of the heart is higher than that of the blood from the left side,—is valuable as additional evidence of drowning, and is very easily carried out; but decomposition rapidly removes any difference which may have existed, and the blood is not necessarily diluted during death by drowning.

**CYTOLOGY.**—The different kinds of cells found under various conditions in the serous cavities form a most inviting field of study. Thus, in syphilitic hydrocele we have endothelium, in gonorrhœal hydrocele, marked polymorphonuclear leukocytosis, in tuberculous hydrocele, lymphocytosis, in mechanical hydrocele, few or no leukocytes. Naturally, the age of the process has much to do with the number and variety of the cells.

**SEMEN.**<sup>1</sup>—The Florence test should be first applied, a reaction common to all semen, and then the material studied microscopically. Seminal stains remain intact for years under favorable circumstances and give the biologic blood test.

Schütze<sup>2</sup> finds in the use of the precipitin method of discovering spermatozoa that the animal need not be inoculated with semen or testicular cells, but that any albuminous fluid of the animal's semen to be proved will produce a serum capable of giving rise to the reaction.

**TOXICOLOGY.**<sup>3</sup>—The presence of poisons in the animal economy may be recognized clinically, chemically, pharmacologically, and pathologically. While we have chiefly to do with the latter method, the success of the chemist and the pharmacologist depends largely upon the procedures adopted for the preservation of material by the pathologist at the time of the performance of the autopsy. There are certain poisons which may kill without leaving in the tissues any specific alterations to be found *post mortem*, especially when the examination is postponed for several days.

A poison is any substance which, when taken into the system and

<sup>1</sup> See SIMON's *Clinical Diagnosis*, 1904, p. 664, or any other recent work for the method of applying this test.

<sup>2</sup> *Zeitsch. f. Hyg. u. Infektionskrank.*, 1901, vol. xxxvi, p. 5.

<sup>3</sup> Much of the material in this section is taken from ROBERT's *Lehrbuch der Intoxikationen*, Stuttgart, 1902, and GLAISTER's *Medical Jurisprudence*, 1903.

either being absorbed or by its direct chemic action upon the parts with which in contact, or when applied externally and entering the circulation, is capable of producing deleterious results. (Wormley.) Poisoning commonly results from alcohol, morphine, lead, arsenic, phosphorus, oxalic acid, carbolic acid, etc.; from food (bromatotoxismus); from meat (kreotoxismus); from milk products (galactotoxismus); from fish and shell-fish (ichthyotoxismus, mytilotoxismus); and from grain (sitotoxismus); of the latter poisoning there are three kinds,—ergotism, lathyrism, and pellagra.

It should always be remembered that conditions which we are apt to regard as being alone produced by strictly pathologic processes are often due to poisons. Thus, toxic inanition may be produced by chronic poisoning with mercury, lead, arsenic, etc.; fatty degeneration, by phosphorus, alcohol, *Amanita phalloides*, etc.; calcification of the renal epithelium, by corrosive sublimate; and amyloid degeneration, by repeated injections of turpentine.

Suspicious undissolved foreign bodies may be found in the vomit and in the contents of the alimentary tract, as arsenic (white, metallic, and various salts), antimony, sulphide of antimony, mercury and its preparations, as calomel, oxid, and bichlorid, chrome salts, oxalates, cantharides, nux vomica beans, heads of matches, and parts of poisonous plants. In one of my cases diagnosed as a heat-stroke, with a temperature of over 110° F., the finding of leaves of *Datura stramonium* in the stomach led to the correct diagnosis. Morphine even when given hypodermically may be found in the stomach contents. Certain chemicals may be detected by odors coming from the body or from the various cavities when opened, as alcohol, ether, chloroform, aromatic oils, formalin, phosphorus, turpentine, nitrobenzol, benzene, wood alcohol, hydrocyanic acid, paraldehyde, camphor, chloral, carbolic acid, nicotine, bromin, chlorin, iodine, ammonia, hydrochloric acid, opium, sulphuretted hydrogen, etc. (See also p. 21.)

When the acidity or alkalinity of the gastric contents is abnormally increased, certain reagents are to be suspected, such as acids, alkalies, and potassium cyanid. The liver especially shows poisoning by phosphorus, antimony, arsenic, and toxins, while the kidney is affected by hæmolytic and methæmoglobinic poisons, by oxalic acid, oxamid, mercury, silver salts, preparations of cantharides, etc. The spectroscopic picture of the blood should always be obtained as soon after death or removal from the body as possible. The addition of



a little distilled water is admissible in methæmoglobinæmia, but even here it is better at once to seal hermetically in glass tubes with exclusion of air as far as practicable. If the blood coming from veins is fluid and scarlet, suspect carbon monoxid poisoning; if a laky purple fluid, not changing on the exposure to oxygen, suspect cyanid. If the muscles of the abdominal walls are drawn and contracted spirally, we may suspect any of the instant poisons, as strychnine or potassium cyanid. I have for a long time had a bottle of blood from a case of cyanid poisoning, and have many times exposed it to the air by removing the cork, yet it is apparently still in a perfect state of preservation.

The left heart is found markedly contracted in death from overdoses of members of the digitalis group, veratrine, and barium salts. As already stated, the odor of the poison may sometimes be detected on exposing the brain. In one of my cases of ammonia poisoning a rod dipped in hydrochloric acid gave off fumes when introduced into the cranial cavity after removal of the brain. Much attention has been paid to the actions of poisons on the central nervous system, and the rapid diagnosis of hydrophobia by this method should not be forgotten. For a description of the Negri bodies in hydrophobia, see the *Zeitschrift f. Hyg. u. Infektionskrank.*, 1903, vol. xliv, p. 519. The joints are alleged to be inflamed after poisoning by colchicum. Testicular atrophy is said to be induced by the long-continued use of capsicum, solanus pseudocapsicum, and conium maculatum.

The mucous membrane of the stomach is irritated and stained by many poisons, as sulphuric acid (black), nitric acid (yellow), oxalic acid (white), bromin (red), iodin (purple), and by a large number of metallic salts, as sulphid of arsenic (yellow), chromate of potassium (red), etc. I have, however, seen several cases of arsenical poisoning with but little inflammation of the gastric mucosa.

Among the questions to be answered in every case of suspected poisoning are: Was death caused by a poison originating within or without the body? What poison caused death? Is the substance found by the chemist the poison which killed the person in whose body it was found? Might not the poison have been administered as a medicine? Is the poison present in such quantity as always causes death? Were there attendant circumstances which conduced to the fatal result? Was more than one poison given? How and when was the toxic substance administered? Could poison have been given



and yet not be discovered? Was the fatal dose taken for purposes of suicide? Was it administered with the object of killing? Was it administered accidentally? Did the person for whom it was intended receive the poison? Could the toxic symptoms be simulated? Was cremation practised in order to destroy evidences of poisoning? Was there any motive for homicide? Are there any accomplices? What became of the vehicle in which the poison was administered? Was there any poison found? Was any poison destroyed?

In case of poisoning the district attorney must prove three things, in order to convict of murder in the first degree: first, that the person is dead; second, that death was caused by the poison under consideration; and, third, that the party or parties on trial administered the drug with felonious intent.

On request of ex-Judge Stevenson, the lawyer for the defence, in the case of the Commonwealth of Pennsylvania *vs.* John and Emma Williams, Judge McMichael issued the following order, under which I made a post-mortem examination of the bodies of the three children, after the experts for the Commonwealth had already made an original examination and two disinterments.

"And now, to wit, this sixteenth day of February, A. D. 1903, it is ordered and directed that the defendants through their experts shall have access to the bottles and prescriptions taken from 1135 Vienna Street, and also that they shall be permitted to exhume the bodies of Anna, Josephine, and Laura Williams, and to make such examination as they shall deem necessary and proper, and to remove such portions of said bodies as may be necessary to a proper and adequate chemical examination and analysis to determine the causes of death. It being understood, however, that the Commonwealth shall during these investigations be represented by an expert that they may select."

Nearly every toxicologist has his own classification of poisons. Thus, one divides them into mineral, vegetable, animal, and mechanical groups, another into irritants, narcotics, and narcotic irritants, a third into chemical and vital poisons, etc. All such divisions are arbitrary, as quickly becomes evident on attempting to place the various poisons in their proper subclasses.

#### SCHEME FOR THE DIVISION OF POISONS.

INORGANIC	{	Irrespirable gases : carbon monoxid, coal gas, chlorin, bromin,
		hydrofluoric acid, sulphur dioxide, etc.
		Chemic : sodium hydrate, sulphuric acid, etc.
		Irritant : arsenic, antimony, mercury, phosphorus, etc.

SCHEME FOR THE DIVISION OF POISONS.—( *Continued.* )

ORGANIC	{	Irrespirable gases : chloroform, ether, formalin, etc.
		Chemic : carbolic acid, acetic acid, pyrogallic acid, etc.
	{	Irritant { Vegetable : gamboge, colchicum, squill, etc.
		Animal : cantharides, etc.
	{	Alkaloidal { Narcotic : opium, hyoscyamus, belladonna, cannabis indica, etc.
		Sedative : digitalis, hydrocyanic acid, aconite, conium, etc.
		Excitomotor : strychnine, ergot, etc.
{	Synthetical	Antiseptics : creolin, lysol, etc.
		Antipyretics : antipyrin, acetanilid, etc.
		Hypnotics : sulphonal, trional.
{	Toxic	Bacterial : toxins, hæmolysins, cytolsins.
		Animal : snakes, scorpions, ptomaines, etc.
		Vegetable : ricine, abrine, etc.

SYMPTOMS OBSERVED AFTER THE ADMINISTRATION OF THE MORE COMMON POISONS.<sup>1</sup>

ACUTE SYMPTOMS :	THINK OF :
1. Death within a few seconds or minutes.	Hydrocyanic acid; potassium cyanid; carbonic acid; carbolic acid.
2. Deep coma.	Alcohol; morphine; opium; chloral hydrate and its derivatives; sulphonal; chloroform and its derivatives; carbon monoxid; anilin oil; oxybutyric acid.
3. Collapse.*	Corrosive acids; corrosive alkalies; nicotine; arsenic; antimony; colchicine.
4. Feverish rise of temperature.	Phosphorus; cocaine; under certain circumstances any of the powerful convulsive remedies; enzymes.
5. Mania; furious delirium; psychic excitement.	Chronic alcoholism; atropine; cannabinone; camphor; physostigmine; veratrine; lead (in animals).
6. Mental disturbances of the most diverse kind.	Alcoholism; morphinism; cocainism; pellagra; ergotism; inhalation of ether; saturnism; mercurialism; poisoning by bromid; iodoform; carbon bisulphid.
7. Violent; at times, tetanic convulsions.	Strychnine; toxin of tetanus; salts of ammonia; cytisine; cornutine; picrotoxin; cicutoxin; active principles of digitalis; cocaine; santonine; *aconitine; gelsemine; filicic acid.

<sup>1</sup> This table and the following one are from KOBERT's *Compendium der Toxikologie*, 1903.

## SYMPTOMS OBSERVED IN CASES OF POISONING.

## ACUTE SYMPTOMS:

## THINK OF:

- |   |  |
|---|--|
| 8. General paralysis, for the most part ascending.                          | Coniine; curarine; colchicine.   |
| 9. Paralysis of individual groups of muscles.                               | Lead; arsenic; carbon bisulphid.   |
| 10. Dilatation of the pupil.  | Atropine; hyoscyamine; scopolamine; cocaine; ephedrine; aconitine; coniine; gelsemine; sausage poisoning.  |
| 11. Contraction of the pupil.   | Muscarine; pilocarpine; nicotine; arecoline; morphine; codeine; opium; physostigmine.  |
| 12. Amaurosis.  | Quinine; salicylic preparations; extract of male fern; belladonna; uræmic poisoning.   |
| 13. Diplopia and ptosis.  | Sausage and fish poisoning.  |
| 14. Conjunctivitis.   | Irritating vapors (sulphurous, hydrochloric, nitric, and osmic acid; nitrogen dioxid; hydrofluoric acid; chlorine; bromine; carbonyl chlorid; ammonia; ethereal oil of mustard; croton oil vapor); irritating kinds of dusts, as root of ipecacuanha, quillaja bark, pepper, chromate, picrate; arsenism; phenylendiamin, chrysarobin, [formalin]. |
| 15. Moist skin.   | Opium; morphine; aconitine; muscarine; pilocarpine; nicotine; physostigmine; lobeline; antimony.   |
| 16. Skin conspicuously dry, even in a warmed bed. Mouth and throat parched. | Atropine, as well as belladonna, stramonium, and hyoscyamus; hyoscyamine; scopolamine; sausage and fish poisoning.   |
| 17. Urticaria or scarlatiniform erythema.                                   | Atropine; hyoscyamine; antipyrin; quinine; balsam of copaiba; cubebene; chloral hydrate; iodine; morphine; and many internal remedies; handling of nettles (urtica).   |
| 18. Eczematous eruptions of the skin.                                       | Croton oil; curcas oil; cardol; rhus toxicodendron; powdered cinchona bark; carbolic acid; tar.  |
| 19. Diffuse dermatitis, with perspiration of the hands.                     | Anilin colors; aurantia; chrysoidin; malachite green; Bismarck brown; butter yellow; anilin yellow.  |
| 20. Acne pustules.  | Bromid; arsenical and antimonial preparations; powdered ipecacuanha.   |
| 21. Blisters containing clear serum on the skin, or even in the mouth.      | Spanish fly; ranunculus acris; ranunculus sceleratus, etc.   |
| 22. Dark, dirty discoloration of the skin, which is not, however, cyanotic. | Argyria; mercurialism; saturnism; arsenical melanosis; bronzed diabetes.   |



SYMPTOMS OBSERVED IN CASES OF POISONING.

ACUTE SYMPTOMS:

23. Bluish discoloration of the peripheral portions of the body [like those seen in Raynaud's disease].
24. Cyanosis.
25. Yellowish-brown discoloration of the conjunctiva, alone or in combination with that of the skin.
26. Discoloration primarily of the tongue and the mucous membrane of the mouth.
27. Secondary discoloration of the gums.
28. Specific odor to the breath.
29. Coryza.
30. Salivation.
31. Metallic cough and aphonia.
32. Edema of the glottis.
33. Edema of the lungs.
34. Luminosity of the breath and vomitus.
35. Increased liver dulness.
36. Diarrhœa with vomiting.

THINK OF:

Gangrenous ergotism; carbolicism; phosphorism.

Nitrobenzol; benzokoll; anilin; toluidin; antifebrin; exalgin.

Phosphorus; helvellic acid; potassium chlorate; nitroglycerin; sodium nitrite; amyl nitrite; pyrogallol; arseniuretted hydrogen; ietrogen [lupinotoxin], in animals. (In picric acid and picrates the discoloration is a pure yellow.)

Reddish yellow—chromic acid and the bichromates. Yellow—nitric and picric acid. Brown — iodine; bromine. Greenish-blue — salts of copper; Schweinfurth-green. Whitish — corrosive alkalies; corrosive acids; corrosive metallic salts; carbolic acid.

Lead; silver; mercury; bismuth. [Bring out for diagnostic purposes by the direct application of sulphuretted hydrogen gas to the gums.] (See pages 21 and 418.)

Iodine; bromine.

Pilocarpine; muscarine; arecaline; nicotine; cornutine; physostigmine; cytisine; mercury; ammonia; saponine [a glucoside contained in *Saponaria officinalis*]; cantharidine; caustics.

Atropine; hyoscyamine; scopolamine; sausage poisoning.

All caustic poisons.

Morphine; muscarine; pilocarpine; ammonia; nitric acid vapors, etc.; inhalation during the swallowing of sublimate and other corrosive poisons.

Phosphorus.

Phosphorus; agaricus bulbosus; poley oil; alcohol.

Salts of antimony; arsenic; digitaline; pilocarpine; nicotine; muscarine; colchicine; corrosive poisons; salts of copper; zinc salts; colocynthis; emetine; cephaline; croton oil, etc.

## SYMPTOMS OBSERVED IN CASES OF POISONING.

ACUTE SYMPTOMS:	THINK OF:
37. Vomiting without diarrhoea.	Apomorphine; lobeline; cytisine.
38. Colic with constipation.	Lead salts.
39. Colic with diarrhoea.	Barium salts.
40. Diarrhoea without vomiting.	Jalap; podophyllotoxin; croton oil; calomel, etc.
41. Pulse continuously and markedly slowed.	Opium; morphine; muscarine; arecaline; physostigmine; baryta; all narcotics.
42. Pulse paroxysmally and markedly slowed and thread-like.	Lead salts, but only during an attack of lead colic.
43. Pulse first slowed, then irregular, lastly accelerated.	Digitalis; hellebore; adonis; coronilla; cheiranthus; nerium; scilla; trophanthus; convallaria; pilocarpine; nicotine; scopalamine.
44. Pulse greatly accelerated.	Belladonna, hyoscyamus; atropine.
45. Abortion.	Sabina; thuja; rue; mentha pulegium; phosphorus; ergot; lead.
46. 6-12-hour period of good health between the poisoning and the appearance of the symptoms.	Most of the poisonous fungi, but especially <i>Amanita phalloides</i> ; also combined arsenic.

## TABLE OF THE MOST STRIKING CHANGES WHICH TAKE PLACE IN THE URINE AFTER THE ADMINISTRATION OF THE MORE COMMON POISONS AND MEDICINES.

URINARY CONDITION:	THINK OF:
1. Very acid reaction.	Mineral acids; acid salts of the metals.
2. Reaction strongly alkaline.	Corrosive alkalies; alkaline carbonates; salts of organic acids, with the exception of oxalic acid.
3. Odor like violets.	Oil of turpentine and related ethereal oils when employed medicinally.
4. Odor like garlic.	Preparations of tellurium when employed medicinally.
5. Odor of methylmercaptan.	Asparagus, sometimes used medicinally in the form of a syrup.
6. Odor of rotten eggs.	Cystinuria exists, or the thiosulphate of sodium has been taken medicinally in large doses.
7. Odor of ammonia.	Ammoniaemia; cystitis caused by strong bases [and certain bacteria].
8. Achromatic crystals, with acid urine.	Oxalic acid; binoxalate of potassium; oxamid; parabanic acid.
9. Leucocytes and epithelial casts.	Cantharidine; potassium cantharidinate; virus of scarlet fever.
10. Yellow to yellowish-red color.	Picric acid; picrates.

EFFECT OF POISONS UPON THE URINE.

URINARY CONDITION:	THINK OF:
11. Urine icteric brown.	Phosphorus, toluylendiamin; cephalanthine; ictrogen.
12. Urine, reddish.	Senna leaves; rhubarb root; campecia wood (logwood); hæmatoxylin; fuchsin; pyramidon; antipyrin.
13. Urine colored wine-red by hæmatoporphyrin.	Sulphonal; trional; tetranol; lead (rarely).
14. Urine becomes scarlet upon putrefaction.	Santonine; santonica seeds; chenopodium.
15. Urine contains albumin and red blood-corpuscles.	Corrosive poisons of all sorts.
16. Urine contains blood pigment in solution.	Arseniuretted hydrogen; helvella esculenta (helvellac acid); cyclamine; solanine; and other saponiferous substances.
17. Urine contains methæmoglobin.	Potassium chlorate; sodium nitrite; amyl nitrite; pyrogallol; chrysarobin; kairin; quinine.
18. Urine contains urobilin.	Lead.
19. Urine becomes black-green on exposure to the air.	Carbolic acid; cresol; lysol; creosote; guaiacol.
20. Urine is green when voided.	Methylen blue.
21. Urine on exposure to the air becomes blackish-brown or even pure black.	Melanuria, associated with melanotic tumors and with hæmochromatosis. It may be produced artificially by injections of melanin.
22. Urine reduces Fehling's solution and gives off carbon dioxid with yeast.	Phloridzin; salts of uranium; curarine; hydrocyanic acid; atropine; amyl nitrite; chromates and bichromates; bichlorid of mercury; cantharidine.
23. Urine reduces Fehling's solution, but yields with yeast little or no carbon dioxid.	Chloral hydrate; menthol; thymol; many of the ethereal oils; carbon monoxid; chloroform; formic acid and formates; free oxalic acid; benzaldehyd; morphine.
24. Urine polarizes light to the <i>right</i> .	Phloridzin; salts of uranium; curarine; hydrocyanic acid; atropine; amyl nitrite; chromates and bichromates; bichlorid of mercury; cantharidine.
25. Urine polarizes light to the <i>left</i> .	Chloral hydrate; menthol; thymol; many of the ethereal oils.
26. Urine contains increased number of paired sulphuric acids and diminished number of sulphates.	Carbolic acid; cresol; lysol; creosote; guaiacol; kairin; antifebrin; anilin; paramidophenol.
27. Urine contains leucin and tyrosin.	Phosphorus; acute yellow atrophy of liver; pellagra.



found, particularly about the big toe. The brain is sometimes shrunken and dry, the blood-vessels being constricted; or these organs may be pale and extremely firm, or pale and œdematous, as in cases of uræmia. The small intestines may show areas of extreme contraction. For the detection of lead in urine and post-mortem specimens, the reader is referred to the *Lancet*, September 12, 1903, p. 746.

**MERCURIAL POISONING.**—The mucous membranes of the gastrointestinal tract, especially the small intestine and cæcum, show extensive desquamation, with hyperæmia, ecchymoses, and grayish-white eschars. The bowel generally contains large quantities of liquid of a yellowish-brown or blood-stained character. The macroscopic appearances are those of dysentery. In some acute cases decalcification of the bones occurs, with a deposit of lime elsewhere in the body, especially in the kidneys. The number of mercurial salts is legion, many forming with albumin an insoluble albuminate of mercury. Chronic cases of poisoning occur, ulcerative stomatitis being one of the chief lesions. Sebillotte, in 1891, collected one hundred and forty-eight cases of poisoning from post-partum vaginal douches of bichlorid of mercury. He expressed his belief that the poison was not absorbed through the healthy mucous membrane, but through laceration of tissue due to the process of labor. Hamburger, however, has found that potassium iodid appeared in the urine in twenty-four hours when tampons of cotton saturated with this substance were placed in the healthy vagina, and potassium ferrocyanid or salicylic acid in three hours. H. C. Wood, Jr., reports a case of poisoning with bloody urine from the use of a douche containing 1 to 2000 of the bichlorid of mercury.<sup>1</sup>

**METHYL ALCOHOL.**—Blindness or impairment of vision may occur not only from the ingestion of wood alcohol, but also from inhalation of its fumes, as methyl alcohol seems to have a predilection for the retina and the optic nerve. A number of cases of fatal poisoning from this source have recently occurred throughout America. These have been tabulated by Buller, of Montreal, and Wood, of Chicago.<sup>2</sup>

**NITROBENZOL POISONING.**—Besides the odor of the artificial oil of bitter almonds, the blood and muscles are of a brownish color and the mucous membrane of the stomach is ecchymotic and injected. The body is cyanosed and of a leaden hue.

---

<sup>1</sup> *Amer. Med.*, December 27, 1902, p. 1006.

<sup>2</sup> *Jr. Amer. Med. Assoc.*, October 1, 8, 15, 22, and 29, 1904.

**NUTMEG POISONING.**—For a description of this rare but interesting form of poisoning, the reader is referred to Wallace's article in Vaughan's dedication volume of "Contributions to Medical Research." Grated nutmeg is used by some as an emmenagogue.

**OPIUM POISONING.**—In acute poisoning there is nothing to distinguish the condition of the brain from that in other cases of cerebral congestion. Extreme passive congestion of the bases of the lungs may take place, as in cerebral apoplexy (Osler). Cases of uncomplicated chronic poisoning are rare. The most important lesion is fatty degeneration of the heart. The liver may show similar changes. If laudanum has been used, the characteristic odor may be present. I know of no drug which is more apt to escape detection at the postmortem than morphine, as there are absolutely no characteristic lesions and chemic analyses are difficult and at times inaccurate. It seems strange that one of the most common and easily accessible poisons is thus so hard to detect. The pupillary reaction is of no value after death, and the clotting of blood in the right heart is by no means constant. Many, if not all, of the chemic tests for morphine may be simulated by the effect of putrefactive bodies. Kippenberger's method is not considered reliable by Clift.<sup>1</sup>

**PELLAGRA POISONING.**—The lesions found are in the posterior columns and the crossed pyramidal tract. The cells in the anterior horn are deeply pigmented, and pigment is found in the internal organs and the skin. The brain presents general wasting; the ventricles are somewhat distended and contain an excess of fluid.

**PHOSPHORUS POISONING.**—In acute phosphorus poisoning the gastro-intestinal tract, especially in the stomach, shows an intense degree of inflammation. Hemorrhages are common and the stomach may contain grumous (coffee-ground) blood. The mucous membrane is the seat of numerous ecchymoses as well as more or less extensive necroses. The skin, the serous membranes, the muscles, and the adipose tissues all show numerous small hemorrhages. The blood is liquid and dark. The skin is jaundiced. The liver, in the early stages increased in size, soon—in from ten to fourteen days—becomes small (from one-half to one-third of the normal bulk), the capsule is wrinkled and shrunken, the color is pale yellowish, and on section the organ presents yellowish patches in the midst of which are areas of deep

<sup>1</sup> *Jr., Amer. Med. Assoc.*, April 23, 1904.

congestion. Drops of fat are seen upon the knife. The kidneys are large, their cortex pale, and the medullary portions congested. The epithelium often shows marked granular degeneration. As a rule, the spleen is not markedly altered. In chronic poisoning by phosphorus wide-spread fatty degeneration is the rule. In cases of workers in phosphorus having defective teeth, necrosis of the jaw is not uncommon. It is the yellow phosphorus that is poisonous and not the red variety. Bug exterminators often contain phosphorus. The coating from the ends of matches is sometimes taken with suicidal intent.

PICROTOXIN.—Carel<sup>1</sup> gives the proceedings in three cases of homicidal poisoning by picrotoxin, derived from the *cocculus indicus* berries added to the liquor of half-drunken men for the purpose of robbery after the production of unconsciousness.

POTASSIUM CHLORATE POISONING.—The blood has the color and consistence of chocolate, the oxyhæmoglobin having been reduced to methæmoglobin. There is usually a hemorrhagic nephritis, especially of the glomeruli.

PTOMAIN AND TOADSTOOL POISONING.—Such cases are of especial interest to the toxicologist, as the symptoms produced and the lesions found at the postmortem are similar to those caused by many alkaloidal and irritant poisons, and the possibility of the case under consideration in a trial being due to one or other of these substances is always suggested by the defence.

RICIN POISONING.—In dogs the eosinophile cells are increased in number. There is no marked positive degeneration of the liver, though the organ is congested and areas of necrosis are seen. In the kidneys the epithelial cells show degeneration.<sup>2</sup>

SILVER NITRATE POISONING.—I have been fortunate enough to see one case of this rare form of poisoning. The darkening of the necrosed mucous membrane on exposure to light was the chief diagnostic point. The child had an inspiration pneumonia.

SNAKE POISONING.—After death caused by cobra bite rigor mortis occurs as usual. The areolar tissue in the region of the bite is infiltrated with a pinkish fluid and the vessels are injected. The blood presents no demonstrable change. The veins of the pia mater are usually engorged, and the ventricles often contain turbid fluid. The

<sup>1</sup> *Merck's Archives*, July, 1904.

<sup>2</sup> MÜLLER, *Ziegler's Beiträge*, vol. xxvii, p. 331.



lungs are generally congested and the lining of the bronchi injected. The appearance of the kidneys varies from normal to one of intense congestion. After death following the bite of an Australian snake the appearances are much the same as those just described. The blood may contain soft coagula, the lungs are sometimes the seat of hemorrhages, and the mucous membranes may be intensely congested and hemorrhagic. The central nervous system shows engorgement of the blood-vessels. At autopsy, after the bite of a viperine snake, the region of the wound is seen to be the seat of intense œdema and extravasation of blood, and the underlying muscles are frequently disorganized and even diffuent from the latter cause. Hemorrhages may also be found in any of the organs and along the alimentary tract. The kidneys are acutely congested or hemorrhagic. The blood is fluid. Snake venom alone is not poisonous, but it takes a serum complement, like lecithin, to make it so. (Flexner and Noguchi.) Keyes<sup>1</sup> describes a method of preparing a pure crystalline compound of the toxin. The use of cryoscopy in this and other forms of poisoning may prove of value.

**STRYCHNINE POISONING.**—Rigor mortis is intense and persistent and the blood is dark and fluid as in asphyxia. Be sure to save the urine if any be present; a frog placed in it will have convulsions, even if but a small amount of strychnine be present. Marshall<sup>2</sup> reports the method of analysis used in a recent case with success.

**TANNIN.**—This substance, so useful as an antidote in various forms of poisoning, may itself produce violent diarrhœa and vomiting.

---

<sup>1</sup> *Berl. klin. Wchnschr.*, 1903, nos. 42 and 43.

<sup>2</sup> *Amer. Med.*, June 18, 1904.

## CHAPTER XXVII

### THE PRUSSIAN REGULATIONS FOR THE PERFORMANCE OF AUTOPSIES IN MEDICOLEGAL CASES

THE Prussian regulations governing the performance of postmortems by the legally appointed officers of the court are of great historic interest, as they bear the imprint of Virchow, and, though put in force February 13, 1875, are still observed throughout Prussia. These regulations also form the basis of similar statutes in other German states and in many countries throughout the world; indeed they are so well defined that it is advisable, though one may chafe under their apparently unnecessary restrictions, to depart from them only in exceptional instances. This is especially the case if the one performing the autopsy is a beginner in medicolegal work.

#### I. GENERAL CONSIDERATIONS.

- |  |  |
|--|--|
| The Physicians making the Autopsy, and their Duties          | § 1. According to the present law, an examination of a corpse for medicolegal purposes may be made only in the presence of a magistrate by two practitioners, one of whom should be a state-appointed physician and the other a district surgeon. Upon those performing the autopsy devolve the duties of medicolegal experts. If doubt should arise in the technical performance of the autopsy, the physician or his deputy decides the question under consideration conditionally upon the right of the surgeon to state upon the protocol his dissenting opinion.  |
| Substitutes  | § 2. The medical officers are permitted to appoint substitutes only when legitimately detained from the performance of their medicolegal duties. If possible, the deputy chosen is to be a physician who has passed his <i>pro physiatu</i> examination.   |
| Time after Death at which the Post-mortem is to be performed | § 3. As a rule, postmortems should not be performed until twenty-four hours after death; the mere inspection of a corpse, however, may be made earlier than this.  |
| The Examination of Decomposed Bodies                         | § 4. Generally, post-mortem examinations must not be neglected nor their performance refused by the legally appointed physicians because of the presence of decomposition, for even in a badly decomposed cadaver abnormalities and injuries to the bones may still be detected; many facts of value in the identification of a body may be ascertained, such as the color and appearance of the hair, the absence of limbs, etc.; and substances which have entered the body from without may be discovered, as well as pregnancy or poisoning proved. On the same grounds, when for one reason or another the advisability of disinterring a body is under consideration, the physicians are to approve of such exhumation without regard to the time which has elapsed since death. |

§ 5. The legally appointed physicians are to be careful to have the following instruments in readiness and in good condition: Instruments  
 from four to six scalpels, of which the two smaller ones are to possess a straight and the two larger ones a rounded cutting edge; one razor; two strong cartilage-knives; two forceps; two double hooks; two pairs of scissors,—the stronger pair should have one blade pointed and the other rounded, while the smaller pair should possess one probe-pointed and one sharp-pointed blade; one enterotome; one injecting nozzle with stopcock; one coarse and two fine sounds; one saw; one chisel and one hammer; one costotome; six curved needles of different sizes; one pelvimeter; a one-metre rule divided into centimetres and millimetres; a measuring-glass divided into one hundred, fifty, and twenty-five cubic centimetres; one pair of scales capable of weighing up to ten pounds; one good magnifying-glass; blue and red litmus paper. The cutting instruments must be perfectly sharp. Those performing the postmortem are recommended to have ready for use a microscope with two objectives, so as to be able to magnify at least four hundred diameters, and the required instruments, glassware, and reagents necessary for the preparation of microscopical slides.

§ 6. A sufficiently large, well-lighted room is to be chosen for the autopsy, and all possible care is to be taken in the selection of a suitable place on which to lay the body and in the avoidance of all disturbing surroundings. Post-mortem examination by artificial light, except where postponement is impracticable, is not allowed; should it be done, the reason therefor must be expressly stated in the protocol (§ 27). Place for the Autopsy  
and its Lighting

§ 7. If the body be frozen, it must be brought into a heated place and the autopsy postponed until the cadaver has sufficiently thawed; the employment of warm water or other warm articles to hasten the thawing process is forbidden. Frozen Bodies

§ 8. If possible, when for any reason the body is moved, especially if transported from one place to another, there is to be no excessive pressure made upon any of the individual parts, nor any marked departure from the horizontal position of the organs in the larger cavities. Transportation  
of Corpses

## II. TECHNIC OF THE POSTMORTEM.

§ 9. Those performing the postmortem must hold steadfastly to the object in view, which is to make the investigation with accuracy and completeness. All important findings must be shown to the magistrate by the obducent before they are entered in the protocol. Medicolegal Aspects  
of the Postmortem

§ 10. In those cases in which this appears to be necessary, the examiners are required, as early as feasible before the performance of the autopsy, to ask the magistrate for permission to visit the place where the body was found, and they are to ascertain the position in which the body was discovered and be given an opportunity to examine the clothing which the deceased wore at the time of his or her death. As a rule, however, it is sufficient for them to await the solicitation of the magistrate to undertake these investigations. They are also obliged to ask for information from the magistrate in regard to any disclosures which might be of use to them in the performance of the autopsy or in helping them to make up their deductions therefrom. Duties of the Obducent  
in regard to the Ascertainment  
of Special Circumstances connected  
with the Case  
under Investigation



**Microscopical  
Examinations**      § 11. In cases in which a doubtful finding is to be quickly and definitely settled,—as, for example, the differentiation between blood and a fluid which is merely stained with hæmatin,—a microscopical examination is to be then and there undertaken. When circumstances render this impossible or when difficult microscopical investigations which cannot be made at once are required,—as, for example, of certain tissues of the body,—portions of such tissue are to be preserved under legal protection and as quickly as possible thereafter to be thoroughly examined. It is to be distinctly stated in the report of such findings when the examinations were performed.

**The Postmortem: its  
two main divisions**      § 12. The postmortem is divided into two main parts: A. External examination (inspection). B. Internal examination (section).

**External  
Examination**      § 13. In the external inspection of the body its appearance in general and that of its individual parts in particular are to be noted. In this general examination of the body the following points, in so far as possible, are to be brought out and recorded. 1. Age; sex; size; development; general condition of nutrition; any signs of previous illnesses,—e.g., ulcers of the foot; special abnormalities,—e.g., moles, scars, tattoo markings; increase or absence of limbs. 2. The signs of death and the changes that have already taken place from decomposition.

After removal by washing of any contaminations of the body in the way of blood, fæces, dirt, etc., record is to be made of the presence or absence of post-mortem rigidity; the general color of the skin of the corpse; the kind and degree of coloration and discoloration brought about by putrefaction; and the color, situation, and extent of any areas of hypostatic congestion, which are to be incised and then carefully examined and described, in order to prevent their being mistaken for extravasations of blood.

The following particulars are to be considered in the study of the individual parts. 1. In unidentified persons, the color and other appearances of the hair (head and beard), as well as the color of the eyes. 2. The possible presence of foreign substances in the normal openings of the head, the arrangement of the teeth, and the situation and appearance of the tongue. 3. An examination is next to be made of the neck, the breast, the abdomen, the back, the anus, the external genitalia, and finally of the limbs.

If an injury is found in any of these parts, its shape, situation, and direction with relation to fixed points of the body are to be described and the length and breadth of the injury given in the metric system. In solution of continuity of tissue, probing is, as a rule, to be avoided in the external inspection, because after the internal examination of the body and of the injured spot the extent of the injury becomes apparent. Should the obducent decide that the introduction of a sound is necessary, this procedure is to be done with great care and special mention of the reason therefor is to be made in the protocol (§ 27). When wounds are present, a description of their borders and the adjacent tissues is to be given, and after such an examination and description of the lesions in their original condition the same are to be enlarged in order that the internal appearance of the borders and of the bottom may be disclosed. As to wounds and injuries which clearly did not conduce to, originate from, or have any connection with death,—for example, markings produced in the endeavor to restore life, gnawing by animals, and the like,—a summary description of the findings is sufficient.

§ 14. In the internal examination the three main cavities of the body—the cranial, the thoracic, and the abdominal—are to be opened. Opening of the vertebral column or of the individual joints is not to be omitted in cases where important findings might be secured thereby. When there is a definite suspicion as to the cause of death, the postmortem is to be commenced with that cavity in which the chief changes are suspected. Otherwise the head is to be examined first, the thorax next, and the abdominal cavity last.<sup>1</sup> The situation of the organs found in each of the above-named cavities is first to be determined, then the color and the appearance of the exposed surfaces. The presence is to be noted of any unusual contents, such as foreign bodies, gases, fluids, or clots, and in the last two cases measured and weighed, and finally each individual organ is to be examined externally and internally.

Internal Examination;  
General Considerations

§ 15. When no injuries are present, the opening of the cranial cavity is accomplished by making an incision from one ear to the other directly over the skull, after which the skin-flaps are displaced forward and backward. (In case injuries are present, they should be as much as possible circumvented by the knife, thus giving rise to a different procedure.) As soon as the appearance of the soft parts and the surface of the bony cranium has been described, the latter is cut through with a saw by a circular incision, and the section, the inner table, and the other appearances of the calvarium are described. The external surface of the dura mater is next examined, the longitudinal sinus opened, and its contents estimated. The dura mater is then to be separated on one side and laid back, and the internal surface of the same described, as well as the appearance of the exposed pia mater. After this has been done on the opposite side, the brain is to be removed in as perfect a condition as possible, and the presence of abnormal contents in the skull is to be noted, and the appearance of the dura and pia mater at the base and sides of the skull and the condition of the large arteries are to be described. After the opening of the transverse sinuses (and, in case reason therefor exists, of the remaining sinuses), the size and shape of the brain are noted and an examination is made of its individual parts by means of a series of well-ordered incisions. Such parts include both cerebral hemispheres, the large ganglia (optic thalamus and corpus striatum), the corpora quadrigemina, the cerebellum, the pons Varolii, and the medulla oblongata, in the description of which are to be included especially the color, the fulness of the vessels, the consistency, and the structure. In addition, the tissue and the vessels of the choroid plexus are always to be described. The size and the contents of the different ventricles as well as the appearance and fulness of the different vascular plexuses in the individual sections of the brain are constantly to be kept in mind, and especial note is to be made of the presence of any clotted blood outside of the blood-vessels. The dura mater over the base of the skull and the sides is then to be removed and the condition of the bones in these regions described.

Cranial Cavity

§ 16. When it is required to open the internal portions of the face, to examine the parotid gland, or to inspect the auditory apparatus, the initial incision extending over the skull is continued behind the ear and down the neck, and the skin, for appearances' sake, is dissected

Face, Parotid Gland, and Ear

<sup>1</sup> As to autopsies on the new-born see §§ 23 and 24.

away from beneath towards the part to be investigated. In this examination special attention is to be paid to the condition of the large arteries and veins.

Spinal Column  
and Cord

§ 17. The opening of the spinal column (§ 14) is usually made from behind, the skin and the subcutaneous fatty tissue being cut directly over the spinous processes and the musculature dissected away from the side of the latter and from the vertebral arches. During this examination hemorrhages, lacerations, and similar changes, especially fractures of bones, are to be carefully searched for. Then a chisel, or, if one is at hand, a vertebral saw (rhachiotome) is used for the purpose of separating the spinous processes with the adjacent portions of the arches throughout their entire extent. When they are removed, the external surface of the dura mater, which is now brought into view, is examined. It is next to be carefully opened by means of a longitudinal incision, and any abnormal contents, especially fluid or extravasated blood, are to be described, also the color, appearance, and similar characteristics of posterior portions of the pia mater, and by means of a gentle passage of the fingers over the spinal cord its degree of consistency is to be determined. Next, on both sides, by means of a longitudinal incision the nerve-roots are cut through; then with one hand the lower end of the spinal cord is carefully grasped, and, after dividing the anterior attachments one after another, its upper end is finally drawn out of the occipital foramen. In all these proceedings special care should be taken not to make pressure on the spinal cord or to bend it. When the cord has been removed, the anterior surface of the pia mater is to be examined; next the external appearance of the cord as to size and color is to be described, and finally, by a considerable number of transverse incisions with a sharp and thin knife, the internal appearance of the spinal cord, both as to its white and its gray matter, is to be noted. Finally the dura mater of the vertebral bodies is to be removed, and they are to be examined in order to determine if there have been any hemorrhages, injuries, or changes in the bones or in the intervertebral discs.

Neck, Thoracic and  
Abdominal Cavities;  
General Considera-  
tions

§ 18. The neck and the thoracic and abdominal cavities usually are opened by means of a single long incision from the chin to the pubic symphysis, passing to the left of the navel. Most commonly the incision in the abdomen is made deep enough to penetrate the abdominal cavity, care being taken to avoid injuring the organs contained therein. This is best begun by cutting a small nick in the peritoneum, at the same time observing whether any gas or fluid escapes. One finger is introduced into the opening and then another, the abdominal wall is elevated from the intestines, and the further opening of the peritoneum is made between the two fingers. The situation, the color, and other appearances of the intestines are to be immediately observed, as well as any abnormal contents within them, and the condition of the diaphragm is to be determined by palpation of its under surface.

The examination of the abdominal organs is to be proceeded with at this time only where a strong suspicion exists that the cause of death may be found within the abdomen (§ 14). As a general rule, the thorax is to be opened and inspected before any further scrutiny of the abdominal cavity.

Thoracic Cavity

§ 19. In opening the thoracic cavity the soft parts of the breast are dissected slightly beyond the junction of the osseous and cartilaginous portions of the ribs. Next with a strong knife



the cartilages are incised a few millimetres within their attachment to the ribs, care being taken to avoid cutting the lungs or the heart. If the cartilages be ossified, the ribs are to be separated with a saw or a costotome somewhat beyond the cartilaginous junction. The attachments of both clavicles to the sternum are then separated by vertical semicircular sections, and the junction of the first rib, be it cartilaginous or ossified, is loosened with the knife or costotome, great care being taken to avoid injuring the vessels which lie beneath. The diaphragmatic attachments along the line of incision are severed close to the false cartilages and the ensiform process. The sternum is turned upward and the mediastinum is cut through, with careful avoidance of any injury to the pericardium or the large blood-vessels. When the sternum has been separated, the condition of the pleural cavity is to be determined, especially as to any abnormal contents, which are to be measured and their characteristics described; also the extent and the appearance of any portions of the lung which are in view. If any vessels have been injured in the removal of the breast-bone, they are to be tied or a sponge is to be placed beneath the bleeding points to catch the blood which if it were allowed to enter the pleura would later obscure the observation of the parts therein. The condition of the mediastinum and especially that of the thymus gland are to be noted, as well as the appearance of the large blood-vessels lying outside of the pericardium, which are not yet incised. The pericardium is next to be opened and examined and the exterior of the heart inspected. Before the heart is incised or removed from the body its size, the filling of the coronary vessels and its individual cavities (auricles and ventricles), its color, and its consistency (rigor mortis) are to be estimated. While the organ is still in its natural position, the ventricles and auricles are to be separately opened and the contents of each chamber determined as to their amount, coagulation, and appearance, and the dimensions of the auriculoventricular openings are to be ascertained by the introduction of two fingers through the auricle. The heart is then to be removed from the body and the condition of the arterial vessels tested, first by filling them with water and next by incising their walls. Finally the color and exact appearance of the heart muscle are to be described. In every case wherein it is suspected that extensive changes—*e.g.*, fatty degeneration—have occurred in the muscular tissue a microscopical investigation is to be made. To this examination belongs that of the large vessels, with the single exception of the descending aorta, which is to be examined after the lungs have been excised. A minute inspection of the latter is not undertaken until they have been removed from the thoracic cavity. During this procedure great care is to be taken to avoid tearing or pressing upon the tissues. Should there be any extensive, especially old, adhesions, these are not to be broken down, but the attached pleura at this point is to be excised at the same time. When the lungs have been removed, their surface is again to be carefully examined for recent changes, so that nothing shall be overlooked,—for example, the commencement of inflammatory exudations; then the air contents, color, and consistency of the individual lobes are to be given. Finally large, smooth sections are to be made in order to determine the appearance of the cut surface and the air, blood, and fluid contents, as well as any solid contents of the air-vesicles, the condition of the bronchi and the pulmonary arteries, the latter being examined with special care to detect any obstructions, etc. For this purpose the air-passages and the large pulmonary vessels are to be opened with scissors and their finer ramifications followed out. When the suspicion arises that foreign materials are present in the air-passages or substances are therein found the nature of which cannot with certainty be determined by the naked eye, a microscopical examination is to be made.

## Neck

§ 20. The examination of the neck may, according to the nature of the case, be made either before or after the opening of the thorax or the removal of the lungs. The obducent may also sever the larynx and the bronchus before the further inspection of the remaining parts when it seems to them especially desirable so to do, as is the case in drowning or hanging. As a rule, it is wise first to examine the large vessels and the nerve-trunks, then the larynx and trachea, by means of an anterior incision, and note their contents. If this observation should appear to be of especial importance, it is to be made before the removal of the lungs, which are at the same time to be carefully pressed upon to see if any fluid, etc., arises in the trachea. The larynx, the tongue, the velum palati, the pharynx, and the œsophagus are to be removed together; the individual parts are to be entirely opened and their contents and especially the mucosa thoroughly examined. At the same time the thyroid, the tonsils, the salivary glands, and the lymph glands of the neck are to be observed. In every case where injuries of the larynx or of the bronchus have been found or important changes therein are suspected, the air-passages are to be opened after their removal from the body and they are then to be examined from their posterior aspect. In cases of hanging or in suspicious cases of strangulation the carotids are to be opened in order to ascertain whether or not their inner coats have been injured. This examination is to be undertaken while the vessels are still in their natural situation. Finally the condition of the cervical vertebræ and of the deep musculature is to be determined.

§ 21. The abdominal cavity and its viscera are now to be critically inspected in such order that the removal of one organ does not prevent the exact determination of its relations to another. Thus, the duodenum and the gall-ducts are to be examined before the scrutiny of the liver. As a rule, the following order of examination commends itself: 1. Omentum. 2. Spleen. 3. Kidneys and adrenals. 4. Bladder. 5. Organs of generation: in the male, prostate, seminal vesicles, testicles, and penis with the urethra; in the female, ovaries, Fallopian tubes, uterus, and vagina. 6. Rectum. 7. Duodenum and stomach. 8. Gall-ducts. 9. Liver. 10. Pancreas. 11. Mesentery. 12. Small intestine. 13. Large intestine. 14. The large blood-vessels in front of the vertebral column, whose condition as to blood contents is to be ascertained and noted.

## Spleen

In every case the spleen is examined in regard to its length, breadth, and thickness, not while held in the hand, but when placed on a solid surface and without pressure by the instrument used in measuring. It is to be divided throughout its entire length, more incisions being made in different directions if diseased areas are found.

## Kidneys

Each of the kidneys is to be removed after cutting vertically through the peritoneum externally and behind the ascending or descending colon, which is shoved back. The capsule is then incised longitudinally through its convex border and slowly peeled off, and the exposed surface of the kidney is examined in regard to size, form, color, condition of blood, and other appearances. Next a longitudinal incision is made through the entire kidney to its pelvis, and the cut surfaces are washed with water and described, in which description medullary and cortical substances, vessels, and parenchyma are to be distinguished.

## Pelvic Organs

The pelvic organs (bladder, rectum, and genitalia connected therewith) are removed preferably *en masse*, the bladder being opened and its contents examined while it is still in its natural situation. After their removal these organs are again inspected, the reproductives



being examined and opened last. The slitting of the vagina is to precede that of the uterus. In puerperæ the venous and lymphatic vessels both in the internal surface of the uterus and in its walls and adnexa require special attention as to their width and contents.

When their external condition has been determined, the stomach and duodenum are with a pair of scissors opened in their natural situation, the duodenum on its anterior surface and the stomach along its greater curvature. After a careful inspection of their contents, the permeability and the presence of any matter in the opening of the gall-passages are determined and these parts are then removed for further examination.

Stomach and  
Duodenum

The liver is first described externally in its natural situation, and after its secretory ducts have been examined (as mentioned in the preceding paragraph) the gland is excised. Smooth incisions are now made through the entire length of the organ and its capacity for blood and the condition of the parenchyma determined. In the description a short account is always to be given of the general relations of the individual lobes, noting especially the condition of the inner and outer portions.

Liver

The small and large intestines, after their individual portions have been examined externally as to dimensions, color, and other peculiarities worthy of mention, are removed together, their mesenteric attachments being severed with a knife close to the bowels, which are then opened with a pair of scissors at the place where the mesentery was attached. During these incisions the contents of the several parts are observed and described. Next the intestines are cleansed and the condition of the individual portions, especially of the small intestine, is inspected with special regard to the Peyer's patches, the solitary follicles, the villi, and the intestinal folds. At least in every case of inflammation of the peritoneum the appendix is to be carefully examined.

Small and Large  
Intestines

§ 22. In those cases in which poisoning is suspected the internal examination is to begin with the abdominal cavity. Before anything else is done the external appearance of the upper abdominal viscera, their situation and extent, the filling of their vessels, and the presence of any odor are to be determined. In regard to the vessels, here as in other important organs, we are to ascertain whether we are dealing with arteries or veins, whether only the main trunks and their branches or the smaller ramifications also are filled to a given degree, and whether the extent of the vascular thinning is considerable or otherwise. Then to the portion of the œsophagus just above its entrance into the stomach and to the duodenum just below the entrance of the gall-duct double ligatures are to be applied and both parts incised between them. Next the stomach with the duodenum attached is carefully removed from the body and opened in the manner described in § 21. The contents are immediately examined as to their amount, consistency, color, composition, reaction, and odor, and placed in a clean porcelain or glass vessel. Then the mucosa is washed and its thickness, color, surface, and condition are determined, the state of the blood-vessels and the structure of the mucous membrane being particularly noted and each main portion separately described. Of especial importance is it to ascertain whether the blood which is present lies within the vessels or is exuded therefrom, whether it is fresh or changed by decomposition or by digestion, and whether in these conditions the neighboring tissues are permeated therewith. If such imbibition has occurred, it is to be noted whether it is found only upon the surface or in the tissue also, whether

Cases of Poisoning



it is coagulated or not, etc. Finally it is of especial importance to decide, in the inspection of the surface, whether loss of substance, erosions, and ulcers are present. The question whether certain changes might not have resulted from natural processes of decomposition after death, especially from the action of the fermentative juices of the stomach, is always to be considered. After the completion of this examination, the stomach and duodenum are to be placed in the same vessel with the gastric contents (see above) and given to the magistrate for further investigation. An anatomical examination having been made of the œsophagus, it is tied high up in the neck, severed above the ligature, and placed in the same vessel. In those cases in which but a small amount of stomach contents is present the contents of the jejunum are also to be preserved. Finally other substances and portions of organs, as blood, urine, pieces of the liver and of the kidney, etc., are to be removed from the body and given to the magistrate for further examination. The urine is to be placed in a separate vessel, and the blood is to be preserved separately only in those cases where spectroscopic examination might disclose facts of interest. All of the remaining portions are to be placed together in a single receptacle. Each of these vessels is closed, sealed, and labelled. In every case where the macroscopical examination shows special alteration and swelling of the mucous membrane of the stomach, a microscopical examination thereof is to be made as soon as possible, especial attention being given to the condition of the peptic glands. Whenever suspicious bodies are found in the stomach contents, as portions of leaves or other parts of plants, remnants of animal food, etc., these also are to be viewed with a microscope. Where trichinosis is suspected, not only a microscopical examination of the contents of the stomach and of the upper portion of the small intestine is to be made, but portions of muscular tissue from the diaphragm, the neck, and the thorax are also to be laid aside for future study.

The New-born;  
Determination of the  
Maturity and Period  
of Intra-uterine  
Gestation

§ 23. In postmortems on the new-born, besides the points previously given, there are to be determined, first of all, the data upon which the maturity and the intra-uterine developmental period of the child depend. For these purposes consider the length and weight of the body, condition of the general coverings and of the umbilical cord, length and appearance of the hair of the head, size of the fontanels, longitudinal, transverse, and diagonal measurements of the head, appearance of the eyes (pupillary membrane), condition of the nasal and auricular cartilage, length and characteristics of the nails, transverse diameter of the shoulders and hips; in boys the condition of the testicles and the appearance of the scrotum, and in girls any peculiarities of the external genitalia. It still remains to be noted whether there be present, and if so to what extent, an ossifying centre in the inferior epiphysis of the femur. To determine this the patella is removed through a horizontal incision made just below it while the knee-joint is strongly flexed, and thin transverse sections are made continuously through the cartilage until the greatest transverse diameter of any centres of ossification which may there be present is found, which is then to be measured in millimetres. When from an examination of the offspring it seems to have been born before the thirtieth week, the postmortem may be discontinued unless a special request is given by the magistrate for its completion.

The Determination  
as to whether or not  
the Child has  
breathed

§ 24. If it be determined that the child was born after the thirtieth week, the following data must be obtained in order to decide whether it breathed during or after birth. For this purpose the respiratory tests are to be applied in the following order:  
(a) Immediately after the opening of the abdominal cavity the

condition of the diaphragm in relation to the corresponding ribs is to be determined. Hence in every case of examination of the new-born the abdominal cavity is to be opened first and afterwards the thoracic and cranial cavities.<sup>1</sup> (b) Before opening the thoracic cavity the trachea is to be once ligatured above the sternum. (c) The thoracic cavity is next to be opened and the extent and consequent situation of the lungs, the latter especially in regard to the pericardium, determined, and also the color and consistency. (d) The pericardium is to be incised and both its condition and the external appearance of the heart are to be described. (e) The individual cavities of the heart must be laid open, their contents noted, and other appearances determined. (f) The larynx and the portion of the trachea above the ligature are to be slit, and their contents as well as the appearance of their walls determined. (g) The trachea is to be cut through above the ligature and removed in connection with the other organs of the thorax. (h) After the removal of the thymus gland and the heart, the lungs are to be tested as to whether or not they float in a large vessel filled with pure cold water. (i) The lower portion of the bronchus and its branches are to be opened and their contents specially examined. (j) Incisions are to be made into both lungs, the presence or absence of crepitation being carefully noted as well as the amount and appearance of any blood which may exude under slight pressure upon the cut surfaces. (k) The lungs are also to be incised under water in order to determine if any air-bubbles arise from the cut surfaces. (l) The lobes of both lungs are next to be cut apart, each lobe subdivided, and every separate portion tested as to its sinking or floating in water. (m) The œsophagus is to be opened and its condition ascertained. (n) Finally, in those cases where it is suspected that the pulmonary tissues may have been filled with the products of disease (hepatization) or with foreign bodies (vernix caseosa and meconium), so as not to permit of the entrance of air, the same are to be examined microscopically.

§ 25. Lastly, it is the duty of the obducent to examine all organs not mentioned in these regulations in case injuries or other abnormalities are discovered.

Further  
Examinations

§ 26. The district surgeon, with the second physician acting as a consultant, is required, after the ending of the autopsy and as far as possible the removal of waste, to undertake the proper closure of those cavities of the body which have been opened.

Closure of the Body

### III. THE DRAWING UP OF THE PROTOCOL OF THE POSTMORTEM AND THE FINAL REPORT OF THE SAME.

§ 27. A post-mortem protocol is to be made by the magistrate, at the time and place of performing the autopsy, concerning all matters relating thereto. The medical officer must, therefore, be careful that the technical findings which have been determined at the examination are faithfully recorded in the protocol. In order to accomplish this, it is recommended to the magistrate that the description and findings of each individual organ be written down before another part is examined.

The Post-mortem  
Protocol

<sup>1</sup> But in no case shall section of the organs of the abdominal cavity be undertaken before the opening and examination of those of the thorax.

Arrangement and Form of the Protocol      § 28. The technical findings given in the post-mortem protocol by the medical officer must be stated clearly, definitely, and in such a manner as to be understood by one who is not a physician. for this purpose the use of foreign expressions is to be avoided except where these may be needed to make clear the description of the findings. The chief divisions, the external and internal examinations, are to be designated with capital letters (A and B). The findings for the openings in the cavities are to be given, in the order in which they were examined, with Roman numerals (I, II.); but the organs in the thorax and abdominal cavity are to be entered under a single number. The descriptions of the organs of the thorax and abdominal cavity, named in § 18, are to be designated by the letters a and b. The results of the examination of each individual part are to be designated with Arabic numerals, such numbers running consecutively from the beginning to the end of the protocol. The record of the examination must be given in the protocol with special reference to the actual observations, and not in the form of mere statements of opinion,—as, for example, inflamed, gangrenous, healthy, normal, wound, ulcer, and the like. The obducent has the option, however, in those cases in which it seems necessary for clearness, to add such observations, inclosed in parentheses. In every case a note must be made of the blood contents of each important part, and a short description thereof must be given, and not simply a name,—as considerable, moderate, middling amount, much reddened, rich in blood, poor in blood. Before any part is incised its size, form, color, and consistency are to be noted, in the order here named.

Provisional Opinion      § 29. At the close of the postmortem the obducent is to give in the protocol their provisional opinion of the case, without stating their reasons therefor. If anything be known by means of which the diagnosis is influenced, in the way of previous history or the like, this must be briefly noted. Should the magistrate ask any special questions, the answers should be distinctly entered in the protocol, with the statement that they are given at his request. In every case the opinion as to the cause of death is to be stated, first with special reference to the facts bearing on the objective findings and then as to the question of criminal motive. If the cause of death is not determined, this fact must be recorded. It is never sufficient to say that death resulted from internal causes or from disease. The latter, whatever it is, must be specifically named. Special mention is to be made, with the reason therefor, in cases where further technical examinations are needed or where doubtful conditions exist.

Supplemental Observations on Instruments      § 30. Should injuries be found on the body which were presumably the cause of death, and if suspicion be aroused that a specially discovered instrument might have inflicted such injuries, the obducent, at the request of the magistrate, are obliged to investigate and to express an opinion as to whether any and, if so, what injuries might have been caused by the instrument, and what conclusions from the situation and appearance of the wound are to be drawn as to the manner in which the one performing the act might have committed the deed, and also as to the strength with which it was performed. When definite weapons are not found, the obducent, as far as it is possible from the conditions present, are to give their opinion as to how the injuries were caused and especially as to what instruments might possibly have been used.



§ 31. If the obducents be requested to present a report, this should be introduced without useless formalities by a condensed but Post-mortem Report exact review of the case, with the conclusion reached by them and the facts on which it is based. Then so much of the post-mortem protocol as they think necessary for the explanation of the case is to be given verbatim, with the number of the protocol. Any change made therein must be expressly stated. The style of the post-mortem report must be plain and concise, and the proof which led to the formation of the opinion therein expressed so set forth as to be understood by and convincing to one not a physician; for this purpose, the obducents are to use, as far as possible, German expressions and ordinarily accepted meanings. Especial attention to literary sources of knowledge is, as a rule, to be avoided. When as medical experts the obducents are asked certain questions by the magistrate, these are to be answered fully and as directly as possible, or, if this cannot be done, the reasons therefor are to be given.

Both obducents must sign their report, which must also bear the official seal of the district physician if he has taken part in the autopsy. When such a post-mortem account is requested, it must be delivered by the obducents within four weeks at the latest.

## CHAPTER XXVIII

### USUAL CAUSES OF DEATH; THEIR NOMENCLATURE, COMPLICATIONS, AND SYNONYMS

As morbidity and mortuary statistics are intimately associated the one with the other, uniformity in their nomenclature throughout the world is greatly to be desired. At the Eighth International Congress of Hygiene and Demography, held in Paris, August 18 to 21, 1900, a modification of the old Bertillon classification was adopted and called the "International System of Nomenclature of Diseases and Causes of Death."<sup>1</sup> It is here added but slightly altered in a few minor particulars as to the causes of death.

#### I. GENERAL DISEASES.

1. TYPHOID FEVER (Abdominal Typhus). *Include:* Dothienenteritis; mucous, continued, enteric, ataxic, or adynamic fever; abdominal typhus.—*Do not include:* Adynamia (179); ataxo-adynamia (179).—*Frequent complications:* Pneumonia; pulmonary congestion; intestinal perforation; peritonitis; intestinal hemorrhage; sloughing; albuminuria.
2. EXANTHEMATOUS TYPHUS.<sup>2</sup> *Include:* Petechial fever; petechial typhus.—*Do not include:* Abdominal typhus; typhus.
3. RELAPSING FEVER. *Include:* Recurrent fever; recurrent typhus.
4. INTERMITTENT FEVER AND MALARIAL CACHEXIA. *Include:* Paludal fever; pernicious fever; *accesso pernicioso*; remittent fever; malaria; ague; etc.
- 4a. MALARIAL CACHEXIA. *Include:* Paludism; pernicious cachexia; etc.
5. SMALLPOX. *Include:* Variola, varioloid.—*Do not include:* Varicella (19).—*Frequent complications:* Meningitis; endocarditis; suppuration; albuminuria.
6. MEASLES. *Include:* Eruption of measles; morbilli.—*Do not include:* Rubeola or German measles (19).—*Frequent complications:* Bronchitis; bronchopneumonia.
7. SCARLATINA. *Include:* Puerperal scarlatina; scarlatinous angina.—*Frequent complications:* Albuminuria; eclampsia; œdema of the glottis; hemorrhage; endocarditis; pericarditis; paralysis; convulsions; diphtheria.
8. WHOOPING COUGH (PERTUSSIS). *Frequent complications:* Bronchitis; spasms.
9. DIPHTHERIA AND CROUP. *Include:* Diphtheritic, buffy, pseudomembranous, infectious, malignant, or toxic angina. Diphtheria under all its forms, especially diphtheria of wounds, cutaneous diphtheria; conjunctival diphtheria;

<sup>1</sup> Supplement to Public Health Reports, vol. xv, no. 49. Translated by Passed Assistant Surgeon H. D. GEDDINGS. WILLIAM A. KING, chief statistician of vital statistics of the United States Census, has added to this list the indefinite and unsatisfactory synonyms used for causes of death in the returns to the United States Census Office for 1890 and 1900. Washington, 1902.

<sup>2</sup> The word "typhus," without qualification, will be taken in the sense which is usual to it in each country,—viz., in the sense of "abdominal typhus" in German-speaking countries, or as "exanthematous typhus" in French-speaking ones.

buccal diphtheria; pseudomembranous bronchitis; pseudomembranous laryngitis; malignant laryngitis; diphtheritic paralysis, etc.—*Do not include:* Stridulous croup (88); spasmodic croup (88); false croup (88).—*Frequent complications:* Pneumonia; albuminuria; paralysis.

- 9a. DIPHTHERIA.
10. INFLUENZA. *Include:* Grippe; grippe pneumonia; grippe bronchitis; epizootic; and grippe bronchopneumonia.
11. SWEATING OR MILIARY FEVER. *Include:* Sudor.
12. ASIATIC CHOLERA. *Include:* Indian cholera; cholera (when epidemic); epidemic cholera.
13. CHOLERA NOSTRAS.<sup>1</sup> *Include:* Sporadic cholera; cholerine; choleriform enteritis or diarrhoea; cholera (when not epidemic).—*Do not include:* Cholera infantum (105 or 106); antimonial cholera (175); hernial cholera (108).
14. DYSENTERY. *Include:* Choleriform dysentery; Chinese dysentery; dysentery of tropical countries.
- 14a. EPIDEMIC DYSENTERY.
15. BUBONIC PLAGUE (Plague or pest).
16. YELLOW FEVER. *Include:* Vomito negro; amarilla fever; black vomit.
17. LEPROSY. *Include:* Elephantiasis Græcorum.—*Do not include:* Elephantiasis Arabum (145d); Morvan's disease (63); syringomyelitis (63).
18. ERYSIPELAS. *Include:* All surgical erysipelas or medical erysipelas, without regard to seat; St. Anthony's fire.—It is disputed whether to classify gangrenous or phlegmonous erysipelas here or under 144.
19. OTHER EPIDEMIC AFFECTIONS.<sup>2</sup> *Include:* Mumps; rubeola; acrodynia; varicella; beriberi; and any other epidemic affections which may not be included in this nomenclature.—*Do not include:* Epidemic dysentery (14a); epidemic cerebrospinal meningitis (61).
20. PURULENT AND SEPTICÆMIC INFECTION.<sup>3</sup> *Include:* Pyohæmia; purulent absorption; putrid absorption; putrid infection; putrid fever; anatomic (dissection) wounds; streptococ hæmia.—*Do not include:* Puerperal septicæmia (137); infectious fever (55).
21. GLANDERS AND FARCY.
22. MALIGNANT PUSTULE. *Include:* Charbon; splenic fever. In France as 143.
23. RABIES. *Include:* Hydrophobia.—*Do not include:* Sitiophobia (68).
24. ACTINOMYCOSIS, TRICHINOSIS, ETC. *Include:* Dystoma hepaticum; cysticerci.—*Do not include:* Cyst or hydatid tumor of the liver (111) or of the lungs (99); intestinal parasites (107).
25. PELLAGRA.
26. TUBERCULOSIS OF THE LARYNX. *Include:* Tuberculous laryngitis; laryngeal phthisis.
27. TUBERCULOSIS OF THE LUNGS.<sup>4</sup> *Include:* Pulmonary tuberculosis; pulmonary phthisis; phthisis (without qualification); phymia; phymatosis; pneumio-phymia; acute, galloping, or miliary phthisis or tuberculosis; pulmonary cavities; consumption; caseous pneumonia; tuberculous, bacillary, specific,

<sup>1</sup> The word "cholera morbus" will be taken in its ordinary signification in each country, as in the sense of "cholera nostras" in North America, and as "Asiatic cholera" in France and in other countries.

<sup>2</sup> In cases where epidemics arise, it will be necessary here to adopt a special provisional title.

<sup>3</sup> When a female of childbearing age is returned as affected with "septicæmia," "sepsis," or any similar term, send the report back in order that the physician may state whether or not the disease was puerperal.

<sup>4</sup> See observation on No. 93, relative to "apical pneumonia."



- granular, neoplastic, or heteroplastic bronchitis or pneumonia; bacillosis; tuberculous pleurisy; tuberculous hæmoptysis; tuberculosis (without qualification).—*Do not include:* Hæmoptysis (without qualification) (99); pulmonary hemorrhage (99); bronchorrhagia (without qualification) (99); apical pneumonia (93); laryngeal phthisis (26); pulmonary anthracosis (99).—*Frequent complications:* Hemorrhage; pneumonia; pleurisy; uncontrollable diarrhœa.
28. TUBERCULOSIS OF THE MENINGES. *Include:* Meningeal tuberculosis; tuberculous meningitis; granular, miliary, caseous, bacillary, specific, neoplastic or heteroplastic meningitis; tuberculous meningitis of spinal cord.—*Do not include:* Meningitis (without qualification), even for children of tender age (61).
29. ABDOMINAL TUBERCULOSIS. *Include:* Tuberculous, granular, bacillary, or specific peritonitis; peritoneal tuberculosis; tuberculous enteritis.
30. POTT'S DISEASE. *Include:* Vertebral caries; vertebral polyarthritis.—*Frequent complications:* Cold abscess, or abscess by congestion; paraplegia.
31. COLD ABSCESS (ABSCESS BY CONGESTION). *Include:* Ossifluent abscess.
32. WHITE SWELLING. *Include:* Fungous growths of joints; coxalgia; scapulargia; tuberculosis of joint.
33. OTHER TUBERCULOUS AFFECTIONS. *Include:* Tuberculosis of the skin, eye, bone, genital organs, etc.; tuberculous adenitis; lupus; esthiomene; bacillary abscess; tuberculous ulcer.—*Do not include:* Pott's disease (30).
34. GENERALIZED TUBERCULOSIS. *Include:* Tuberculosis showing itself simultaneously in any two or more organs. Often better placed under 27.
35. SCROFULA. (An unsatisfactory title.) *Include:* Struma; King's evil; lymphatism; scrofulides.—*Do not include:* Scrofulous or lymphatic keratitis and blepharitis (75).
36. SYPHILIS. Of which are recognized: (1) Primary, (2) secondary, (3) tertiary, (4) hereditary. These divisions are intended for mortuary statistics alone. *Include:* (1) Indurated or infecting chancre; chancre of the mouth or face; primary accident or infection; (2) Secondary manifestations—mucous plaques; syphilitic amygdalitis; angina or laryngitis; (3) Tertiary manifestations—specific manifestations; gummata; ulcerations; exostoses, etc. Osteocopic pains; all these diseases to be specified as "syphilitic."—*Do not include:* Soft, simple, or phagedenic chancre (36a).
- 36a. SOFT CHANCRE. *Include:* Chancroid; chancrelle; simple chancre; phagedenic chancre or bubo; bubo of soft chancre; venereal, virulent, or absorption buboes.—*Do not include:* Infecting or syphilitic chancre or bubo (36, 1); chancre of the mouth (36, 1); scrofulous bubo (35); suppurating bubo (144); plague bubo (15); bubo (without qualification) (144). (Morbidity statistics only.)
37. GONORRHOEA (five years and over). *Include:* Blennorrhœa; ardor urinæ; urethritis; military drop; balanitis; balanorrhagia; balanoposthitis, vaginitis; gonorrhœal cystitis, orchitis, buboes, arthritis, rheumatism, or conjunctivitis of the adult; or gonorrhœal or blennorrhagic ophthalmia of the adult.—*Do not include:* Vaginismus (132); vaginalitis (126).—*Frequent complications:* Bubo; adenitis; cystitis; orchitis.
38. GONORRHOËAL AFFECTIONS OF THE CHILD.<sup>1</sup> *Include:* Blennorrhagic or gonorrhœal conjunctivitis of the child; gonorrhœal vulvitis, etc.

<sup>1</sup> This title takes account only of children under five years of age.

39. CANCER AND OTHER MALIGNANT TUMORS OF THE BUCCAL CAVITY. *Include:* Cancer of the mouth, lips, tongue, roof of the mouth, velum of palate, maxilla, jaw, parotid gland, and tonsil; sarcoma of soft palate; epithelioma, or carcinoma, or canceroid of these organs; smoker's cancer.
40. CANCER AND OTHER MALIGNANT TUMORS OF THE STOMACH AND LIVER.<sup>1</sup> *Include:* Cancer of the œsophagus; cancer of the cardia; cancer of the pylorus; carcinoma or scirrhus, or colloid or encephaloid tumor of these organs; gastrocarcinoma; tumor of the stomach; scirrhus of liver or stomach; hepatic cancer.—*Do not include:* Hæmatemesis (104).
41. CANCER AND OTHER MALIGNANT TUMORS OF THE PERITONEUM, INTESTINES, AND RECTUM. *Include:* Cancer, carcinoma, scirrhus, encephaloid, canceroid, or epithelioma of the colon and anus; retroperitoneal sarcoma.
42. CANCER AND OTHER MALIGNANT TUMORS OF THE FEMALE GENITAL ORGANS. *Include:* Cancer of the uterus (womb), ovary, vagina, vulva; carcinoma, or encephaloid, or colloid tumor, or heteromorphous or neoplastic growth, or canceroid, or sarcoma, or epithelioma of these organs.
43. CANCER AND OTHER MALIGNANT TUMORS OF THE BREAST. *Include:* Carcinoma, or scirrhus, or encephaloid, or heteromorphous or neoplastic growth, or canceroid, or epithelioma of the breast or nipple.
44. CANCER AND OTHER MALIGNANT TUMORS OF THE SKIN. *Include:* Canceroid (without qualification); epithelioma or epitheliomatous tumor (without qualification); cancer of the ear, face, nose, or cervicofacial region; "noli-me-tangere;" rodent ulcer.—*Do not include:* Esthiomene (33); lupus (33).
45. CANCER AND OTHER MALIGNANT TUMORS OF OTHER ORGANS, OR OF ORGANS NOT SPECIFIED. *Include:* Abdominal cancer; pelvic cancer; cancer of the lung, of the kidney, of the bladder, and of the prostate; cancerous goitre; thyro-sarcoma; sarcohydrocele; cancer of the bone; osteosarcoma; cancerous or sarcomatous tumor of the neck; carcinoma, or scirrhus, or encephaloid, or cancerous ulcer, or malignant tumor, or sarcoma, or malignant fungus of these organs, or of other organs not specified; chimney sweeps' cancer; Lobstein's cancer; fungus hæmatodes; sarcoma of leg; lymphosarcoma, etc.—*Do not include:* Cancer of the œsophagus (40); cancer of the anus (41); cancer of the ovary, vagina, or vulva (42).
46. OTHER TUMORS (Tumors of the Female Genital Organs excepted). *Include:* Tumor (without qualification); abdominal tumor; intestinal tumor; vascular or erectile tumor; angioma; lymphoma; lymphadenoma; lymphatocele; adenoma; chondroma; osteoma; myoma; lipoma; wen; grub; sebaceous tumor; cystoma.—*Do not include:* Cancer and its synonyms (39-45); tumor of the stomach (40); stercoraceous tumor (108); tumor of the uterus (129); hydatid tumor (111); cyst of the ovary (131); aneurismal tumor (81); varicose tumor (83); polyp of the ear (76); polyp of the nasal or nasopharyngeal fossæ (87); uterine polyp (129); osteoma (146).
47. ACUTE ARTICULAR RHEUMATISM. *Include:* Rheumatic arthritis; rheumatic meningitis; abdominal or cerebral rheumatism; rheumatic vertigo; rheumatic endocarditis, pericarditis, pleurisy, peritonitis, etc.—*Do not include:* Organic heart disease of rheumatic origin (79); rheumatic iritis (75) arthritis deformans (48); gonorrhœal rheumatism (37 and 38).

<sup>1</sup> In countries where the words "organic lesion of the stomach" always signify "cancer of the stomach" classify these diagnoses under No. 40. In countries where, on the contrary, this is not always so, classify them under No. 104.

48. CHRONIC RHEUMATISM AND GOUT. *Include:* Arthritis deformans; articular rheumatism; Heberden's disease, podagra; rheumatic gout; rheumatism (unqualified).
49. SCURVY. *Include:* Scorbutus; Werlhoff's disease; Barlow's disease.
50. DIABETES. *Include:* Diabetes insipidus and mellitus; Hirshfeld's disease; diabetic gangrene and coma; glycosuria.—*Frequent complications:* Pneumonia; furunculosis; gangrene; cerebral hemorrhage and cerebral softening; tuberculosis.
51. EXOPHTHALMIC GOITRE. *Include:* Exophthalmia; Basedow's, Graves's, Parry's disease; exophthalmic cachexia.—*Frequent complications:* Hypertrophy of the heart; cachexia.
52. ADDISON'S DISEASE. *Include:* Adrenal disease. *Freq. complic.:* Cachexia; ascites.
53. LEUKÆMIA. *Include:* Adenoleukæmia; leucocythæmia; Hodgkin's disease; pseudoleukæmia.—*Freq. complic.:* Hemorrhage; ascites; apoplexy; cachexia.
54. ANÆMIA; CHLOROSIS. The cause of the anæmia should always be given. *Include:* Pernicious anæmia.—*Do not include:* Cerebral anæmia (74b).
55. OTHER GENERAL DISEASES. *Include:* Autointoxication; infectious fever; virulent disease (without explanation); visceral steatosis; acromegalia; amyloid or generalized fatty degeneration; adiposis; obesity; polysarcia.
56. ALCOHOLISM, ACUTE OR CHRONIC. *Include:* Drunkenness; ethylism; alcoholic intoxication; alcoholic delirium; alcoholic dementia; delirium tremens; absinthism; absinthæmia; dipsomania; "mania-a-potu."—*Do not include:* Alcoholic cirrhosis (112); general alcoholic paralysis (67); atheroma (81); or any other disease attributable to alcohol; intoxication amblyopia (75).
57. SATURNISM. *Include:* Saturnine colic; lead colic; painter's colic; lead encephalopathia; lead paralysis; chronic lead poisoning; all conditions characterized as "saturnine."
58. OTHER TRADE OR OCCUPATION INTOXICATIONS. *Include:* Mercurial (hydrargyris); phosphorus, arsenical, or other intoxication, when special mention by the physician makes it clear that the intoxication is the result of a trade. Failing in this specific declaration, it should be classed in one of the conditions under No. 59.—*Do not include:* alcoholism (56).
59. OTHER CHRONIC POISONINGS. *Include:* Morphinism; cocaineism; chronic ergotism.—*Do not include:* Amblyopia from intoxication (75). Note the remark under No. 58.

## II. DISEASES OF THE NERVOUS SYSTEM AND OF THE ORGANS OF SPECIAL SENSE.

60. ENCEPHALITIS. *Include:* Cerebral fever; phrenitis; poli-encephalitis.
61. SIMPLE MENINGITIS. *Include:* Meningitis (without qualification); meningo-encephalitis; pachymeningitis.
- 61a. EPIDEMIC CEREBROSPINAL MENINGITIS. *Do not include:* Tuberculous meningitis (or other synonym) (28); rheumatic meningitis (47).
62. PROGRESSIVE LOCOMOTOR ATAXIA. *Include:* Duchenne's disease; posterior sclerosis; tabes dorsalis; cerebral ataxia; posterior spinal sclerosis; progressive ataxia; progressive spastic ataxia.
63. OTHER DISEASES OF THE SPINAL CORD. *Include:* Disease of the cord; sclerosis in plaques; symmetrical sclerosis; lateral sclerosis; sclerosis (without qualification); Charcot's disease; Morvan's disease; syringomyelitis; hemorrhage into the spinal cord; hæmatomyelitis; hæmatorrhachia; myelitis;



- medullary congestion; affections of the bulb; bulbar paralysis; spinal paralysis; paralysis agitans; trembling paralysis; ascending paralysis; essential paralysis of infancy; fatty or amyloid degeneration of the cord; Parkinson's disease; Friedreich's disease; medullary compression or compression of the cord; progressive muscular atrophy; fatty degeneration of muscle; atrophic muscular paralysis; amyotrophia; amyotrophic paralysis; atrophic paralysis; pseudohypertrophic paralysis; etc.
64. CEREBRAL CONGESTION AND HEMORRHAGE. *Include:* Apoplexy; cerebral apoplexy; meningeal apoplexy; serous apoplexy; cerebral atheroma; œdema of the brain; cerebral effusion; cerebellar hemorrhage; meningeal hemorrhage; cataplexia; apoplectic dementia; stroke (unqualified); clot on brain.—*Frequent complications:* Hemiplegia; paralysis.
65. CEREBRAL SOFTENING. *Do not include:* Senile dementia (154).—*Frequent complications:* Hemiplegia; paralysis; pulmonary congestion.
66. PARALYSIS WITHOUT SPECIFIED CAUSE. *Include:* Paralysis (without qualification); hemiplegia; facial paralysis; generalized paralysis (not insane or unqualified); palsy.—*Do not include:* Diphtheritic paralysis (9); atrophic muscular paralysis (63); general paralysis (67); paralytic cachexia (marasmus) (67); paralytic dementia (idiocy) (67); shaking (67) or trembling paralysis (63); bulbar paralysis (63); ascending paralysis (63); essential paralysis of infancy (63); labioglossolaryngeal paralysis (74b); paralysis of the velum palati (101); paralysis of the muscles of the eye (75).
67. GENERAL PARALYSIS. *Include:* Paralytic lunacy; paralytic dementia; paralytic cachexia; paralytic marasmus; diffuse meningo-encephalitis; diffuse peri-encephalitis.—*Do not include:* Generalized paralysis (not insane) (66).
68. OTHER FORMS OF MENTAL ALIENATION. *Include:* Dementia; lunacy; unsoundness of mind; hallucinations; mania; megalomania; monomania; delusions of persecution; melancholia; lypomania; nostalgia; insanity; nosophobia; necrophobia; sitiophobia; lycanthropy; homesickness; andromania; nymphomania; priapism; satyriasis; mental disease.—*Do not include:* Alcoholic dementia or delirium (56); delirium tremens (56); delirium (179); uræmic delirium (120); apoplectic dementia (64); paralytic dementia (67); choreic dementia (73); senile dementia (154); hysteria (74a).
69. EPILEPSY. *Include:* "Haut and petit mal;" disease of Hercules; fits; falling sickness.—*Do not include:* Epileptiform convulsions (70-71).
70. ECLAMPSIA (Non-puerperal).<sup>1</sup> *Include:* Epileptiform convulsions (of adults).—*Do not include:* Scarlatinous eclampsia (7); uræmic eclampsia (120); eclampsia of children under five years of age (71).
71. CONVULSIONS OF CHILDREN.<sup>2</sup> *Include:* Eclampsia of young children; contractions of children; spasms.—*Do not include:* Trismus nascentium (72).
72. TETANUS. *Include:* Opisthotonos; emprosthotonos; pleurosthotonos; trismus nascentium or neonatorum; lockjaw; idiopathic tetanus.
73. CHOREA. *Include:* Choreic dementia; Bergeron's disease; St. Vitus's dance.
74. HYSTERIA. *Include:* Hysterical anorexia; hysterical colic; all diseases classified as "hysterical." (Morbidity statistics alone.)
- 74a. NEURALGIA. *Include:* Tic douloureux; sciatica. (Morbidity statistics alone.)

<sup>1</sup> When a female of childbearing age is designated as having been stricken with "eclampsia," return the report to have the physician state whether or not the disease was puerperal.

<sup>2</sup> This title applies only to children under five years of age.

- 74b. **OTHER DISEASES OF THE NERVOUS SYSTEM.** *Include:* Cerebral compression; cerebral tumor; acquired hydrocephalus; neuroma; encephalopathia (without qualification); idiocy; imbecility; cretinism; gatism; amnesia; paramnesia; loss of speech; aphasia; nervous or cerebral accidents; cerebral anæmia; neurosis; tic; convulsive tic; contracture; anæsthesia (not due to external anæsthetic); neurasthenia; migraine; vertigo; somnambulism; catalepsy; boulimia; Landry's disease; symptomatic or Jacksonian epilepsy; athetosis; labioglossolaryngeal paralysis; amyloid or fatty degeneration of the nervous system, etc.—*Do not include:* Senile dementia, imbecility or senile gatism (154); syringomyelitis (63); myxœdema (89); congenital or undescribed hydrocephalus (150).
75. **DISEASES OF THE EYE AND ITS ADNEXA.** *Include:* Ophthalmia; foreign bodies; conjunctivitis (not including diphtheritic or gonorrhœal conjunctivitis); xerosis; pterygion; pinguecula; keratitis of every description; staphyloma; diseases of the cornea; arcus senilis; diseases of the sclerotic; diseases of the iris; iritis; diseases of the choroid; choroiditis; iridochoroiditis; sclerochoroiditis; glaucoma; diseases of the retina; retinitis; optic neuritis; amaurosis; amblyopia; amblyopia from intoxication; hemiopia; hemeralopia; nyctalopia; diseases of the lens; cataract; aphacia; parasites of the eye; ophthalmozoa; coloboma; strabismus; strabotomy; paralysis of the muscles of the eye; nystagmus; styes; chalazion; blepharitis; blepharconjunctivitis; scrofulous blepharitis; blepharophimosis; blepharoplastia; ectropion; entropion; trichiasis; dacryoadenitis; diseases of the lachrymal gland and lachrymal sac; dacryocystitis; dacryolithiasis; dacryoma; lachrymal fistula; diseases and tumors of the orbit (undefined).—*Do not include:* Diphtheritic (9) or gonorrhœal (37-38) conjunctivitis; ocular cancer (45) or tuberculosis (33); exophthalmic goitre (51); exophthalmia (51). Many titles in 75 are never employed as causes of death.
- 75a. **FOLLICULAR CONJUNCTIVITIS.** (Morbidity statistics alone.)
- 75b. **TRACHOMA.** (Morbidity statistics alone.)
76. **DISEASES OF THE EAR.** *Include:* Otitis; otorrhœa; catarrh of the ear; hydrotitis; foreign body in the auditory canal; obstruction of the auditory canal; polyp of the ear; inflammation of the tympanum; "vertigo ab aure laeso." Ménière's disease, or vertigo; caries of the labyrinth (?); deafness; deaf-mutism.—*Do not include:* Mumps (19).

### III. DISEASES OF THE CIRCULATORY APPARATUS.

77. **PERICARDITIS.** *Include:* Cardiopericarditis; hydropericarditis; hydropneumopericarditis; pericardial adhesions.—*Do not include:* Rheumatic pericarditis (47); endopericarditis (78); pleuropericarditis (94); pneumopericarditis (93).
78. **ACUTE ENDOCARDITIS.** *Include:* Endocarditis (without qualification); myocarditis, acute or without qualification; endopericarditis.—*Do not include:* Rheumatic endocarditis (47), or the other cardiac accidents which may supervene in the course of an attack of rheumatism.
79. **ORGANIC DISEASES OF THE HEART.** *Include:* Aortic, mitral, tricuspid, or cardiac affection or lesion; cardiac or valvular insufficiency or stenosis of the valves of the heart; cardiac cachexia; hypertrophy of the heart; dilatation of the heart; cardiectasis; steatosis of the heart; degeneration of the heart; cardiopathy; cardiosclerosis; cardiovascular sclerosis; cardiomalacia; cardiostenosis; labored heart; tachycardia; rupture of the heart; card

- rhexia; cardiac palpitation; asystole; cardiac asthma.—*Do not include:* Cardiac accidents (undetermined) (86); persistence or patency of the foramen of Botallo (150).—*Frequent complications:* Dropsy; bronchitis and pneumonia; albuminuria; embolism; thrombosis.
80. **ANGINA PECTORIS.** *Include:* Cardialgia; sternalgia; neuralgia of the heart.
81. **AFFECTIONS OF THE ARTERIES, ATHEROMA, ANEURISM, ETC.** *Include:* Arteritis; fatty degeneration of arteries; arteriosclerosis; atheroma of arteries; arteriectasis; aortic ectasis; Hodgson's disease; atresia of the pulmonary artery; aortitis; aneurismal tumor.—*Do not include:* Aortic affection (79).
82. **EMBOLISM AND THROMBOSIS.** *Include:* Thrombosis (without qualification); phlegmasia alba dolens (non-puerperal); embolic pneumonia.—*Do not include:* Embolism (puerperal) (140).
83. **AFFECTIONS OF THE VEINS (Varices, Hemorrhoids, Phlebitis, etc.).** *Include:* Pneumophlebitis; varicose ulcer; varicocele.—*Do not include:* Puerperal phlebitis (137); vascular or erectile tumor (46); angioma (46).
84. **AFFECTIONS OF THE LYMPHATIC SYSTEM.** *Include:* Angioleucitis; adenopathy; lymphangitis; adenitis.—*Do not include:* Suppurative adenitis (144); adenophlegmon (144); leucemic adenitis (53); lymphatism (35); bubo (36a and 144); adenoma (46); lymphoma (46); lymphadenoma (46).
85. **HEMORRHAGES.** *Include:* Hemorrhage (without qualification); internal hemorrhage; hæmophilia; epistaxis; stomatorrhagia; cutaneous hemorrhage; purpura hæmorrhagica.—*Do not include:* Cerebral hemorrhage (64); cerebellar hemorrhage (64); meningeal hemorrhage (64); pulmonary hemorrhage (99); hæmoptysis (99); hæmatemesis (104); intestinal hemorrhage (109); hæmaturia (121); uterine hemorrhage (135 or 128, depending on whether it is or is not puerperal); metrorrhagia (128 or 135); umbilical hemorrhage (under three months) (152); traumatic hemorrhage (166).
86. **OTHER AFFECTIONS OF THE CIRCULATORY APPARATUS.** *Include:* Cardiac accidents (undetermined); angiectasis; angiectopia; affections of the great vessels; permanently slow pulse.—*Do not include:* Vascular nævus (150).

#### IV. DISEASES OF THE RESPIRATORY APPARATUS.

87. **DISEASES OF THE NASAL FOSSÆ.** *Include:* Coryza; cold; polypus of the nasal or nasopharyngeal fossa; ozæna; abscess of the nasal fossa; adenoid vegetations.—*Do not include:* Epistaxis (85); syphilitic coryza (36).
88. **AFFECTIONS OF THE LARYNX.** *Include:* Acute, chronic, erysipelatous, œdematous, phlegmonous, or stridulous laryngitis; aphonia; loss of voice; false croup; spasmodic croup; stridulous croup; œdema of the glottis; spasm of the glottis; polypus of the larynx; stricture of the larynx; laryngotomy.—*Do not include:* Tuberculous laryngitis (26); laryngeal tuberculosis (26); croup (9); diphtheritic laryngitis and its synonyms (9); foreign bodies in the larynx (176).
89. **AFFECTIONS OF THE THYROID BODY.** *Include:* Goitre; thyrocele; myxœdema; pachydermic cachexia.—*Do not include:* Exophthalmic goitre (51).
90. **BRONCHITIS, ACUTE.** *Include:* Capillary bronchitis; tracheitis; tracheobronchitis; broncho-œvcolitis.—*Do not include:* Bronchopneumonia (92); specific bronchitis or other synonym of pulmonary tuberculosis (see No. 27); fetid bronchitis (96); summer bronchitis (99).



91. BRONCHITIS, CHRONIC.<sup>1</sup> *Include:* Mucous bronchitis (pituitous); catarrh (without qualification); bronchial, pituitous, pulmonary, or suffocating catarrh; bronchorrhœa; dilatation of the bronchi; bronchiectasis.—*Do not include:* Fetid bronchitis (96); tuberculous bronchitis (27).
92. BRONCHOPNEUMONIA. *Include:* Catarrhal, deglutition, and aspiration pneumonia.—*Do not include:* Capillary bronchitis (90).
93. PNEUMONIA.<sup>2</sup> *Include:* Croupous pneumonia; fluxion of the lung; pleuropneumonia; pneumopleurisy; splenopneumonia; apical pneumonia; peripneumonia; pneumopericarditis; typhoid and alcoholic pneumonia.—*Do not include:* Caseous pneumonia (27); specific, bacillary, or any synonym of pulmonary tuberculosis (27); pulmonary congestion (95).
94. PLEURISY. *Include:* Pleuropericarditis; pleuritic or thoracic effusion; pneumothorax; hydropneumothorax; pyothorax; pleural vomica; pneumopyothorax; hæmothorax; thoracentesis; empyema; pleural adhesions.—*Do not include:* Pleurodynia (99).
95. PULMONARY CONGESTION AND PULMONARY APOPLEXY. *Include:* CEdema of the lungs; hypostatic pneumonia.
96. GANGRENE OF THE LUNG. *Include:* Fetid bronchitis; mortification of lung.
97. ASTHMA. *Do not include:* Cardiac asthma (79); suffocating catarrh (91); hay fever (99).
98. EMPHYSEMA OF THE LUNGS. *Include:* Emphysema (without qualification).—*Do not include:* Subcutaneous emphysema (145).
99. OTHER DISEASES OF THE RESPIRATORY APPARATUS (Phthisis excepted). *Include:* Tracheostenosis; pleurodynia; pneumopathy; hydatids of the lung; pulmonary calculus; abscess of the lung; pulmonary anthracosis; interstitial pneumonia; cirrhosis of the lung; secondary sclerosis; hay fever (summer bronchitis or catarrh). To be also included when their nature is not indicated: Organic lesion of the lung; pulmonary accidents; hæmoptysis; spitting of blood; pulmonary hemorrhage; pneumorrhagia; bronchorrhagia; tracheotomy.—*Do not include:* Cancer of the lung (45).

#### V. DISEASES OF THE DIGESTIVE APPARATUS.

100. AFFECTIONS OF THE MOUTH AND ITS ADNEXA. *Include:* Diseases of the gums; epulis; gingivitis; ulorrhagia; glossitis; diseases of the tongue (except cancer); parotid tumor; parotiditis; salivary fistula; ranula; thrush; diseases of the teeth; odontalgia; dental caries; staphylitis; staphyloplasty; staphylorrhaphy.—*Do not include:* Cancer of the lips or tongue (39); chancre of the mouth (36a); noma (142); mumps (19); gangrene of the mouth (142); diseases of the palate (146 or 36); fracture of the maxilla (164); necrosis of the maxilla (146); paralysis of the velum palati (101).
101. AFFECTIONS OF THE PHARYNX. *Include:* Angina or Ludwig's disease; anginas of all descriptions (except diphtheritic angina and its symptoms; see Diphtheria, No. 9); amygdalitis; quinsy; abscess of the fauces, throat, or

<sup>1</sup> Return to the physician the reports given in as "bronchitis," in order that he may specify acute or chronic. When the physician fails thus to answer, classify under No. 90 all reports relating to children under five years of age, and under No. 91 all reports as to those of greater age.

<sup>2</sup> In countries where "apical pneumonia" is always synonymous with "phthisis," class this diagnosis under No. 27. In countries, on the contrary, where this is not constant, class under No. 93.

- retropharynx; paralysis of the velum palati; elongation of the uvula; pharyngitis; tonsillitis.—*Do not include:* Angina pectoris (80); cardiac angina (80); scarlatinal angina (7).
102. AFFECTIONS OF THE ŒSOPHAGUS. *Include:* Foreign bodies in the œsophagus; wound of the œsophagus; stricture of the œsophagus (except from cancer); spasm of the œsophagus; œsophagotomy.—*Do not include:* Cancer of the œsophagus (40); syphilitic stricture of the œsophagus (36).
103. ULCER OF THE STOMACH. *Include:* Round ulcer.—*Frequent complications:* Hæmatemesis; perforations of the stomach; peritonitis.
104. OTHER AFFECTIONS OF THE STOMACH (Cancer excepted).<sup>1</sup> *Include:* Dilatation of the stomach; paresis of the stomach; dyspepsia; apepsia; gastritis; gastrohepatitis; foreign body in the stomach; gastrotomy; perforation of the stomach (non-traumatic); gastralgia; "vertigo a stomacho læso;" catarrh of the stomach; indigestion. To be also included when their nature is not indicated: Gastrorrhagia; hæmatemesis; gastric hemorrhage.—*Do not include:* Gastro-enteritis (105 or 106, according to age).
105. DIARRHŒA AND ENTERITIS (under two years). *Include:* Gastro-enteritis or gastrocolitis of children; infantile enteritis; cholera infantum; athrepsia. This title only considers these ailments in children under two years.
- 105a. DIARRHŒA AND ENTERITIS, CHRONIC (under two years). *Include:* Athrepsia.
106. DIARRHŒA AND ENTERITIS (two years and over). *Include:* Gastro-enteritis or gastrocolitis of adults; enteritis of adults; diarrhœa of adults; lien-enteritis; intestinal ulcerations; colitis; intestinal colic; flatulent colic; inflammatory colic. *Do not include:* Tuberculous enteritis (29).
107. INTESTINAL PARASITES. *Include:* Helminthæ; oxyuri; tænia, of all kinds and descriptions; solitary worm; ascaris lumbricoides; trematodes; trichocephalus; ankylostomes; colic from worms.
108. HERNIAS AND INTESTINAL OBSTRUCTIONS. *Include:* Internal strangulation; intestinal invagination; stercoral tumors; ileus; intestinal occlusion; volvulus; hernial colic; hernial gangrene. The following to be included when their nature is not specified: Merocele; sarco-epiplocele; sarco-epi-omphalitis; kelotomy; herniotomy; artificial anus; stercoraceous vomiting.—*Do not include:* Laparotomy (without other qualification) (46 and 179).—*Frequent complication:* Peritonitis.
109. OTHER AFFECTIONS OF THE INTESTINES. *Include:* Paralysis or paresis of the intestine; enteroptosis; constipation; stercoræmia; intestinal calculi; intestinal perforation; foreign bodies in the intestine or rectum; rectitis. Include also the following diseases when their nature is not indicated, and these operations when their cause is not specified: Enterotomy; artificial anus; enterorrhagia; intestinal hemorrhage; melæna; prolapsus of the rectum; stricture of the rectum.—*Do not include:* Stercoral tumor (108); intestinal invagination and its synonyms (108); typhlitis and appendicitis (118); perityphlitis (118).
- 109a. DISEASES OF THE ANUS AND FECAL FISTULAS. *Include:* Proctitis; periproctitis; proctocèle; proctoptosis; fissure of the anus; abscess of the margin of the anus; fistula of the anus, either fecal or rectovaginal.—*Do not include:* Urinary fistulæ, even when these involve the rectum (124); artificial anus (108) (morbidity statistics alone); unnatural anus (108); imperforate anus (150).<sup>2</sup> (statistics alone.)

<sup>1</sup> See obs.<sup>2</sup> of the stomach.

110. ACUTE YELLOW ATROPHY OF THE LIVER. *Include:* Pernicious icterus; parenchymatous hepatitis; Weil's disease.—*Do not include:* Icterus (without qualification) (114); chronic icterus (114); icterus of the new-born (under three months) (151).
111. HYDATID TUMORS OF THE LIVER. *Include:* Hydatid cyst: hydatids (without qualification); echinococci.
112. CIRRHOSIS OF THE LIVER. *Include:* Cirrhosis (without qualification); alcoholic cirrhosis; interstitial cirrhosis; biliary cirrhosis; amyloid or fatty degeneration of the liver; slow atrophy of the liver; steatosis of the liver; alcoholic, interstitial, or chronic hepatitis.—*Do not include:* Organic lesion of the liver (114); hypertrophy of the liver (114).—*Frequent complications:* Dropsy; hemorrhage; pneumonia; tuberculosis.
113. BILIARY CALCULI. *Include:* Hepatic calculi; biliary lithiasis; hepatic colic.
114. OTHER AFFECTIONS OF THE LIVER. *Include:* Abscess of the liver; hepatitis: hepatitis, acute; angiocholitis; cholecystitis; hepatocystitis; choluria. To be also included when their precise nature is not indicated: Organic lesion of the liver; tumor of the liver; hypertrophy of the liver; acholia; cholangemia; icterus (over three months); chronic icterus; jaundice; hepatic congestion.—*Do not include:* Acute yellow atrophy of the liver (110); icterus of the new-born (151).
115. AFFECTIONS OF THE SPLEEN. *Include:* Splenitis; splenopathia; megalosplenism; splenocele.—*Do not include:* The affections of the spleen due to leukæmia (53) or malaria (4).
116. PERITONITIS, SIMPLE (Puerperal excepted).<sup>1</sup> *Include:* Peritonitis (without qualification); peritonitis, chronic; peritoneal adhesions; epiploitis; metropéritonitis, pelvipéritonitis.—*Do not include:* Tuberculous peritonitis (29); cancer of the peritoneum (41); puerperal peritonitis (137); rheumatic peritonitis (47).
117. OTHER AFFECTIONS OF THE DIGESTIVE APPARATUS (Cancer and Tuberculosis excepted). *Include:* Diseases of the pancreas (cancer excepted).
118. APPENDICITIS AND ABSCESS OF THE ILIAC FOSSA. *Include:* Iliac phlegmon or abscess; typhlitis; perityphlitis; typhlodidcliditis; appendicitis.—*Do not include:* Pelvic (130) or periuterine abscess; pelvic suppuration (130).

#### VI. DISEASES OF THE GENITO-URINARY APPARATUS AND ITS ADNEXA (NOT INCLUDING VENEREAL DISEASES).

119. NEPHRITIS, ACUTE. *Do not include:* Scarlatinous nephritis (7); chronic nephritis (120); tuberculous nephritis (33); nephritis of pregnancy (138).
120. BRIGHT'S DISEASE. *Include:* Chronic, albuminous, interstitial, or parenchymatous nephritis; albuminuria; amyloid or fatty degeneration of the kidney; amyloid kidney; steatosis of the kidney; renal sclerosis. To be included when their precise nature is not indicated: Uræmia; uræmic eclampsia; uræmic delirium; uræmic coma.—*Do not include:* Organic lesion of the kidney (121); puerperal uræmia (138); cardiac albuminuria (79).—*Frequent complications:* Anasarca; dropsy; convulsions; hemorrhages; cerebral apoplexy; pneumonia.

<sup>1</sup> When a female of childbearing age is returned as having been stricken with "peritonitis," without other explanation, the report should be returned in order that the physician may specify whether or not the condition was puerperal.



121. OTHER DISEASES OF THE KIDNEYS AND THEIR ADNEXA. *Include:* Pyelitis; anuria; renal congestion; renal ectopia; nephroptosis; floating, motile, or displaced kidney; movable kidney; renal cysts; polycystic kidney; hydro-nephrosis; hæmaturia; perinephritis; perinephritic and nephritic abscess; pyelonephritis; nephropyosis. To be also included when their nature is not specified: Organic lesion of the kidney; nephrorrhagia.
122. CALCULI OF THE URINARY TRACT. *Include:* Renal, ureteral, nephritic, vesical, or urinary calculus; nephritic colic; nephrolithiasis; gravel; stone; calculous affections; urinary lithiasis; lithotrixy; lithoclasty.—*Do not include:* Prostatic calculus (125).
123. DISEASES OF THE BLADDER. *Include:* Cystitis, acute or chronic; vesical or ureteral catarrh; cystorrhagia; tumor of the bladder; cystocele; cystop-tosis; foreign body in the bladder; cystotomy; retention of urine; dysuria; paralysis and section of bladder; vesical inertia; incontinence of urine; tenesmus of the bladder.—*Do not include:* Hæmaturia (121); uri-nary fistula, even when they involve the bladder (124); cystosarcoma (45).
124. DISEASES OF THE URETHRA. *Include:* Urinary abscess, etc.; ankylurethria; foreign bodies; urethrotomy; urinary fistula (urethral, urethrorectal, vesi-corectal, or vesicometrorectal); urinary infiltration; urinary intoxication; urethralgia; urethrorrhagia; urinæmia; stricture of the urethra; urethro-stenosis; urethroplasty; urethrorrhaphy; stricture (male).—*Do not in-clude:* Ureteral catarrh (123); retention of urine (123).
125. DISEASES OF THE PROSTATE. *Include:* Hypertrophy of the prostate; prosta-titis; abscess of the prostate; prostatic calculus.—*Do not include:* Cancer of the prostate (45); tuberculosis of the prostate (33).
126. NON-VENEREAL DISEASES OF THE GENITAL ORGANS OF THE MALE. *Include:* Phimosis; paraphimosis; amputation of the penis; seminal losses; sper-matorrhœa; orchitis; epididymitis; funiculitis; hydrocele; hæmatocele of the testicle, cord, or scrotum; castration (in man); Malassez's disease.—*Do not include:* Cancer of the testicle (45); tuberculosis of the testicle (33); sarcydrocele (45); syphilitic sarcocele (36); varicocele (83).
127. METRITIS (non-puerperal or unqualified). *Include:* Ulcer of the uterus; ulceration of the neck (of the womb); endometritis.
128. UTERINE HEMORRHAGE, NON-PUERPERAL. *Include:* Metrorrhagia; menorrha-gia; tamponage of the vagina or uterus.
129. UTERINE TUMOR (not cancerous). *Include:* Fibroid tumor, or fibroid of body of the uterus; hysteromyoma; uterine polypus; fungous or fungoid tumors of the uterus; Huguier's disease.
130. OTHER DISEASES OF THE UTERUS. *Include:* Procidentia of uterus; uterine or vaginal catarrh; deviation, ante flexion, retro flexion, anteversion, falling or prolapse of the uterus; prolapse of the vagina; uterine prolongation; amenorrhœa; hypertrophy of the neck of the uterus; dysmenorrhœa; organic lesion of the uterus; hysterectomy; hysterotomy; metrotomy; ablation of the uterus; abscess of the pelvis; periuterine or retro-uterine abscess or phlegmon; pelvic suppuration; leucorrhœa; fluor albus (whites; vaginal flow; white flux).—*Do not include:* Puerperal diseases (134 and 141); abscess of the iliac fossa (118).
131. CYSTS AND OTHER TUMORS OF THE OVARY. *Include:* Ovariectomy; castration (in the female). Dermoid cyst often classified better here than under 146.

132. OTHER DISEASES OF THE GENITAL ORGANS OF THE FEMALE. *Include:* Vaginitis; tumors of the vagina; ovaritis; salpingitis; salpinx; metrosalpingitis; hæmatosalpinx; pyosalpinx; abscess and tumors of the vulvovaginal glands; vulvitis; periuterine or retro-uterine hæmatocele.—*Do not include:* Urinary fistulæ (124); stercoral fistulæ (109a); even when they involve the genital organs. King makes a subheading of "Diseases of tubes."
133. NON-PUERPERAL DISEASES OF THE BREAST (Cancer excepted). *Include:* Mammitis; abscess of the breast (non-puerperal); cyst of the breast; cystic disease of Reclus; tumor of the breast (without qualification, or non-cancerous); amputation of the breast.—*Do not include:* Fistula of the breast (puerperal, or without qualification) (43); mammary cancer (43).

#### VII. THE PUERPERAL STATE.

REMARKS.—It often happens that physicians neglect to note the puerperal character of the disease; hence the following rule for the guidance of those whose duty it is to collect statistics. "Whenever a female of childbearing age is noted as dead from a disease which *may* be puerperal, the report should be returned to the reporter, in order that he may state explicitly whether or not the disease was puerperal." The following are these diseases: Peritonitis; pelviperitonitis; metroperitonitis; septicæmia; hemorrhage; metrorrhagia; eclampsia; phlegmasia alba dolens; phlebitis; lymphangitis; embolism; sudden death; abscess of the breast.

134. ACCIDENTS OF PREGNANCY. *Include:* Miscarriage (death of mother); abortion (death of mother); hemorrhage of pregnancy; incoercible vomiting; rupture of tubal pregnancy; ablation of the pregnant tube; difficulties and fatigues supervening in the course of pregnancy.
- 134a. LABOR, NORMAL. (Morbidity statistics only.)
135. PUERPERAL HEMORRHAGE. *Include:* Puerperal metrorrhagia; post-partum hemorrhage.
136. OTHER ACCIDENTS OF LABOR. *Include:* Dystocia; Cæsarean section; rupture of the uterus; metrorrhæxia; laceration or rupture of the perineum; perineorrhaphy; placenta prævia; malposition, retention, detachment, or apoplexy of the placenta; cephalotripsy; embryotomy (adult); symphyseotomy; version; application of forceps; uterine inversion.
137. PUERPERAL SEPTICÆMIA. *Include:* Puerperal fever; puerperal infection; puerperal endometritis; puerperal salpingitis; perimetrosalpingitis, or phlegmon of the broad ligament, or diffuse pelvic puerperal cellulitis; puerperal peritonitis, metroperitonitis, phlebitis, lymphangitis, or pyohæmia.—*Do not include:* Septicæmia (without qualification) (20).
138. PUERPERAL ALBUMINURIA AND ECLAMPSIA. *Include:* Puerperal uræmia; nephritis of pregnancy; eclampsia of women in labor; epileptiform convulsions of women in labor; puerperal tetanus.
139. PHLEGMASIA ALBA DOLENS, PUERPERAL. *Do not include:* Phlegmasia alba dolens, non-puerperal (82).—*Frequent complications:* Gangrene; embolism.
140. OTHER PUERPERAL ACCIDENTS; SUDDEN DEATH. *Include:* Puerperal embolism; puerperal thrombus; sudden death in the puerperium; consequence of labor (without other explanation); subinvolution of uterus.—*Do not include:* Sudden death, non-puerperal (178); puerperal scarlatina (7).
141. PUERPERAL DISEASES OF THE BREAST. *Include:* Fissure of the nipple (puerperal); circumscribed abscess; abscess of the breast (puerperal); fistula of the breast (puerperal or without further indication).

## VIII. DISEASES OF THE SKIN AND CELLULAR TISSUE.

142. GANGRENE. *Include*: Eschar; sphacelus; gangrene, dry; gangrene, senile; gangrene of the extremities; gangrene of the mouth; gangrene of the vulva, etc.; noma; Raynaud's disease.—*Do not include*: Gangrene of the lung (96); hernial gangrene (108); gangrenous erysipelas (18 or 144).
143. FURUNCLE (Carbuncle).<sup>1</sup> *Not included*: Biskra, Aleppo, or Medina button (145).
144. PHLEGMON; WARM ABSCESS. *Include*: Abscess (without qualification); phlegmonous tumor; adenophlegmon; suppurative adenitis; bubo (without qualification); suppurating bubo; diffuse phlegmon; phlegmonous or gangrenous erysipelas; panaris; whitlow; abscess of the mediastinum; vomica (without any other indication).—*Do not include*: Bacillary abscess (33); abscess of the fauces, throat, or retropharynx (101); of the liver (114); of the iliac fossa (118); of the female pelvis (130); of the prostate (125); urinary (124); periuterine (130); of breast, non-puerperal (130); cold (31); by congestion (31); ossifluent (31); angioleucitis (84).
145. TINEA FAVUS. (Morbidity statistics alone.)
- 145a. TINEA TONSURANS, TRICHOPHYTON. *Include*: Tinea (without qualification). (Morbidity statistics alone.)
- 145b. PELADES. (Morbidity statistics alone.)
- 145c. ITCH. (Morbidity statistics alone.)
- 145d. OTHER DISEASES OF THE SKIN AND ITS ADNEXA. *Include*: Erythema; urticaria; prurigo; pityriasis; lichen; psoriasis; dermatitis; eczema; impetigo; aphtha; herpes; ecthyma; elephantiasis Arabum; pachydermatitis; polysarcia; scleroderma; cheloids; fungoid mycosis; seborrhœa; trophoneuroses; zona; Wardrop's disease; Biskra, Aleppo, or Medina button; Penjdeh ulcer; Cochín-China ulcer; pemphigus; myiasis.—*Do not include*: Pachydermatous cachexia (89); elephantiasis Græcorum (17).

## IX. DISEASES OF THE ORGANS OF LOCOMOTION.

146. AFFECTIONS OF THE BONES (Non-tuberculous). *Include*: Periostitis; periostosis; osteitis; osteoperiostitis; osteomyelitis; caries; necrosis; sequestrum; perforation of the palatine vault; necrosis of the maxilla (non-phosphoric or without qualification); exostosis (without qualification); osteoma; osseous tumor; cranial tumor; foreign bodies in the frontal or other sinuses; mastoiditis; abscess of the frontal or maxillary sinus; osteomalacia; softening of bone; rhachitis; scoliosis; lordosis; kyphosis.—*Do not include*: Caries of the petrous bone (76); dental caries (100); osteocopic pains (36); osteosarcoma (45); phosphorus necrosis (58).
147. ARTHRITIS AND OTHER DISEASES OF THE JOINTS (Tuberculosis and Rheumatism excepted). *Include*: Arthritis; polyarthritis (non-vertebral); hydrarthrosis; foreign bodies in joints; arthrodynia; arthropyosis; arthrophytis; ankylosis; arthralgia; arthrocele; genu valgum.—*Do not include*: Rheumatic arthritis (47).
148. AMPUTATION.<sup>2</sup> *Include*: Only those cases in which the lesion, the cause for amputation, is not specified.—*Do not include*: Amputation of the breast

<sup>1</sup> See No. 22.<sup>2</sup> Amputation, surgical operation, shock, and surgical shock, unqualified, are not sufficiently specific terms as causes of death.



(133); amputation of the penis (126).—*Frequent complications:* Septicæmia; erysipelas; tetanus; hemorrhage.

149. OTHER AFFECTIONS OF THE ORGANS OF LOCOMOTION. *Include:* Hygroma; perichondritis; disarticulation; tarsalgia; painful talipes valgus; retraction of the fingers or of the palmar aponeurosis; Dupuytren's disease; non-traumatic muscular rupture; muscular diastasis; myodiastasis; non-traumatic rupture of a tendon; diseases of tendons; tenophytes; tenosynovitis; tenotomy; tenorrhaphy; torticollis; lumbago; curvature.

#### X. MALFORMATIONS.

150. MALFORMATIONS (Stillbirths not included). *Include:* Malformation; monstrosity; anomaly; arrest of development; congenital hydrocephalus; hydrocephalus (without qualification); megalcephalus; hydrorachia; spina bifida; encephalocele; podencephalia; congenital eventration; omphalocele; exomphalos; ectopia; imperforate anus, etc.; hare-lip; cleft palate; anaspadias; hypospadias; cryptorchid; vascular nævus; polydactylia; syndactylia; congenital club-foot; talipes valgus, varus, or equinus, congenital; congenital deafness or blindness; persistence of the foramen of Botalli (foramen ovale).—*Do not include:* Coloboma (75); painful flat-foot (149); acquired hydrocephalus (74b); tuberculous hydrocephalus (28). King makes a separate subheading for hydrocephalus and cyanosis.

#### XI. EARLY INFANCY.

- 150a. THE NEW-BORN AND NURSINGS DEPARTING FROM HOSPITALS WITHOUT HAVING BEEN SICK. (Morbidity statistics alone.)
151. CONGENITAL DEBILITY, ICTERUS, AND SCLEREMA.<sup>1</sup> *Include:* Premature birth (not stillborn); atrophy (infantile); icterus or hepatitis of the new-born; atelectasis of the lungs in the new-born; œdema of the new-born.
152. OTHER DISEASES OF EARLY INFANCY. *Include:* Umbilical hemorrhage; inflammation of the umbilicus; cyanosis of the new-born. (This title has reference to children not more than three months old.)
153. LACK OF CARE.

#### XII. OLD AGE.

154. SENILE DEBILITY. *Include:* Senility; old age; cachexia (of the old); senile exhaustion; senile dementia.—*Do not include:* Senile gangrene (142).

#### XIII. AFFECTIONS PRODUCED BY EXTERNAL CAUSES.

Among suicides there should only be classed those in whom suicide or attempted suicide is clearly demonstrated. In collective suicides there should only be counted those who have attained their majority. Minors ought to be regarded as the victims of assassination and placed under 176.

155. SUICIDE BY POISON. *Include:* Voluntary poisoning; voluntary absorption of sulphuric acid (or any other corrosive substance).—*Do not include:* Cocainism (59); morphinism (59).
156. SUICIDE BY ASPHYXIA. *Include:* Suicide by the vapor of charcoal.
157. SUICIDE BY HANGING OR STRANGULATION. *Include:* Hanging.
158. SUICIDE BY DROWNING.

159. SUICIDE BY FIREARMS.
160. SUICIDE BY CUTTING INSTRUMENTS.
161. SUICIDE BY JUMPING FROM HIGH PLACES.
162. SUICIDE BY CRUSHING.
163. OTHER SUICIDES.
164. FRACTURES. *Include:* Separation of the epiphyses; fracture of the cranium.
165. SPRAINS. *Include:* Strains; ligament-stretching. (Morbidity statistics only.)
- 165a. LUXATIONS. *Include:* Subluxations; dislocations.
166. OTHER ACCIDENTAL TRAUMATISMS. *Include:* Stabs; contusion; bites (non-venomous, non-virulent); crushing; railroad accidents (suicide excepted); wounds by cutting instruments (suicide not demonstrated); accidental falls; concussion of the brain; perforation of the cranium; traumatic hemorrhage; traumatic fever; traumatic eventration; perforation of the abdomen or chest; all acute affections designated as "traumatic;" wounds by firearms. King subdivides into: accidental gunshot wounds; injuries by machinery; injuries in mines and quarries; railroad accidents and injuries; injuries by horses and vehicles; and other accidental traumatism.
167. BURNS AND SCALDS. *Include:* Burns and scalds from steam, petroleum, gasoline, boiling liquid, etc.—*Do not include:* Conflagration (174).
168. BURNS FROM CORROSIVE SUBSTANCES. *Include:* Burns by vitriol.
169. INSOLATION. *Include:* Sunstroke.
170. FREEZING. *Do not include:* Effects of cold (new-born) (153).
171. ELECTRICAL SHOCK. *Include:* Death from lightning.
172. ACCIDENTAL SUBMERSION. *Include:* Drowning (non-suicidal).
173. PROSTRATION. *Include:* Fatigue. (Morbidity statistics alone.)
- 173a. INANITION. *Include:* Hunger; insufficient food (new-born excepted); misery.—*Do not include:* Lack of care (new-born) (153); lack of nutrition (new-born) (153); sitiophobia (68); hysterical anorexia (74a).
174. ABSORPTION OF DELETERIOUS GASES (Suicide excepted). *Include:* Asphyxia, accidental (pathological asphyxia and suicidal asphyxia excepted); asphyxia by illuminating gas; asphyxia by stoves (fixed or portable); absorption of carbonic oxid; conflagration; absorption of ammonium sulphid; asphyxia by night-soil; absorption of chloroform; absorption of nitrous oxid.—*Do not include:* Asphyxia of the adult (without qualification) (179); asphyxia (under three months) (152).
175. OTHER ACUTE POISONINGS. *Include:* Every acute poisoning (suicide excepted); antimony cholera; acute ergotism; absorption of venom; bite of serpent; accidental absorption of sulphuric acid or other corrosive substances.—*Do not include:* Saturnism (57); hydrargyrism, etc. (58 or 59); morphinism, chronic ergotism, etc. (59).
176. OTHER EXTERNAL VIOLENCE. *Include:* Accident (without other qualification); bad treatment (upon a child); capital punishment; foreign body in the larynx; foreign body in the trachea. King subdivides into: suffocation; injuries at birth; homicide; and other external violence.

#### XIV. ILL-DEFINED DISEASES.

The following titles will include only those conditions ill-defined by the physician, whether from lack of sufficient data, or because the disease was ill-defined, or because the physician was negligent in making a complete diagnosis.

177. **DROPSY.** *Include:* Anasarca; ascites; œdema of the extremities or generalized œdema; organic lesion (not defined).—*Do not include:* Œdema of the new-born (151); œdema of the glottis (88); œdema of the lungs (95); œdema of the brain (64).
178. **SUDDEN DEATH.** *Include:* Syncope (followed by death).—*Do not include:* Puerperal sudden death (140), nor sudden death followed by an explanation, as "diabetic" (50) or "apoplectic" (64).
179. **ILL-DEFINED OR UNSPECIFIED CAUSES OF DEATH.** *Include:* Exhaustion or cachexia or debility (of adults); asthenia; adynamia; ataxo-adynamia; coma; asthenic, hectic, colliquative, synochal, gastric, bilious, or pituitary fever; gastric involvement; fever of dentition; paralysis of the heart (in German "herzlahmung" or "herzschlag," in English "heart failure"); cyanotic asphyxia (without indicated cause, the new-born excepted); or any other insufficient diagnosis.—*Do not include:* Exhaustion, cachexia or debility of the old (154); fever, ataxo-adynamic (1), continued (1), summer, or hay (99); asphyxia by external cause (156 or 174); cyanosis of the new-born (152).
180. **STILLBIRTHS.** Stillbirths are not included among deaths, as a stillborn child is one born dead. Still, it is wise to have a separate heading for them, including under congenital debility (151) and unknown (179) cases where the child lived after birth and a definite cause of death cannot be assigned.

An endeavor is now being made to adopt the following death certificate throughout America.

## RETURN OF A DEATH

IN THE CITY, TOWN, OR TOWNSHIP OF.....

### Physician's Certificate.

1. Full Name of Deceased,.....
2. Color,..... State if { Chinese,  
Japanese,  
Indian.
3. Sex,.....
4. Single, Married, ..... State if { Widow,  
Widower,  
Divorced.
5. Age, { Years, ..... 6. Date of Death. { Year, .....  
Months, ..... Month, .....  
Days, ..... Day, .....  
(If age is less than one day, give hours.....)
7. Cause of Death, { Chief, .....  
Contributing, .....

No Certificate will be accepted which is **MUTILATED, ILLEGIBLE, INACCURATE, or any portion of which has been ERASED, INTERLINED, CORRECTED, or ALTERED,** as all such changes impair its value as a Public Record.

*This Certificate must not be issued for any other purpose than as a report to the Board of Health. Should the Physician issue a duplicate, it must be distinctly marked "Duplicate," and state why issued.*

..... M.D.  
Residence, .....

Write plainly, and with ink; fill in every blank space.



## CHAPTER XXIX

### LITERATURE

#### Post-mortem References

- ADLER, N. Whether post-mortem examinations under any circumstances should be allowed in the Jewish Hospitals. *The Occident*, June, 1856, v. XIV, n. 3.—ADMINISTRATION des hôpitaux et les autopsies. *Arch. gén. de méd.*, 1842, v. I, p. 114.—ALBERS, J. F. H. Bericht über 84 Leichenöffnungen. *Deutsche Klinik*, 1850, v. II, p. 413.—ALLEN, H. Synopsis of autopsies made at Lincoln General Hospital. *Proc. Path. Soc. of Phila.*, 1864, v. II, p. 160.—ALLISON, H. E. On a general system of reporting autopsies in American asylums for the insane. *Am. J. Insan.*, 1889-90, v. XLVI, p. 216.—AMANO, F. Zusammenstellung der Resultate von der letzten 100 Sektionsfällen nebst einigen interessanter Befunden. (Japanese text.) *Okayama Igaku Kwai Zasshi*, 1898, p. 67; p. 102.—AMES, D. *Brief notes on the conduct of post-mortem examinations*. Baltimore, 1897.—AMMANN, P. *Oratio de autopsia medica*. Lipsiæ, n. d.—AMON. Zwei Sectionen aus der Privatpraxis. *München. med. Wchnschr.*, 1895, v. XLII, p. 1183.—AUTOPSIES cadavériques des rois de France. *Clinique*, 1829, v. I, p. 136.—AUTOPSIES at the Leichenhof, Vienna. *Med. Times and Gaz.*, 1872, v. II, p. 605.—AZEVEDO, N. *Técnica d'autopsias clinicas*. *Med. contemp.*, Lisbon, 1903, v. XXII, p. 177.
- BAADER, J. Observationes medicæ incisionibus cadaverum anatomicis. *SANDEFORT'S Thesaurus dissertationum*. Lugd. Batv., 1778, v. III.—BADES, V. Post-mortem examinations with a view to bacteriological research. *Arch. roum. de méd. et chir.*, Par., 1887, v. I, p. 157.—BAETENS, H. J. *Dissertatio medico-legalis de occisi hominis cadaveris inspectione, a inde renunciandâ lethalitate*. Par., 1808.—BANDOUIN, M. Les autopsies dans les hôpitaux. *Gaz. méd. de Par.*, 1901, v. XII, p. 1.—BAREZ. Leichenöffnungen. *Wchnschr. f. d. ges. Heilk.*, 1897, v. II, p. 102.—BEALE, L. S. *Suggestions for taking cases and for making post-mortem examinations*. Lond., 1867.—BEATSON, W. B. Précis of medicolegal post-mortem examinations in Dacca during the year 1865. *Indian M. Gaz.*, 1866, v. I, p. 82.—BENEKE. Zur Technik der Oberkiefer und Nasenhöhlensection. *Centr. f. allg. Path. und path. Anat.*, Oct. 25, 1896; Zur Technik der Bauchsection, *ibidem*, 1900, v. II, p. 433.—BERICHT der vom V. deutschen Aerztetag gewählten Commission für Herstellung der Vorarbeiten zur Einführung der allgemeinen Leichenschau in Deutschland. *Aerzt. Vereinsbl. f. Deutschl.*, 1878, v. VII, p. 3.—BERLINER, P. Plastische Reproduction path.-anat. Präparate. *D. med. Wchn.*, 1892.—BERTRAND, L. C. *Quelques faits d'anatomie pathologique*. Par., 1837.—BIERMAYER. Bericht über die Leichenöffnungen. *Med. Jahrb. d. k. k. österr. Staates*, 1816, v. III, p. 73.—BILL to provide for inquests under national authority. Washington, 1886.—BLACKBURN, I. W. *A manual of autopsies designed for the use of hospitals for the insane and other public institutions*. Phila., 1892; *Intercranial Tumors among the Insane. A report of 1642 autopsies in cases of mental disease*. Washington, 1903.—BLEICHRODER. Die Funktionsprüfung der Mitralklappe bei der Herzsection. *Virchow's Archiv*, 1902, v. CLXIX.—BLUMER, G. Examining and recording pathological material. *Albany M. Ann.*, 1902, v. XXIII.—BOILEAU, J. P. H.—Gravimetry in the mortuary, etc. *British Med. Journ.*, 1881.—BOLLES, W. P. Cases with autopsies. *Med. and Surg. Rep. Bost. City Hosp.*, 1877, p. 270.—ROLLINGER, O. *Atlas und Grundriss der pathologischen Anatomie*. München, 1901; *Beiträge zur pathologischen Anatomie*. Wiesbaden, 1903; *Arbeiten aus dem pathologischen Institut zu München*. München, 1886; Bericht über die Leichenöffnungen im pathologischen Institut während der Jahre 1897-1898-1899. *Ann. d. städt. allg. Krankenh. zu München*, 1901, v. XI, p. 352.—BONET, T. *Sepulchretum sive anatomia practica ex cadaveribus morboedatis*. Genevæ, 1700.—BOOBY, G. Post-mortem observations on the appendix. *Bull. Iowa Inst.*, 1903, v. V, p. 314.—BOURNEVILLE and P. BRICON. *Manuel de technique des autopsies*. Par., 1885.—BOWDITCH, H. P. On the collection of data at autopsies. *Boston M. and S. J.*, 1882, v. BOYD. Post-mortem appearances in 75 cases. *Edinb. M. and S. J.*, 1904, p. 304.—BRESLAU. Über eine neue Methode der Eröffnung

den Leichen Neugeborner. *Wien. med. Wchnschr.*, 1862, v. XII, p. 145.—BROFFEAU. Ouverture cadavérique. *Dict. d. sc. med.* Par., 1819, v. XXXVIII, p. 55.—BRODIER, L. Étude statistique de 716 autopsies. *Arch. gén. de méd.*, 1901, n. s. v. VI, p. 76.—BRUNETTI, L. Lettera sopra il suo nuoro rachiotomo e sul metodo di aprire lo speco vertebrale adottato dalla scuola di anatomia patologica in Padova. *Gazz. med. ital., prov. venete.* Padova, 1863, v. VI, p. 133. See also *Scuola anat. patol. d. r. Univ. di Padova*, 1878. *La tannizzazione dei tessuti animali dai patologhi.* Padova, 1888.—BÜCHNER, L. Sectionsbericht über 145 Leichenöffnungen. *Arch. f. path. Anat.*, 1859, v. XVII, p. 343.—BUHLIG, W. H. General and special methods in the post-mortem examination of the brain and spinal cord. *Cleveland M. J.*, 1904, v. III, p. 28.—BULLARD, W. N. The importance of well made and accurately reported autopsies in the determination of the etiology of weak-mindedness and idiocy. *J. Psycho-Asthenics.* 1903-1904, v. VIII, p. II.—BURNS, A. E. Autopsies. *Med. Sentinel*, Portland, Oregon, 1896, v. IV, p. 24.—BURT, S. S. Diagnosis in the light of a necropsy. *Postgraduate*, N. Y., 1900, v. XV, p. 1092.—BUSH, M. J. and BRANDTS, J. G. Rapport sur le résultat d'autopsies cadaveriques. *Biblioth. méd.*, Brux., 1827, v. IV, p. 64.—BUSSE, O. *Das sektionen-Protokoll.* Berlin, 1903.

CADAVER in volume II of the *Index Catalogue*, 1898, will be found to contain some additional post-mortem references. CARPENTER, I. L. On Autopsies. *Tr. N. Hampshire M. Soc.*, Concord, 1897, p. 220.—CARRINGTON, R. E. *Notes on Pathology. A handbook for the Post-mortem room.* London, 1892.—CASSEBOHN, J. F. *Methodus secandi et contemplandi viscera hominis.* Halæ, 1740.—CATTELL, H. W. Some special points on the performance of autopsies on the new-born. *Am. Gynec. and Pædiat.*, 1892, v. VI, p. 758; *Some practical post-mortem points* Phila., 1893; *Notes on the demonstrations in morbid anatomy, including autopsies* Phila., 1893; *Post-mortem pathology.* Phila., 1903.—CATTELL, T. The precautionary measures which may be employed in dissections and post-mortem examinations. *Lancet*, Lond., 1847, v. I, p. 645.—CAUDEREAU. Sur un procédé nouveau de dissociation des glandes. *Compt. rend. Soc. de biol.*, 1879, v. I, p. 291.—CAUTIONS for the pathological anatomist. *Med. Mag.*, Bost., 1834, v. VII, p. 491.—C. E. De l'importance des autopsies pour les familles. *Lyon méd.*, 1878, v. XXVII, p. 169.—CÉRÉBROTOME du Dr. Gavoy. *Compt. rend. Soc. de biol.*, 1885, v. II, p. 114.—CERNÉ. Une salle d'autopsie. *Normandie méd.*, 1891, v. VI, p. 369.—CHAILLÉ, S. E. Post-mortem changes versus ante-mortem lesions. *N. O. M. and S. J.*, 1873-1874, n. s., v. L, p. 639.—CHIARI, H. *Pathologisch. anatomische Sektionstechnik.* Berl., 1894.—CLARKE, J. J. *Post-mortem examinations in medico-legal and ordinary cases.* Lond., 1893.—CLEMENT, A. W. *Veterinary post-mortem examinations.* New York, 1891.—COCKS, W. P. *Forms for facilitating the records of post-mortem appearances.* Lond., 1832; *Pathological anatomy of the liver, spinal cord, and the membranes.* (With cases.—Morbid appearance.) Lond., 1831.—CONNELL, W. T. Some post-mortem notes. *Kingston M. Quart.*, 1898-1899, v. III, p. 172.—COPLIN, W. M. L. *Manual of pathology, including bacteriology, the technique of postmortems and methods of pathological research.* Phila., 1900.—COUNCHMAN, W. T., and MALLORY, F. B. A study of the lesions in selected autopsies. *Med. and Surg. Rep. Bost. City Hosp.*, 1896, p. 216.—COX, G. W. Some practical post-mortem points. *Indian M. Rec.*, 1898, v. XV, p. 47.—CRAWFORD. *Post-mortem examination on Mr. Crawford.* (N. n., n. d.)—CRUSE, P. Obduktionsbefunde nach den Protokollen des Dorpater pathologischen Instituts. *Dorpat. med. Ztschr.*, 1871, v. I, p. 255.

DALAND, J. A new post-mortem case. *Tr. Path. Soc. Phila.*, 1898, v. XVIII, p. 482.—DALTON, J. C. A ready method of making brain sections for post-mortem examinations. *Boston M. and S. J.*, 1880, v. CIV, p. 57.—DALTON, N. Report of the pathological department: Analysis of post-mortem examinations performed from October 1, 1896, to September 30, 1897. *King's Coll. Hosp. Rep.*, 1896-1897, Lond., 1898, v. IV, p. 273; Report of the pathological department: Analysis of post-mortem examinations performed from October 1, 1895, to September 30, 1896. *Ibid.*, 1895-1896, Lond., 1897, v. III, p. 251.—DANIELS, C. W. Notes on a series of postmortems made in the Public Hospital, Georgetown, from April 1 to December 31, 1893. *Brit. Guiana M. Ann.*, Demerara, 1894, v. VI, p. 79; Further notes on a series of postmortems in the Public Hospital, Georgetown, April, 1893, to March, 1895. *Ibid.*, 1895, p. 50.—DAVY, J. On a new method of preserving anatomical preparations for a limited time. In his: *Researches and Anat.*, Lond., 1839; Some directions for making and keeping an-



preparations in hot climates. *Ibid*—DE ANGELIS, M. G. *Tecnica d autopsie*. Milan, 1893.—DÉGRANGES, E. *Souvenirs de la cour d' assises de la Gironde*. *Union méd. de la Gironde*, 1863, v. VIII, p. 401.—DELAFIELD, F. *A hand-book of post-mortem examinations and morbid anatomy*. New York, 1872.—DEMMÉ, T. A. A description of a craniotome for post-mortem examinations. *Med. and Surg. Reporter*, 1860, v. IV, p. 295.—DE RYCKERE and CORIN. A quel moment est-il permis au médecin légiste de pratiquer l'autopsie? *Ann. Soc. de méd. lég. de Belg.*, Charleroy, 1900-1901, v. XII, p. 126.—DEZEIMEZ's Autopsy. *Dis. répert. de médecine*. Par., 1834, v. VI, p. 160.—DOUGLASS, S. H. Method of conducting post-mortem examinations in cases of suspected poisoning. *Perins J. M.*, Ann Arb., Mich., 1853-1854, v. I, p. 49.—DOWLER, B. Post-mortem researches. *West. J. M. and S.*, Louisville, 1843, v. VII, p. 241; v. VIII, p. 241.—DUBINI, A. Una parola sui falsi risultati che si ottengono dalle sezioni cadaveriche incomplete. *Gazz. med.*, Milano, 1845, v. IV, p. 197; Dell' arte di fare le sezioni cadaveriche. *Ibid.*, 1847, v. VI, p. 319.—DUCHAUSSOY. Surprises que certaines autopsies nous réservent. *Bull. Soc. de méd. prat. de Par.*, 1891.—DWIGHT. The inquest. *Med. Rec.*, 1879.

ECKARD, E. How to conduct an autopsy. *Peoria M. J.*, 1896, v. I, p. 344.—EISENLOHR, L. Bericht über 3749 während der Jahre 1878-84 im Pathologischen Institut zu München vorgenommene Sectionen. *Arb. u. d. path. Inst. zu München*. Stuttgart, 1886, p. 462.—ENGEL. Methode bei Leichenuntersuchungen. *Wien. med. Wchnschr.*, 1870, v. XX, p. 1003; Aus der Leichenkammer der k. k. Josef's Akademie. *Spitals-Ztg.*, Wien, 1864, p. 209; *Sections-Beschreibungen*. Wien, 1861; Sectionsergebnisse an der Prager pathol.-anatomischen Anstalt vom Oct. 1, 1849-Feb. 1, 1850. *Vrtljschr. f. d. prakt. Heilk.*, Prag, 1850, v. XXVII, p. 9; *Darstellung der Leichenerscheinungen und deren Bedeutung*. Wien, 1854; Die Bestimmung der Blutmenge bei Leichenuntersuchungen. *Wchnbl. d. Ztschr. d. k. k. Gesellsch. d. Aerzte*. Wien, 1866, v. XXII, p. 341.—ERICHSEN, J. Bericht über die im Hebammeninstitut, Ihrer Kaiserlichen Hoheit der Grossfürsten Helena Pawlowna ausgeführten Sectionen für die Jahre 1862 und 1863. *St. Petersburg. med. Ztschr.*, 1865, v. VIII, p. 257.—ESCOLAR, S. Extracto del resultado de las necropsias practicadas en el mes de agosto en la sala de Santo Domingo del hospital general de este corte. *Bol. de med., ciruj. y farm.*, Madrid, 1850-1853, v. V.

FABRICIUS, P. C. *Methodus cadavera humana rite secandi*. Halae and Helmstadii, 1774; *De autopsiae in medicina utilitate et praestantia*. Helmstadii, 1784.—FEARN, J. R. Post-mortem Revelations. *Calif. M. J.*, 1896, v. XVII, p. 243.—FELDMANN, J. The relations of practising physicians to pathological post-mortems and the question of prosecution in our country. *Gyógyászat*, Budapest, 1902, v. XLII, p. 10.—FENGER, C. Beretning om 422 Sektioner. *Nord. med. Ark.*, Stockholm, 1873, v. V, No. 18.—FÉRÉ. Procédé de coup du crâne. *Progrès méd.*, 1877, v. V, p. 495.—FINLAYSON, J. Method of performing post-mortem examinations. *Clin. Diagnosis*, Phila., 1878.—FINNELL. Post-mortem cases. *Med. Rec.*, 1869, v. IV, p. 260.—FLEISCHMANN, D. G. Leichenöffnungen. *Repert. f. d. ges. Med.*, Jena, 1842, v. V, p. 273.—FLEMING, R. A. Experience gained from post-mortems in relation to the practice of medicine and surgery. *Internat. Clin.*, 1902, 12th S., v. II, p. 142.—FOCHIER, A. Le droit aux autopsies dans les hôpitaux. *Lyon méd.*, 1878, v. XXVII, p. 287.—FODÉRÉ, F. E. Règles de l'autopsie cadavérique des blessés. *Dict. a. sc. méd.*, Par., 1820, v. XLIII, p. 119.—FORM of record of autopsy. U. S. War Department, Surgeon-General's Office, Washington, 1882.—FORSTER, A. Uebersicht von 639 in den Jahren 1849-56 verrichteten Sectionen. *Schmidt's Jahrb.*, 1858, v. XCVII, p. 89; Fortsetzung der Instruction für die öffentlichen angestellten Aerzte und Wundärzte in den k. k. Oesterreichischen Staaten wie sie sich bei gerichtlichen Leichenbeschauen zu benehmen haben. *Med. Jahrb. d. k. k. österr. Staates*, Wein, 1816, v. III, p. 17.—FRIEDREICH, N. Über den werth der Leichenöffnungen. *Ephem. d. Heilk.*, Hamb. and Würzb., 1814, v. VIII, p. 333; Werth der Leichenöffnungen zur Bestimmung Typhus und Hirnentzündung. *Gesammelte med. Progr.*, Würzburg, 1824, p. 38.—FUCHS, C. H. Leichenöffnungen. *Heidelb. klin. Ann.*, 1833, v. IX, p. 50.—FÜHRER. Bericht über die vom 1 Juni 1852 bis zum 1 Jan., 1854 zu Jena angestellten Leichenuntersuchungen. *Deutsche Klinik*, 1854, v. III, p. 181.

GAIRDNER, W. T. S. Post-mortem examinations. *Edinb. med. and surg. J.*, 1841, v. V.—GALSTER. *Gesellsch. d. Aerzte*.

GAIRDNER, W. T. S. The performance of post-mortem examinations. *Edinb. med. and surg. J.*, 1841, v. XI, p. 302.—GANSSET. Post-mortem examinations. *Medical Sciences*, N. Y., 1887.—GANSSET. *Leichenöffnungen*. *Ztschr. d. k. k. Gesellsch. d. Aerzte*, Wien, 1887, v. XIII, p. 181.—GERSICH, A. Az 1900 ik



év fobyamán a Szt István Kórházban végzett bonczolások közül néhány. *Budapest székess. közkeztev.*, 1900-1901, p. 259.—GENOVES, J. Breve reseña sobre la grande importancia de la inspeccion cadaverica. *Bol. de med., cirug. y farm.*, 1880, v. IV, p. 250.—GERBER, B. R. *De legitima cadaveris occisi sectione occasione an.* Halae Magdeb., 1761.—GERICKE, P. *Pr. quo inspectionem cadaveris in hominibus apud Romanos olim in usu fuisse ostendit simulque publicas sectiones et demonstrationes cadaveris foemini.* Helmstadii, 1739; *Programma in quo inspectionem cadaveris in homicidio apud Romanos olim in usu fuisse ostendit.* Helmstadii, 1738.—GERLACH, L. Ueber ein neues Verfahren, kleinere anatomische Objecte zum Zwecke zu fixiren und zu conserviren. *Sitzungsber. d. phys.-med. Soc. zu Erlang.*, 1881.—GLANTURCO, V. *Manuale della tecnica delle autopsie e della diagnostica anatomica pathologica.* Napoli, 1890.—GILBRIDE, J. J. A simple method for closing the body after an autopsy. *Am. Med.*, Phila., 1902, v. IV, p. 768.—GLATTER, E. Ueber Leichenobduction. *Ztschr. f. Nat. u. Heilk. in Ungarn.*, 1851-52, v. II, p. 1.—GOUBERT, *Manuel de l'art des autopsies cadavériques.* Par., 1867.—GRAINGER, T. Analysis of the post-mortem examinations in the Murtridabad district during the past five years. *Indian M. Gaz.*, 1902, v. XXXVII, p. 101.—GRANDJUX, Les autopsies à la guerre, à la marine et aux colonies. *J. de Méd. de Par.*, 1901, v. XIII, p. 239.—GRAUPNER, R., AND ZIMMERMANN, F. *Technik und Diagnostik am Sektionstisch.* Zwickau, 1899.—GREISING, J. E. *De cadaveris inspectione sive sectione legali.* Jena, 1742.—GREEN, C. R. M. Return of sudden deaths requiring coroner's inquests for five years (from 1896 to 1900) in the city of Calcutta. *Indian M. Gaz.*, 1902, v. XXXVII, p. 301.—GREEN, W. A. Synopsis of symptoms and post-mortem appearances in six cases of fever; in five cases of cholera; in two cases of death from exposure to the sun; in one case of death from drinking. *Tr. M. and Phys. Soc. of Bengal, Calcutta*, 1838-1840, p. 25.—GREENFIELD, W. S. Jottings from the post-mortem room. *St. Thomas Hosp. Rep.*, Lond., 1875, v. VI, p. 239.—GRIESINGER, W. Über die Untersuchungs-Methode des Schädel Inhaltes. *Arch. f. Psychiat.*, 1868, v. I, p. 317.—GUITÉRAS, J. Notes upon the lectures on general pathology. Phila., 1890-1891.—GÜNTZ, E. W. *Der Leichnam des Neugeborenen in seinen physischen Verwandlungen nach Beobachtungen und Versuchen dargestellt.* Leipzig, 1827.—GUYOT, C. *De cadaverum sectionibus pathologicis, et recto ex illis ferendo judicio.* Groningae, 1818.—GVOZDEFF, I. First external examination of the cadaver of a grown individual. Kazan, 1887.—HASBRAND, Obductionsbericht und Gutachten über die Ursache des Todes eines bald nach einer erlittenen Misshandlung Verstorbenen. *Arch. f. med. Erfahr.*, Berl., 1818, v. II, p. 167.—HAUCK, F. H. G. *De autopsia conditionibus.* Halis, 1836.—HARTMANN, R. Rede über die Methode des Unterrichtes in der Anthropotomie. Berl., 1873.—HARRIS, R. O. Autopsies and physical examinations. *Boston M. and S. J.*, 1900, v. CXLIII, p. 420.—HARRIS, T. *Post-mortem Handbook for Clinical and Medico-legal Purposes.* Lond., 1887.—HARKE, THOMAS Die Section der oberen Athmungswege. *Berl. klin. Wchnschr.*, July 25, 1899; Ein neues Verfahren die nasen Rachenhöhle mit ihren pneumatischen Anhängen am Leichnam ohne äussere Entstellung freizulegen. *Virchow's Archiv.*, 1891, p. 125, pt. 2; Athmungswege Sectionstechnik. *Centralblatt für Pathologie.*, 1893, v. IV, p. 179.—HAMILTON, D. J. *A textbook of pathology.* Chap. I, v. I (Seco Cadaveris). Lond. and New York, 1889.—HALLER, A. *Pathological observations treating chiefly of dissections of morbid bodies.* Lond., 1756.—HAINES, W. D. Post-mortem examinations. *J. Am. M. Ass.*, 1903, v. XL, p. 442.—HEBB, R. G. Report of post-mortem examinations made in 1887. *Westminst. Hosp. Rep.*, Lond., 1888, v. IV, p. 175; Report of the postmortems made in 1897. *Ibid.*, 1899, v. XI, p. 211; Report of the post-mortem examinations made in 1898. *Ibid.*, 1899, v. XI, p. 205.—HEBENSTREIT, J. E. *De medicis cadavera secantis religione dissertatio.* Lipsiae, 1741; *De corpore delicti.* Lipsiae, 1743.—HERTOEN, L. *The technique of post-mortem examinations.* Chicago, 1896; A series of necropsies held at the Cook County Hospital. *Chicago Clin. Rev.*, 1895-96, v. V, p. 212.—HELLER, Über die Notwendigkeit, die meist übliche Sektions-technik zu ändern. *Verhandl. d. deutsch. path. Gesellsch.*, Jena, 1904, v. XV; See also *Vierteljahrsschrift f. gericht. Mediz.*, 1904, v. XXVII, no. 1, p. 110. (This writer has a book on technique which will soon be in the hands of the printer.)—HENKE, A. Über die Obduktion bereits begrabener und faulender Leichname. *Ztschr. f. d. Stau.*, 1824, v. VII, p. 1.—HERBERT, C. A. Procedure in post-mortem examinations. *Canad. Med. Rec.*, Montreal, 1899, v. XXVII, p. 1; p. 12.—HESCHL, R. *Sections Technik.* Wien, 1859.—HESSELT



*buch für Gerichtliche Aerzte und Wundärzte bei gesetzmässigen Leichenöffnungen, mit der Beschreibung der hierbei nöthigen Instrumente und Geräthschaften.* Giessen, 1819.—HINZE, A. Über das neue Königl. preussische Criminalgesetz, die medizinischgerichtlichen Leichenöffnungen betreffend. *Ztschr. f. d. Staatsarznk.*, Erlang., 1824, v. VII, p. 1.—HIRSCHFELD, L. Le médullotome. *Bull. acad. de méd.*, Par., 1865-66, v. XXXI, p. 7.—HODGKIN. On the object of post-mortem examinations. *Lond. M. Gaz.*, 1828, v. II, p. 423; Holding of post-mortem examinations without notice to company. *J. Am. M. Ass.*, 1903, v. II, p. 805.—HOMMEL. *De inspectione.* Leipzig, 1747.—HUBER, J. M. Het schouwen der lijken naar het standpunt der nieuwere wetenschap, ter voor koming van het levend begraven worden. *N. pract. Tijdschr. v. d. Geneesk.*, Gorinchem, 1853.

IBRAHIM, H. A. *De l'examen du cadavre en médecine légale.* Par., 1869.—INDEX to abstracts of post-mortem examinations. *Middlesex Hosp. Rep.*, 1901. Lond., 1903, p. 122.—INGELS, B. C. Autopsies faites à l'Hospice Guislain. *Ann. Soc. de méd. de Gand.*, 1869, v. XLVII, p. 139; Autopsies pratiquées à l'Hospice. Guislain. *Ibid.*, 1875, v. LIII, p. 215.—INGERLE, S. Bericht über 3 596 während der Jahre 1854-1864 in der Pathol.-anatom. Anstalt zu München vorgenommene Sectionen. *Arb. a. d. path. Inst. zu München.* Stuttg., 1886, p. 403.—INQUESTS at the Radcliffe Infirmary. Oxford, 1893.

JAMAKEWA, K. Necessity of autopsy for studying medicine. *Tokyo Iji Shinshi*, 1898, No. 1030, p. 5.—JAMESON, W. B. A tabular report of 254 autopsies. *Phila. Hosp. Rep.*, 1896, v. III, p. 184.—JANEWAY, E. G. On the necessity of post-mortem examinations. *Med. Rec.*, 1872, v. VII, p. 536; Post-mortem examinations in relation to public health. *Am. Pub. Health Ass. Rep.* 1879. Boston, 1880, v. V, p. 75.—JOHNSTON, W. *Syllabus of post-mortem methods.* Montreal.—JOCON, F. *Comment faire une autopsie.* *Arch. prov. de chir.*, Par., 1894.

KAISERLING, C. Weitere Mittheilungen über die Herstellung möglichst naturgetreuer Sammlungs präparate. *Arch. f. path. Anat.* Berl., 1897, v. CXLVII, p. 389; *Über Konservierung und Aufstellung pathologisch-anatomischer Präparate für Schau und Lehrsammlungen.* München, 1899. Berl., 1900; His method. *Verh. d. pathol. Ges.*, 1900, p. 203; Ueber konservierung von Präparaten für diagnostische und Unterrichtszwecke. *Appar. d. klin. med.* Wiesb., 1901, p. 53.

—KAPATSENSKI, N. On cadaveric spots and differential diagnosis of ante- and post-mortem excoriation and strangulation furrows. St. Petersburg, 1882.—KAST, A. and RUMPEL, T. *Pathologisch-Anatomische Tafeln nach frischen Präparaten mit erläuterndem-anatomisch. klinischen text.* Wandsbik, 1896-97.—KELLY, H. A. On a method of post-mortem examination of the thoracic and abdominal viscera through vagina, perineum, and rectum, and without incision of the abdominal parietes. *Med. News*, 1883, v. XLII, p. 733.—KELYNACK, T. N. *The pathologist's handbook; a manual for the post-mortem room.* Lond., 1899.—KEMPF, E. J. How to make a postmortem. *Am. Pract. and News*, Louisville, 1888, n. s., v. V, p. 392.—KERR, E. K. Statistical study of post-mortem findings in 171 cases of lobar pneumonia. *Trans. Chicago Path. Soc.*, 1901, v. V, p. 274.—KITT, TH. Section der Schädelhöhle. *Jahrb. der kgl. C. Thierarzneischule.* München, 1881-1882.—KITTSTEINER, W. Bericht über 3846 während der Jahre 1865-1874 in der Patholog.-anatomisch. Anstalt zu München vorgenommene Sectionen. *Arb. a. d. path. Inst. zu München.* Stuttg., 1886, p. 432.—KNEPPELHOUT, C. J. *Sectiones cadaverum pathologicae.* Lugd. Batav., 1805.—KOHLHAAS. Eine Bemerkung zu Virchow's Sektionstechnik. *Deutsche med. Ztg.*, 1893, v. XIV, p. 1083.—KOLSTER, R. Seltene Sectionsbefunde. *Arch. f. path. Anat.*, 1899, v. CLV, p. 388.—KÖRBER. *Die Sektionstechnik für neugeborene Kinder.* Dorpat, 1888.—KORNFELD, H. Der zweite Obducent. *Ztschr. f. med. Beamte*, 1901, v. XIV, p. 719; Ueber überflüssige Sectionen. *Arch. f. krim. Anthropol. u. Kriminalist.* Leipz., 1901-1902, v. VIII, p. 192.—KRANNHALS, H. Bericht aus dem allgemeinen Krankenhause zu Riga über die im Jahre 1884, ausgeführten Leichenöffnungen. *St. Petersburg. med. Wchnschr.*, 1885, v. II, p. 385.—KRAUS, L. A. *Praktische Anweisung zu gerichtlichen Leichenuntersuchung.* Helmstadt, 1837.—KRIMER, W. Leichenöffnungen in pathologischer und praktischer Beziehung. *J. d. pract. Heilk.*, 1820, v. LI, p. 3, 1821, v. LII, p. 29.—KRUPP.—Über die Unterbindung der grösseren Gefässe der Brusthöhle bei der gerichtlichen Untersuchung der neugeborenen Kinder. *Allg. med. Ann., Altnb. u. I.*

—KUROKI, T. Autopsies. *Kokka Igaku Kwaishi*

pit

... necroscópicas hechas en el Hos-  
estud. clin. de la Habana, 1882,



- v. II, p. 109.—LAMARCHE, A. *Autopsies cadaveriques*. N. t. p. 1881.—LAMB, D. S. Autopsy, necropsy, necroscopy. *J. Am. M. Ass.*, 1897, v. XXVIII, p. 661.—LAMB, Klinische obductie. *Tijdschr. v. inland. Genees. k.* Batavia, 1898, v. VI, p. 35.—LARIN. Traumatic excoriations produced during life and after death. St. Petersburg, 1894.—LEA, H. Report of fatal diseases and morbid appearances observed in post-mortem examinations made at St. George's Hospital during year 1843. *London M. Gaz.*, 1844, v. II, p. 154.—LEBRUN AND MACUIER. *Nouveau manuel complet du mouleur*. Paris, 1850.—LEFORT, P. Aide-mémoire d'anatomie pathologique, d'histologie pathologique et de technique des autopsies pour la préparation du cinquième examen. Par., 1891.—LEGG, J. W. Report from the post-mortem room. *St. Barth. Hosp. Rep.* Lond., 1875, v. XI, p. 63.—LESHTSHINSKI, M. On the penetration of liquids into the respiratory passages of dead bodies. St. Petersburg, 1868.—LETHEBY, H. On the observance of method in conducting post-mortem examinations, especially when they are intended for legal purposes. *Lancet* Lond., 1846, v. I, p. 1040.—LETTERMAN, J. Autopsies. *Calif. M. Gaz.*, 1869-1870, v. II, p. 100.—LETULLE, M. *La pratique des autopsies*. Par., 1903; Le dispositif instrumental des autopsies. *Bull. et mém. soc. anat. de Par.*, 1902, v. LXXVII, p. 1007; L'autopsie. *Presse méd.*, 1903, v. L, p. 321; Autopsie du carcum. *Presse méd.*, 1904, v. I, p. 1.—LEY. *De frustranea cadaveris inspectione in homicidio dessert.* Helmstadil, 1723.—LIBAVIUS. *Tractatus de cruentatione cadaverum in jure cuade factorum, presente, qui occidisse creditur*. Francfort, 1594.—LIEBTAUD, J. *Historia anatomica medica*. Par., 1767.—LITTERSKI, BLEICH, BECK, VOLLMER, MULERT UND KLOSE. Der zweite Obducent. *Ebend.* No. 24.—LITTLE, JOHN, H. D. Post-mortem examination. *Edinb. M. J.*, 1876, v. XXI, p. 673.—LOBEDANK. *Hilfsstafel zum Gebrauch bei Sektionen und zur Abfassung des Sektionsprotokolls*. Leipzig, 1903.—LOGAN, T. M. Data for calculating the lapse of time since the death of an individual. *Pacific M. and S. J.*, 1863, v. VI, p. 38.—LOPES VIEIRA. Precaucoes adoptaveis n'as autopsias de cadaveres euz putrefacção. *Coinbra med.*, 1899, v. XIX, p. 354.—\*LORENZ. *Obduktionsscheern bei Verdacht auf Vergiftung*. Berl. N. d.—LÖWE, Ueber eine neue Sektionsmethode der Nasenhöhle. *Virchow's Archiv*, 1901, v. 163, p. 179.—LUND, E. A form of craniotomy for post-mortem examinations. *Lancet*, Lond., 1858, v. II, p. 520.—LUTAUD, A. Les autopsies dans les hôpitaux; à M. Mourier, Directeur de l'assistance publique. *Rev. de méd. leg.*, Par., 1901, v. VIII, p. 208.—LUYS, J. Procédé pour la conservation et la momification des cerveaux à l'état sec. *Encéphale*, 1881, p. 84; De divers modes de préparation et de durcissement du cerveau. *Bull. acad. de méd.*, 1886, v. XVI, p. 271.—LYMAN, H. M. The Mutual Autopsy Society of Paris. *Chicago M. J. and Exam.*, 1879, v. XXXIX, p. 373.
- MAGNUS. Signs of death. *Virchow's Archiv*, 1872.—MAHON, P. A. O. *An essay on the signs of murder in new-born children*. Lancaster, 1813.—MALLORY AND WRIGHT. *Pathological Technique*. Phila., 1897.—MARCHAND. Autopsie crânienne et rachidienne. *Revue de Psychiatrie et de Psychologie expérimentale*, 1903, v. XV, p. 343.—MARIQUE AND DALLEMAGNE. *Revue du service des autopsies*. *J. de méd., chir. et pharmacol.*, 1886, v. LXXXIII, p. 534.—MARTIN, R. J. A. Der Schädelhatter, ein Beitrag zur Vervollständigung des Sections-apparats. *Mag. f. a. Staatsarznk.* Leipzig, 1844, p. 248.—MAUCLAIRE AND MESUREUR. Les autopsies dans les hôpitaux. *J. méd. de Par.*, 1903, s. 2, v. XV, p. 204.—MCCAW, J. D. Crâne-agrafe for holding the skull while sawing off the calvaria. *Lancet*, Lond., 1883, v. I, p. 508.—MCLEOD, K. Summary of 50 post-mortem examinations of inhabitants of the Jessore district. *Indian Med. Gaz.*, 1868, v. III, p. 205.—MECKEL, J. F. *Tabulæ anatomico-pathologicae*. Lipsiæ, 1817.—MEHNERT, E. Bericht über die Leichenmessungen am Strassburger Anatomischen Institut. *Morphol. Arb.* Jena, 1894, v. IV, p. 1.—MEHRERE. Über die gerichtlichen Leichenöffnungen betreffende Punkte. *Mag. f. d. gerichtl. Arzneik.*, 1832, v. I, p. 18.—MERBACH. Pathologisch-anatomischer Bericht über zweihundert im Dresdener Krankenhause angestellte Sectionen. *J. d. Chir. u. Augenh.* Berl., 1850, v. XXXIX, p. 123.—MERZ, C. H. A synopsis of the various points of interest in post-mortem examinations. *Med. Age*, 1893, v. XI, p. 1.—MESA, E. Sobre las autopsias judiciales. *Corresp. méd.* Madrid, 1898, v. XXXIII, p. 380.—MEYER, A. Preparations for a post-mortem examination. *Chicago Clin. Rev.*, 1893, v. II, p. 32.—MEYNERT. Über die methode der Gehirnwiegunen. *Mittheilungen der Anthropologischen Gesellschaft*, No. 5; Das Gesamtgewicht und die Theilgewichte des Gehirns in ihren Beziehungen zum Geschlecht, dem Lebensalter und dem Irrsinn, untersucht nach einer neuen Wiegunsmethode. *Vlljschr. f. Psychiat.*



1867, pt. II.—MILLS, T. *An account of the morbid appearances exhibited on dissection in disorders of the trachea, lungs, and heart, with pathological observations.* Dublin, 1829.—MINOR, W. C. *Post-mortem examinations made at Knight U. S. A. General Hospital.* New Haven, 1864.—MIURA, S. Method of holding postmortems. Tokyo, 1894.—MOLCHANOFF, V. A. *Study of cadaveric spots.* St. Petersburg, 1894.—MORACHE, G. *Naissance et mort.* Par., 1904.—MORGAGNI, J. B. *The seats and causes of disease, investigated by anatomy: containing a great variety of dissections and accompanied with remarks.* Lond., 1822; Boston, 1824.—MORRIS, M. The method of making autopsies and their proper interpretation. *Internat. M. Mag.*, 1894-95, v. III, p. 825.—MOWAROWSKI, J. F. Autopsy in a case of murder. *Gaz. lek.*, Warszawa, 1879, v. XXVII, p. 61.—MUJASIRO. Post-mortem examinations should be made obligatory for the attending physician. *Torin no Siawori, Tokio*, 1893, No. 9, v. V, p. 10.

NASSE. Leichenöffnungen. Bonn, 1821.—NAUWERCK, C. *Sectionstechnik.* Jena, 1904.—NEGRESCU. Studii asupra unui numar de autopsii practicate im Judetul Ramnicul Sarat. *Bul. asoc. gen. a. med.*, Bucuresci, 1899-1900, v. III, p. 150.—NEUMANN. 50 Krankheits u. Sectionsgeschichten von Wahnsinnigen mit vergleichenden Resultaten. *J. de pract. Heilk.*, 1824, v. LIX, p. 3.—NEW MORTUARY and post-mortem room at the Middlesex Hospital. *Lancet*, Lond., 1889, v. II, p. 1247.—NEWTH, A. H. *Postmortems.* Detroit, 1897; *A Manual of Necroscopy or a Guide to the Performance of Post-mortem Examinations.* Lond., 1878.—NOBILING, A. Zur Sectionstechnik. *Friedreich's Bl. f. gerichtl. Med.*, Nürnberg, 1893, v. XLIV, p. 53.—NOIR, J. Les autopsies dans les hôpitaux. *Progrès méd.*, 1901, v. XIII, p. 91.—NORTH CAROLINA STATE BOARD OF HEALTH. *Method of Performing Post-mortem Examinations.* Raleigh, 1879.

ÖFFNUNG der Kopfhöhle häufig vorkommende Vernachlässigung der erforderlichen praktischen Kautelen und die daraus entstehenden Nachtheile. *Mag. f. d. gerichtl. Arzneiw.*, 1832, v. II, p. 151.—OGSTON, F. Pathological studies on the bodies of the drowned. *Lond. M. Gaz.*, 1851, n. s., v. XVII, p. 288.—OLLIVIER D'AUERS. *Considérations médico légales sur certaines productions résultant de la décomposition des cadavres et qui peuvent, dans quelques cas, aider à découvrir la cause de la mort.* Par., 1819.—ORTH, J. *Compendium der pathologisch-anatomischen Diagnostik nebst Anleitung zur Ausführung von Obductionen.* Berl., 1884-1900. (Prof. Orth informs the writer that a new edition of his work will not appear for several years.)—OSTAPENKO, A. P. Examination of bodies of domestic animals. *Vet. Vestnik*, Charkoff, 1891, v. IX, p. 1.—ORTO. Sectionsberichte mit vergleichender Zusammenstellung der eigenthümlichen der pathologisch-anatomischen Veränderungen der Organe gegenüberstehenden Krankheitserscheinungen. *Arch. d. deutsch. Gessellsch. f. Psychiat.*, Neuwied, 1858, v. L, p. 64.—QUELLY, P. M. A. *De l'indispensable utilité des nécroscopies dans la pratique particulière dans l'intérêt des familles, du médecin et de la science en général, par rapport aux maladies héréditaires.* Par., 1833.—OUGHTON, T. On the Necrometer. *Army M. Dep. Rep.*, Lond., 1871, v. XI, p. 520.

PACKARD, F. R. Method of replacing skull-cap after post-mortem examination so as to prevent disfigurement. *Proc. Path. Soc.*, Phila., 1860, v. II, p. 62.—PARK, R. On a new method of making anatomical preparations. *Ann. Anat. and Surg.*, v. V, 1882.—PASQUIER. Emploi des ciseaux de Brunetti pour l'ouverture du rachis, dans les autopsies. *Bull. méd. du nord*, 1882, v. XXI, p. 383.—PAVENTA. Moduli per autopsie. *Gazz. med. di Torino*, 1885, v. XXXVI, p. 464.—PENTA, P. Alcune note su 35 autopsies di condannati. *Rev. mens. di psichiat. fienese*, Napoli, 1902, v. V, p. 425.—PERTIK, O. Singular cases of Pathological Anatomy observed in 1900 by the Prosecutor at St. Rok's Hospital. *Budapest székcs f. Köz Körhczek*, 1900-1901, p. 161.—PETERSON, F. Report of the pathological results of 215 post-mortem examinations. *Buffalo M. and S. J.*, 1884-85, v. XXIV, p. 212.—PICK, T. P. Pathological observations. *St. George's Hosp. Reports*, Lond., 1868, v. III, p. 277.—PIETRE, JEAN. Au quel e trucidati vulnere antea sicco manat sanguis, prodeum te etiam sicario, sola effluit putredine affirmavit. *Diss. resp. Le tour-neurs*, Par., Nov. 16, 1634.—PINNOCK, R. A. A plea for legislation to empower medical practitioners to make post-mortem examinations in cases of doubtful diagnosis. *Intercolon. M. Cong. Australas. Tr.*, 1899, Brisbane, 1901, p. 495.—PLANER, J. Statistik der Leichenkammer des k. k. allgem. Kraukenhauses vom Monat November, 1854. *Wechnbl. d. Zeitschr. d. k. k. Gesellsch. d. Aerzte in Wien*, 1855, v. I, p. 171.—PODHAJSKY, V. The post-mortem. *Ootvirani mrtvol. Cásop. lek. cesk.*, v. Praha, 1869, v. VIII, p. 121.—POKROVSKI, M. Handbook for



- performing autopsies. Moskva, 1901.—POLIZEI verordnung, betreffend die Öffnung der Särge vor dem Versenken. *Preuss. med. Ztg.*, Berl., 1864, n. f. v. VII, p. 10.—POSTMORTEM. *J. A. M. A.*, 1903, v. II; Post-mortem inspection of head, *See* M. J., 1871, v. II, p. 262.—\*PRAUSNITZ, Zur Sectionstechnik des Herzens.—PRESCOTT, A. C. Remuneration for post-mortem examinations. *Tr. M. Soc. Wisc.*, 1897, v. XXXI, p. 71.—PREUSS, F. W. *Über die Altersbestimmung an Leichen*. Berl., 1895.—PRIVATSECTIONEN. *Allg. med. Ann. Altenb.*, 1807, p. 105.
- RAYNAU, T. *De incorruptione cadaverum, occasione de mortui foeminae corporis post aliquot secula incorrupti, nuper refossi Carpentoracti*. Aravico, 1651.—REID, J. Analysis and details of forty-seven inspections after death. *Edinb. M. and S. J.*, 1839, v. LII, p. 448.—RICHARDSON, B. W. Attempt to restore to its natural appearance putrefied dead body in order to prove its identity. *Lancet*, Lond., 1863, v. I, p. 350; Temporary preservation of post-mortem specimen. *Asclepiad*, Lond., 1864, v. I, p. 156.—RICHARDSON, J. G. New method of preserving tumors and certain urinary deposits during transportation. *Phil. M. Times*, 1872-73, v. III, p. 181.—RICHTER, J. C. *Gerichtsärztliche Obductionen Tabellen*. Lüneburg, 1841.—RICHTER, M. Ueberflüssige Sectionen. *Arch. f. Krim. Anthropol. u. Kriminalist.* Leipz., 1902, v. IX, p. 353.—RIGOT AND TROSSEZ. Recherches nécroscopiques sur quelques altérations que subissent après la mort les vaisseaux sanguins, les poumons et la membrane muqueuse gastro-pulmonaire à l'état sain. *Arch. gén. de méd.*, Par., 1826, v. XII, p. 169.—RIVERO, P. G. Conferencias sobre autopsias. *An. de San. Mel.*, Buenos Aires, 1900, v. II, p. 103.—ROBINSON, B. Observations on the Abdominal Viscera in 17 Autopsies. *Med. Rev.*, 1895, v. XXXI, p. 101; Inspection of the abdominal cavity and its viscera in postmortems. *Am. Surg.*, Phila., 1899, v. XXIX, p. 731.—\*ROHATZSCH, J. G. Taschenbuch für gerichtliche Sectionen und Gutachten zum gebrauchte practischer Aerzte und Wundärzte.—ROKITANSKY, C. *A Manual of Pathological Anatomy*. Lond., 1848, Phila., 1855; Handbuch der speciellen pathologische Anatomie. Wien, 1844; *Lehrbuch der pathologischen Anatomie*. Wien, 1855, 1861; Statistik der Sanitätspolizeilichen und landesgerichtlichen Leichenschauen in den Monaten Juli, Aug., Sept., und Oct., 1855. *Wechnbl. d. Ztschr. d. k. k. Gesellsch. d. Aerzte*. Wien, 1856, v. II, p. 346; Statistik der Leichen Kammer des k. k. Allgem. Krankenhauses vom October, 1854, und zwar der pathologischen Sectionen. *Ibid.*, 1855, v. I, p. 40.—ROLLESTON, H. D., AND KANTHACK, A. A. *Manual of practical morbid anatomy, a handbook for the post-mortem room*. Cambridge, 1894.—ROMANOFF, F. J. Result of the pathological necropsies performed in the University of Tours during 1890 and 1900. *Izviest Imp. Toursk Univ.*, 1902, v. XIX, p. 1.—ROMBERG. Ergebnisse einiger Leichenöffnungen. *Arch. f. med. Erfahr.* Berl., 1822, v. II, p. 379.—ROOSE, T. G. A. Taschenbuch für gerichtliche Aerzte und Wundärzte bei gesetzmässigen Leichenöffnungen. Frank.-a.-M., 1819.—RUDALL, J. Report of post-mortem examinations at the Yarra Bend Asylum, 1861-1862. *Austral. M. J.*, Melbourne, 1862, v. VI, p. 252.—RUGE, G. *Anleitungen zu den Präparierübungen an der menschlichen Leiche*. Leipzig, 1896.
- VON SAMSON HIMMELSTEIN, G. Zusammenstellung der pathologisch-anatomischen Sectionsbefunde am Dorpat'schen Kreishospital aus dem Jahre, 1847. *Beitr. z. Heilk.*, Riga, 1851, v. I, p. 152.—SANDIFORT, E. *Observationes Anatomico-pathologicae*. Lugd. Bat., 1776.—SANGALLI. La metodica sezione del corpo umano alla ricerca delle sue alterazioni. *Gior. di anat. e fisiol.*, Pavia, 1864, v. I, p. 135; 1865, v. II, p. 493; Metodo della sezione degli organi della cavità abdominale. *Ibid.*, 1865, v. II, p. 616.—SANITÄTS-POLIZEILICHEN Leichenobductionen. *Oester. San. Wes.* Wien, 1895, v. VII, p. 183.—SAWYER, T. C. One hundred autopsies. *Rep. St. Lawrence State Hosp.*, 1894, Albany, 1895, v. VIII, p. 159.—SCARENZO, A. Nuovo metodo per ben sezionare lo speco vertebrale. *Gazz. med. ital. Lomb.* Milano, 1860, v. IV, p. 265.—SCHÄLLE. Eine neue Sectionsmethode für die Nasen-, Rachen-, und Gehörorgane. *Arch. f. path. Anat.*, 1877, v. LXXI, p. 266.—SCHERER, M. Zur Section des Nasen-Rachensaums. *Virchow's Archiv*, 1899; Schema zum Einzeichnen pathologischer Befunde der Brust- und Baucheingeweide. *Bern.*, 1891.—SCHINZ, C. S. *Tractatio de cauto sectionum cadaverum usq. ad dijudicandas morborum causas*. Gottingae, 1787.—SCHLOCKOW-ROTH-LEPPMANN. *Der Kreisarzt*. Berlin, 1900.—SCHMALZ, R. *Atlas über die Lage der Eingeweide des Pferdes*. Berl., Verl. v. Schotz.—SCHOTTILIUS. Neun Sectionstafeln. Wiesbaden, 1878.—SCHWENINGER, E. Bericht über die Leichenöffnungen in dem unter Präs. v. Buhl's Leitung stehenden pathologischen Institute pro 1876 u. 77. *Arzt-Städt. Allg. Krankenk. zu München*, 1876-77, p. 348.—SÉE, M. Autopsi



- encycl. d. sc. méd.* Par., 1867, v. VII, p. 411.—SELDEMAN. *De cadavere inspiciendo*. Groningue, 1692.—SEVERI, A. Di un modo rapido per determinare all' autopsia il peso specifico dei vari organi e tessuti. *Sperimentale*, Firenze, 1882, v. XLIX, p. 580.—SLEE, R. Simple method of retaining the calvarium in place after post-mortem examination. *Med. News*, 1892, v. LXI, p. 737.—\*SMETI, Po. *Casop. lek. cesk.* After death. v. *Praxe*, 1873, v. XII, p. 7; Société d'autopsie. *Rev. mens. de l'école d'anthrop. de Par.*, 1893, v. III, p. 233.—SOUTHACK, J. W. Method of making autopsies at the pathological Institute of Vienna. *Oregon M. and S. Reporter*, 1869-1870, v. I, p. 14.—SPITTA, H. *Die Leichenöffnung in Bezug auf Pathologie und Diagnostik*. Stendal, 1826.—SPOTT, C. *Methodus sectiones cadaverum scopo pathologico instituendi*. N. p. 1842.—STAUPA, J. M. *Sistens methodum cadavera humana rite secandi*. Vindobonæ, 1826.—STECKSÉN, A. *Kort handledning i obduktionsteknik*. Stockholm, 1898; Eine Darmseziermethode. *Arb. a. d. pathol.-anat. Institut zu Tübingen*, v. III, 1901.—STEINBERG, S. Ueber den Härtemesser des Gehirns encephalosclerometer. *Arch. d. Heilk.*, 1865, v. VI, p. 524.—STEINER, W. R. Some early autopsies in the United States. *Johns Hopkins Hosp. Bull.*, 1903, v. XIV, p. 201.—STEWART, W. D. Notes on post-mortem examinations conducted in the Porree district. *Indian M. Gaz.*, 1872, v. VII, p. 169.—STEIBEL, J. Leichenbefunde aus dem Kinderhospitale zu Frankfurt. *M. J. f. Kinderkr.*, 1851, v. XVI, p. 364.—SUCHARD, E. Technique générale et sommaire des autopsies cliniques. *Clin. méd. de la Charité*. Par., 1894, p. 1011.
- THACHER, J. S. The pathological conditions found at 1000 autopsies. *Med. and Surg. Rep. Presby. Hosp. N. Y.*, 1902, v. V, p. 170.—THOMAS, A. R. *A practical guide for making post-mortem examinations, and for the study of morbid anatomy*. New York, 1873.—THOMAS, M. Comparative merits of investigating sudden death in England and Scotland. *Glasgow M. J.*, 1857-58, v. V, p. 281.—TILING, T. Sectionsergebnisse in der Irrenheilanstalt. *St. Petersburg. med. Ztschr.*, 1875, v. V, p. 171.—TOMES, A. Postmortems and laws. *Indian Med. Gaz.*, 1880, v. XV, p. 145.—TOTT, C. A. Resultate einiger Leichenöffnungen und Leichenbesichtigungen. *Jahrb. d. ges. Staatsarznh.*, Leipz., 1835, v. I, p. 91.—TOURDES, G. Autopsies. *Dict. encycl. d. sc. méd.* Par., 1867, v. VII, p. 417.—TRETYAKOFF, N. *Brief description of post-mortem examinations*. St. Petersburg, 1897.
- UNITED STATES. War Department, Surgeon-General's Office. Form of Record of Autopsy. (Extract from a report of Dr. Bowditch to the Mass. Med.-Leg. Soc., Feb. 1, 1882, published for medical officers of the Army.) Washington, 1882.
- VALENTIN, G. *Die kunstgerechte Entfernung der Eingeweide des menschlichen Körpers*. Frankfurt-a.-M., 1857.—VALERY, CH. *Tableaux synoptiques pour la pratique des autopsies*. Par., 1902.—VALUE of post-mortem examinations. *J. Am. M. Ass.*, 1903, p. 1425.—VAN DUYSE. Technique des autopsies. *Belgique méd.*, 1896, v. III, p. 673; Premiers éléments de diagnostic anatomo-pathologique au cours des autopsies. *Ibid.*, 1897, v. IV, p. 263; Autopsie de la cavité encephalo-rachidienne. *Ibid.*, 1897, v. IV, p. 455.—VERGA, A. *Delle autopsie Loria nel 1894*. R. I. L., 1895, v. XXVIII, p. 645.—VIRCHOW, R. *A description and explanation of the method of performing post-mortem examinations in the dead-house of the Berlin Charité Hospital, with special reference to medico-legal practice*. Lond., 1876; *Die Sectionstechnik im Leichenhause des Charité Krankenhauses*. Berl., 1874, 1893; *Lernen und Forschen*. Berl., 1892; *Post-mortem examinations with especial reference to medicolegal practice*. Phila., 1880; *Deutsche Klinik*, No. 2, 1859.—VISCONTE, A. *Le necroscopie Loria nel 1896-1897*. Milano, 1898.—VISITE, S. *Necroscopiche*. *Indipendente*. Torino, 1874, v. XXXV, p. 1.—VOLLMER. Sections Bericht und Gutachten über die Todesart der A. G. D. zu C., etc. *Ztschr. f. d. Staatsarznh.* Erlang., 1845, v. I, p. 405; 1847, v. LIV, p. 373.—VOPPEL. Die sections Ergebnisse aus der Irren-Versorgungs-Anstalt zu Corditz. *Ztschr. f. klin. Med.*, Bresl., 1856, v. VII, p. 161.
- VAN WALSEM. Eröffnung der Schädelhöhle. *Virchow's Arch.*, v. CLXIII, 1901, and v. CLXX, 1902.—WALSH, J. P.—Some notes on autopsies. *Med. and Surg. Rep.*, 1896, v. LXXV, p. 167.—WELLS, S. R. AND WILSON, W. H. On post-mortem nerve changes. *J. Path. and Bact.*, 1894-96, v. III, p. 482.—WENDELL, C. A. Post-mortem examinations. *Boston M. and S. J.*, 1865, v. LXXI, p. 111.—WETZLER. Leichenöffnungen. *Jahrb. d. Med. u. Chir.*, 1825, v. IX, p. 2.—WHAT to observe at the bedside and after death in medical cases. *London Med. Soc. of Observation*, Lond., 1853.—WOODHEAD, G. S. *Practical Pathology*. Edinburgh and London, 1892.—WOODS, G. W. Details of necropsies. *Rep. Surg. Gen. Navy*. Wash. 1896, 1897, 1898.



—WRANY, A.—Sectionsergebnisse an der Prager pathologisch-anatomischen Anstalt. *Verh. d. prakt. Heilk.* Prag, 1867, v. XCIV, p. 1.

YAMAKIWA, K. Necessity of autopsy for studying medicine; result of the second session of autopsy; a case of autopsy interesting in pathology. *Tokyo Iji Shinshi*, 1898, no. 1030, p. 5, nos. 1031 and 1032, pp. 13-22.

ZUCKERKANDL, E. *Anleitungen für den Secirsaal*. Wien u. Leipzig, 1897. *Normale und pathologische Anatomie der Nasenhöhle und ihrer pneumatischen Anhangs*. Wien, 1882.

#### Medicolegal References

- ABERCROMBIE, J. *Student's guide to medical jurisprudence*. Phila., 1885.—AGUGLIA, F. *Moduli per autopsie medico-legali*. *Gazz. med. di Torino*, 1885, t. XXXVI, p. 313.—ALBERTI, M. *Commentatio in constitutionem criminalem Carolinam medicam*. Halle, 1739; *Systema jurisprudentia medicæ*. (Bibliography.) Halle, 1736-1747.—ALLEN, I. J. *Address introductory to the lectures on medical jurisprudence*. Willoughby, Ohio, 1846.—AMMAN, P. *Medicina critica*. Erlang., 1870.—ANDRADE, A. *De algunas aplicaciones del microscopio en las pesquisas médico-legales*. Mexico, 1870.—ANGLADA, J. *Traité de toxicologie générale*. Par., 1835.—ANTONIO, R. *Sul progetto di legge per i medici periti giudiziari*. Napoli, 1898.—APPEL, J. M. *Jura aegrotantium*. Erfordia, 1719.—ARDINI, G. *Prelezione al corso di medicina legale*. Catania, 1886.—AUGUSTIN, F. L. *Repertorium für die öffentliche und gerichtliche Arzneiwissenschaft*. Berl., 1810.—AUSIAUX, S. *Oratio de medicinae forensis historia ejusque dignitate*. Liège, 1822.—AUTENRIED, G. *De judicio medici forensis saepe dubio*. Vetebergae, 1798.—VON AUTENRIED, J. H. F. *Anleitung für gerichtliche Aerzte*. Tübingen, 1846; *With von AUTENRIED, H. F. Gerichtlich-medizinische: Aufsätze und Gutachten*. Tübingen, 1876.—B, J. E. L. *Influence de la médecine légale sur la morale et sur le jury*. Par., 1819.—BABES, V. *Bacteriologia si estologia pathologica in medicina legala*. Romania méd., Bucarest, 1898.—BAIER, J. J. *Introductio in medicinam forensis et responsa ejusdem argumenti*. . . Francof. et Lipsiae, 1748.—BAKKER, G. G. *Momenta quaedam medicinae forensis et physiologia et observatione recentiori illustrata*. Groningae, 1825.—BALK, D. G. *Versuch einiger Umrissse der philosophisch-medizinischen Jurisprudenz*. Dorpat, 1803.—BARRETO, L. *Prontuario de medicina legal y jurisprudencia médica*. Bogota, 1890.—BARTLEY, O. W. *A treatise on forensic medicine or medical jurisprudence*. Bristol, 1815.—BARZILLOTTI, G. *Medicina legale secondo lo spirito delle leggi civili e penali regnanti nei governi d'Italia*. Pisa, 1819; 1840.—BAUDRY, S. *Etude médico-légale sur les traumatismes de l'œil et de ses annexes*. Par., 1904.—BAUMER, J. W. *Medicina forensis*. Francof. et Lipsiae, 1778.—BAUMERT, G. *Lehrbuch der gerichtlichen Chemie*. Braunschweig, 1889.—BAVARIA, KÖNIGLICHE STAATSMINISTERIUM DER JUSTIZ UND DES INNERN. *Instruktion . . . bei den gerichtlichen Untersuchungen menschlicher Leichen*. 3rd edit. München, 1897.—BAYARD, H. L. *Etudes pratiques en médecine légale*. Par., 1840; *Manuel pratique de médecine légale*. Par., 1844.—BAYNES, C. R. *Hints on medical jurisprudence*. Madras, 1854.—BEAMTET. *Der Arzt und ärztliche Sachverständige*. Berl., 1900.—BECK, T. R. *Elements of medical jurisprudence*, 1823; *With Beck, J. B.*, Albany, 1835; Phila., 1838.—BECK, J. B. *Researches in medicine and medical jurisprudence*. N. Y., 1835.—BECKER, R. *Sammlung gerichtsarztlicher Gutachten*. Berl., 1892.—BEER, H. *Ueber die Bedeutung der gerichtlichen Medizin für Juristen*. Wien, 1859.—\*BEHRENS, C. B. *Medicus legalis oder gesetzmässige Bestellung der Arzneykunst*. Francofurt et Leipzig, 1696.—BELING, B. E. *Geist der Preussischen Gesetzgebung im Gebiete der gerichtlichen Medizin*. Breslau u. Leipzig, 1819.—BELL, C. *Medico-legal studies*. N. Y., 1889-1898; *Inaugural addresses as president of the Medico-legal Society of the City of New York*. N. Y., 1872; 1873; 1874.—BELL, J. *An address introductory to a course of lectures on the institutes of medicine and medical jurisprudence*. Phila., 1829; *Syllabus of a course of lectures on the institutes of medicine and medical jurisprudence*. Phila., 1829.—BELL, J. S. *The use and abuse of expert testimony*. Phila., 1879.—BELLENGER, G. D. *Considérations médico-légales sur les maladies spontanées qui peuvent être confondues avec l'empoisonnement aigu produit par les substances vénéneuses irritantes ou corrosives*. Par., 1830.—BELLINI, R. AND FELIPPI, A. *Biblioteca medico-legale*. Pisa, 1877-78.—BELLOC, J. J. *Cours de médecine légale théorique et pratique*. Par.,



- 1811.—BEMIS, G. *Report of case of J. W. Webster*. Bost., 1850.—VON BENE, F. *Elementa medicina forensis*. Budæ, 1811.—BENEDIKT, M. *Anatomical studies upon brains of criminals*. N. Y., 1881.—BENOIT. *De l'empoisonnement criminel en général*. Thèse de Lyon, 1888; *Anleitung zur Abfassung medicinisch gerichtlicher Fundscheine und Gutachten*. Wien, 1821; *Beiträge zur gerichtlichen Arzneikunde*. Wien, 1818-1823; *Systematisches Handbuch der gerichtlichen Arzneikunde*. Wien u. Triest, 1817; *Vis a reperta und gerichtlich medicinische Gutachten*. Wien, 1827-1845.—BERGERON, H. *Médecine légale et jurisprudence médicale*. Par., 1898-1904.—BERGMANN, C. *Lehrbuch der Medicina forensis für Juristen*. Braunschweig, 1846.—\*BERNHOLD, J. M. *Medicina legalis Teychmey-eriana tabulis expressa*. Jena, 1760.—\*BERNSTEIN, C. T. J. *Beiträge zur Wund- arzneikunst und gerichtliche Arzneigelehrtheit*. Coblenz u. Frankf., 1804-1812.—BERNT, J. *Ergänzungen des systematischen Handbuches der gerichtlichen Arznei- kunde*. Wien, 1826.—BERTRAND, C. A. H. A. *Manuel médico-légal des poisons introduits dans l'estomac, et des moyens, etc.* Par., 1817.—BETOUILLERES, J. M. G. *Psychologie de l'expertise médico-légale*. Bordeaux, 1900.—BETTI, P. *Studi di medicina pubblica*. Firenze, 1860-1862.—BLALET, J. *Lecciones de medicina legal aplicada á la legislación de la República Argentina*. Córdoba, 1885.—BIBLIOTHEK der gesammten medicinischen Wissenschaften für praktische Aerzte und Special- ärzte. Wien u. Leipz., 1899.—\*BICHAT. *Recherches sur la vie et la mort*.—\*BIERMANN, J. C. A. *Abhandl. naturhistor, gerichtl. und med. Inhalts*. Leipz., 1828.—BIESSY, C. V. *Manuel pratique de la médecine légale*. Lyon, 1821.—BILLIARD, C. R. C. *Considérations médico-légales sur l'empoisonnement par les irritants*. Par., 1820.—\*BISCHOFF, D. *Merkwürdige Criminalrechtsfälle für Rechte, Gerichtsärzte, Vertheidiger und Psychologie*. Hannover, 1833-1835.—BISCHOFF, C. H. E. *Grundriss einer anthropologischen Propädeutik zum Studio der gerichtlichen Medicin*. Bonn, 1827; *Ueber die Bedeutung und des Erford- erniss der gerichtlichen Medicin, als wesentlichen Bestandtheiles rechtswissen- schaftlicher Studien*. Bonn, 1826.—BITSCH, W. *De Hippocrate Justiniano, seu mutuo medicinarum inter et jurisprudentiam nexu*. Argentorati, 1727.—BIZZAZERO, G. *Manuale di microscopia clinica, con aggiunte risguardanti gli esami chimici più utili al pratico e l'uso del microscopio nella medicina legale*. Milano, 1882; Bruxelles, 1883.—BLACKMER, R. C. *Students' manual of medical jurisprudence*. St. Louis, Mo., 1897.—BLYTH, A. W. *Poisons their effects and detection*. Lond., 1895.—BOCK, A. C. *Gerichtliche Sectionen des menschlichen Körpers*. Leipz., 1843, 1852.—BOCKER, F. W. *Die Vergiftungen in forensischer u. klinischer Beziehung*. Iserlohn, 1857; *Memoranda der gerichtlichen Medicin*. Iserlohn u. Elberfeld, 1853-54; *Lehrbuch der gerichtlichen Medicin*. Iserlohn, 1857.—\*BOERNER, F. *Institutiones medicina legalis*. Wittenberg, 1756.—\*BOHN, J. *De officio medicæ duplici, clinico niniiorum ac forensi*. Leipz., 1704; *Specimen primum et tertium medicina forensis*. Lipsie, 1690-92.—BONALUMI, G. *Manuale di medicina legale militare*. Firenze, 1891.—BORGES, J. F. *Institucoes de medicina forense*. Lisboa, 1840.—BORMANS, T. *Répertoire belge de législation, d'instruc- tions, de doctrine et de jurisprudence concernant la médecine légale, l'exercice de l'art de guérir et la police sanitaire; etc.* Bruxelles, 1882.—BORNTREGER, J. B. *Compendium der gerichtsarztlichen Praxis*. Leipz., 1896.—BORRI, L. *Nozioni elementari di medicina legale*. Milano, 1900.—BOSTOCK. *A vindication of the opinions delivered in evidence by medical witnesses for the Crown on a late trial at Lancaster for murder*. Liverpool, 1808.—BOTTO, C. A. *Segunda cadeira de clinica medica*. Rio de Janeiro, 1884.—BRACH, B. *Lehrbuch der gerichtlichen Medicin*. Köln, 1850.—BRANDAU, C. W. R. *Einige Bemerkungen über das Ver- hältniss der gerichtsarztlichen Thätigkeit zur Richtlichen, sowie über Zurechnungs- fähigkeit und deren Princip mit Rücksicht auf den in Kurhessen geltenden Gerechtsgebranch*. Marburg, 1858.—BRENDL, J. H. *De medico ex voluntate aegroti perperam curante*. Wittenberg, 1719.—\*BRENDL, J. G. *Institutiones medicina legalis*. Halle, 1768; *Medicina legalis sive forensis*. Hannoverae, 1789.—\*BRESCHET, G. *Rapp. méd.-légale sur une plaie d'arme à feu*. Par., 1826.—BRIAND, J. *Manuel de médecine légale*. Par., 1821; *Manual completo de medicina legal*. Madrid, 1872-1873; With BROSSON, J. X. *Manuel complet de médecine légale*. Bruxelles, 1830-1858; With CHANDL, E. *Manuel complet de médecine légale*. Par., 1846; 1874.—BRIERRE DE BOISMONT, A. F. *Manuel de médecine légale*. Par., 1835.—VAN DEN BROECKE, J. C. AND P. *De uitoeffening der geregtelijke geneeskunde in Nederland, hare gebreken, middelen tot herstel derzelve*. Utrecht,



- 1845.—BROUARDEL, P. *La mort et la mort subite*. Par., 1895; English translation by BENHAM, New York, 1897; *Le service des autopsies et de l'expertise médicale légale en Allemagne et en France*. Tribune méd. Par., 1879, v. XII, p. 11. *L'infanticide*. Par., 1897; *Les explosifs et les explosions au point de vue médical*. Par., 1897; *La responsabilité médicale*. Par., 1898; *La pendaison, la strangulation, la suffocation, la submersion*. Par., 1887; *Les empoisonnements criminels et accidentels*. Par., 1902.—BROWNE, G. L. AND STEWART, C. G. *Report of trials for murder by poisoning by prussic acid, strychnia, arsenic, and acetic acid. Including the trials of Tawell, W. Palmer Dove, Madeline Smith, Dr. Prichard, Smethurst, and Dr. Lamson*. Lond., 1883.—BUCHNER, E. *Lehrbuch der gerichtlichen Medicin für Aerzte und Juristen*. München, 1867; 1872; Welche Verletzungen sind nothwendig tödtlich? Erlangen, 1861.—BUCHOLZ, W. H. S. *Beiträge zur gerichtlichen Arzneigelahrtheit und zur medicinischen Polizei*. Weimar, 1782.—BUDAEUS, G. *Miscellanea medico-chirurgica, practica et forensia*, etc. Leipzig, 1732–37.—\*BUKHSHTAB, B. G. Tr. On various changes in the legal chemical examination of internal organs. *Framats*. Mosk., 1900, v. VIII.—BURCARR, D. M. *Diss. de inspectione et sectione legali, harumque exemplo speciali*. Lipsæ, 1785.—BURDACH, C. F. *Gerichtsärztliche Arbeiten*. Stuttgart u. Tübingen, 1839.—BURGMEISTER, P. *De antiquissimo illo more, quo veteres reorum innocentium culpam per aquam ferventem ac frigidam probare solebant*. Ulmae, 1680.—BURROWS, G. *Introductory lecture to a course of forensic medicine*. Lond., 1831.—BUSSE, O. *Das Sektions Protokoll*. Berl., 1900.
- CAMPOGRANDE, V. *I diritti sulla propria persona*. Catania, 1896.—CAPURON, J. *La médecine légale relative à l'art des accouchemens*. Par., 1821.—CARDILE-CHOEGLA, G. *La medicina forense*. Palermo, 1872.—CARRANZA, A. *Disputatio de terra humani partus naturalis et legitimi designatione*. Madrid, 1628.—CASPER, J. L. *Atlas zum Handbuch der gerichtlich-medicinischen Leichen-Diagnostik*. Berl., 1857; *Gerichtliche Leichenöffnungen*. Berl., 1852; *Klinische Novellen zur gerichtlichen Medicin*. Berl., 1863; *Practische Handbuch der gerichtlichen Medicin*. Berl., 1860; 1881; *Der Entwurf des neuen Strafgesetzbuchs für die preussischen Staaten vom ärztlichen Standpunkte erläutert*. Berl., 1843; *Beiträge zur medicinischen Statistik und Staatsarzneikunde*. Berl., 1825, 1835; *Denkwürdigkeiten zur medicinischen Statistik und Staatsarzneikunde*. Berl., 1846.—CATECHISM series. *Forensic Medicine*. Edinb., 1896.—DU CAZAL AND CATRIX. *Médecine légale militaire*. Par., 1893.—CELONTI, T. M. *La chirurgia forense esposta in casi pratici, diviso in tre parti, nelle quali si comprendono i trattati delle ferite, delle fratture, e delle lussazioni*. Roma, 1806.—CHAPMAN, H. C. *A manual of medical jurisprudence, insanity, and toxicology*. Phila., 1903.—CHLAUMETON. *Esquisse historique de la médecine légale en France*. Par., 1806.—CHAUSSIER, F. *Observations chirurgico légales sur un point important de la jurisprudence criminelle*. Dijon, 1790; *Recueil de mémoires, consultations et rapports sur divers objets de médecine légale*. Par., 1828; *Médecine légale*. Par., 1838.—\*CHEVALLIER, A. *Traité de toxicologie et de chimie judiciaire*. Par., 1868.—CHEVERS, N. *A manual of medical jurisprudence for Bengal and the Northwestern Provinces*. Calcutta, 1856; *A manual of medical jurisprudence for India, including the outline of a history of crime against the person in India*. Calcutta, 1870.—CHITTENDEN, R. H. *Significance of the absorption and elimination of poisons in medico-legal cases*. N. Y., 1885.—CHITTY, J. *A practical treatise on medical jurisprudence*. Phila., 1835–1836.—CHOULANT, J. L. *Gutachten und Aufsätze im Gebiete Staatsarzneikunde*. Leipz., 1847; *Auswahl von Gutachten medicinisch-forensischen und medicinal-polizeilichen Inhalts*. Dresden, 1853.—CHOULETTE, S. *Observations pratiques de chimie, de pharmacie, et de médecine légale*. Par., 1860.—CHRISTISON, R. *Syllabus of the university course of lectures on medical jurisprudence*. Edinb., 1826; *A treatise on poisons in relation to medical jurisprudence, physiology, and practice of physics*. Edinb., 1829.—CICCONE, A. *Questioni di medicina forense secondo lo spirito del codice delle leggi del regno delle due Sicilie*. Napoli, 1847.—CLARK, H. G. *Medical jurisprudence*. Bost., 1868.—CLÉMENT, E. *Conférences pratiques de médecine légale*. Par., 1880.—CLAUDER, C. E. *Praxis medicolegalis*. Altenburg, 1736.—CLAUSADE, A. *Essai sur la médecine légale considérée comme science*. Montpellier, 1838.—COETSEM, C. A. VAN. *Elementa medicinae forensis*. Gandavi, 1827.—COLEY, H. *Treatise on medical jurisprudence*. New York, 1832.—COMBES, J. M. L. H. *Cours d'hygiène et de médecine légale*. Toulouse, 1841.—COOPER, T. *Tracts on medical jurisprudence*. Phila., 1819.—COR-



- PENS, C. *Moral principles and medical practice. The basis of medical jurisprudence.* New York, Cincin., and Chic., 1897.—COUTAGNE, J. P. H. *Manuel des expertises médicales en matière criminelle à l'usage des magistrats instructeurs et des officiers de police judiciaire.* Lyon, 1887; *Précis de médecine légale.* Lyon and Par., 1896.—CRAIG, J. *On the law of the coroner, and on medical evidence in the preliminary investigation of criminal cases in Scotland.* Edinb., 1855.—DE CRECCHIO, L. *Lezioni di medicina legale secondo i codici del regno d'Italia.* Napoli, 1871.—CROMMELINCK, C. *Annales medico-legales Belgas.* Bruxelles, 1843, 1844.—CSOKOR, J. *Lehrbuch der gerichtlicher Thierheilkunde.* Wien, 1889.—CUESTA Y CKERNER, J. *Medicina legal.* Madrid, 1868.—CUGINI, A. *Profilazione al corso di medicina legale nell'anno scolastico 1889-90 letta nella r. Università di Parma.* Parma, 1890.
- DAMBRE, A. *Traité de médecine légale et de jurisprudence de la médecine.* Bruxelles, 1885.—DANCOURT, J. *Essai sur les moyens que nous offre la médecine légale pour distinguer l'assassinat du suicide dans la cas de plaier.* Par., 1808.—DANGER, F. P. AND FLANDIN, C. *De l'arsenic, suivi d'une instruction propre à servir de guide aux experts dans les cas d'empoisonnement.* Par., 1841.—DANIELS, C. F. *Sammlung medicinischer Gutachten und Zeugnisse.* Leipz., 1776; *Entwurf einer Bibliothek der Staatsarzneikunde oder der gerichtlichen Arzneikunde und medicinischen Polizey, von ihrem Anfange bis auf das Jahr 1784.* Halle, 1784; *Institutionum medicinarum publicarum edendarum adumbratio cum specimen de vulnere letalitate.* Lipsie, 1778.—DARDANO, P. A. *Degli ufficii dell'igiene e della medicina legale.* Genova, 1836.—D'ASSIER, S. P. L. A. *Quelques mots sur la médecine légale et sur la liberté morale.* Par., 1829.—DEAN, A. *Principles of medical jurisprudence.* Albany, 1854; *Manual of medical jurisprudence.* Albany, 1840.—DEASE, W. *Remarks on medical jurisprudence.* Phila., 1819.—DÉGRANGES, E. AND LAFARGUE, E. *Observation de médecine légale.* Bordeaux, 1851.—DEINLIN, J. J. F. *De medico inter sententias medicolegales discrepantes arbitro tertio, germanice.* Obmann. Altorfii, 1751.—\*DENONVILLIERS, NÉLATON, AND TARDIEU. *Questions médico-légales de responsabilité médicale.* Par., 1857.—DERVIEUX, F. *Les expertises médico-légales.* Par., 1901.—DESFOSSÉS, E. *Étude médico-légale sur les blessures par revolver.* Par., 1886.—DESMAZE, C. *Histoire de la médecine légale en France d'après les lois, registres et arrêts criminels.* Par., 1880.—DESSART, J. B., AND THIEBAULD, C. *Traité de médecine légale vétérinaire.* Bruxelles, 1879.—DEVERGIE, M. G. A. *Médecine légale, théorique et pratique.* Par., 1852; *Des experts en justice et de l'expertiste médico-légale.* Par., 1878; *Mémoire sur la combustion humaine spontanée.* Par., 1857.—DEV, K. L. *Medical jurisprudence.* Calcutta, 1876.—DIEHL, J. L. *Anatomischer Atlas der gerichtlichen Praxis zum Gebrauche bei Legaluntersuchungen für Aerzte und Richter.* Heidelberg, 1839-1840.—DIATRIBE *juridico-practica de testimoniis sanitatis.* N. p. 1710.—DISERTACIONES medico-legales. Mex., 1850.—DITTRICH, P. *Lehrbuch der gerichtlichen Medicin für Studierende und Aerzte.* Wien und Leipz., 1897.—\*DORN, A. *Die gerichtliche Arzneiwissenschaft in ihrer Anwendung.* Munich, 1813.—DORNBLÜTH, A. L. *Die medicinal Ordnung im Grossherzogthum Mecklenburg-Schwerin.* Güstow, 1840.—DORNIER, M. *De l'homicide conjugal.* Lyon and Par., 1904.—DOUAY, E. *Le suicide ou la mort volontaire.* Par., 1870.—DRAGENDORFF, G. J. L. *Beiträge zur gerichtlichen Chemie einzelner organischer Gifte.* St. Petersburg, 1872; *Die gerichtlich-chemische Ermittlung von Giften in Nahrungsmitteln, Luftgemischen, Speiseresten, Körpertheilen, etc.* St. Petersburg, 1868; Göttingen, 1888.—DRAPER, F. W. *The medical profession and the commonwealth.* Bost., 1892.—DUBRAC, F. *Traité de jurisprudence médicale et pharmaceutique.* Par., 1893.—\*DUFLOS, A. *Handbuch der angewandten gerichtlich-chemischen Analyse der chemischen Gifte.* Breslau u. Leipz., 1873.—DUMAS, F. M. P. I. *Exposition historique et appréciation des secours empruntés par la médecine légale à l'anatomie et à la physiologie.* Strasbourg, 1838; *Rapports de la médecine légale avec la législation.* Strasbourg, 1840.—DUMONT, J. *Aperçu sommaire de la physiologie médico-légale, ou nécessité de l'intervention médicale dans certains faits déferés aux tribunaux.* Angers, 1860.—DUNCAN, A. *Heads of lectures on judicial medicine.* Edinb., 1801.—DUNGLISON, R. *Syllabus of the lectures on medical jurisprudence and on the treatment of poisoning and suspended animation.* University of Virginia, 1827.—DUPOSCHÉ, E. *Traité de médecine légale militaire.* Par., 1890.—DWIGHT, E. W. *Epitome of medical jurisprudence.* Phila., 1903.—DWIGHT, T. *Identification of the human skeleton.* Bost., 1878.

- B. BEVERLY, M. G. *The trial of J. Reynolds medico-legally considered*. New York, 1870.—\*BENHART, G. *Magazin f. d. technische Heilkunde*. Öffentl. Arzneiwissenschaft und med. Gesetzgebung. Ulm, 1805.—EKAMA, S. *Tracta juridica*. Lugd. Bat., 1844.—\*EISENBERG, J. A. *Untersuchungen über die Veränderungen im Körper der Neugeborenen durch Asphyxie und Luftvergiftungen in anatomischer und gerichtlicher Hinsicht*. Stuttgart, 1853.—EVERT, E. G. *Einige Fälle aus der gerichtlichen Arzneikunde*. Tübingen, 1792.—EWELL, J. J. *A medico-legal treatise on malpractice and medical evidence*. New York, 1860.—EMMER, C. *Lehrbuch der gerichtlichen Medizin*. Leipz., 1900.—ENTRES, J. *Handbuch der gerichtlichen Obduktions-Technik*. München, 1901.—ERICHSEN, J. E. *On surgical evidence in courts of law, with suggestions for its improvement*. Lond., 1878.—ERMEI, F. A. *Rhapsodia supplementa*. Lipsiæ, 1770.—\*VAN ERMINGEN. *Congres de médecine légale*. Bruxelles, 1897.—\*ESCHENBACH, C. E. *Medicus legalis brevissumus*. Rostock, 1746.—ESTLANDER, J. A. *Den medicolegala indelningarna af kroppstadiet, med särskildt afseende å det i svenska och finska läroförekommande uttrycket "lifsfarlig sjukdom"*. Helsingfors, 1872.—EYSEARTH, C. S. *De recto rationis atque experientiae usu in praxi clinico-forensi*. Lipsiæ, 1793.—FABRE, A. F. H. *Bibliothèque du médecin-praticien*, v. XV. Par., 1851.—\*FABRICIUS, P. C. *Sammlung verschiedener medic. responsorum und Sectionsberichte*. Halle, 1772.—\*FAHNER, J. C. *Beiträge zur pract. und gerichtl. Arzneikunde*. Stendal, 1799; *System einer vollständigen medicinischen Polizey*. Berlin, 1792.—FALCK, F. A. *Lehrbuch der praktischen toxiologie*. Stuttgart, 1880; *Art in Handbuch der gerichtl. med.*. Tübingen, 1881.—\*FALK. *Juristische Encyclopædie*. Kiel, 1821.—FANTONETTI, G. B. *Delle malattie in rapporto alla medicina legale*. Ferrara, 1863.—FANZAGO, F. L. *Discorso inaugurale del P. P. di patologia*. Trattati teorico-practici e medicina legale. Padova, 1807.—FARR, S. *Elements of medical jurisprudence*. Lond., 1788.—FASLIUS, J. F. *Elementa medicina forensis*. Jenæ, 1767.—FAURE, R. *Exposition historique et appréciation des secours empruntés par la médecine légale à la médecine proprement dite*. Strasbourg, 1838.—FERDINANDUS. *Contra jurisprudentia vituperatores oratio, qua manifeste declarat leges plurimum medicina philosophicæ artibus antefereudas esse*. Bononiæ, 1566.—FERNANDEZ DEL VALLE, J. *Cirurgia forense general y particular*. Madrid, 1796.—FERRARESE, L. *Programma di psicologia medico-forense*. Naples, 1834; *Quistioni di psicologia medico-forense*. Napoli, 1837.—FERKER Y GARCÉS R. *Tratado de medicina legal*. Barcelona, 1867.—FERRI, E. *Sul nuovo codice penale*. Napoli, 1889.—VON FEUERBACH, P. J. A. *Translation Narratives of remarkable criminal trials*. Lond., 1846.—FIDELIS, F. *De relationibus medicorum*. Lipsiæ, 1674.—FIELD, G. W. *Medico-legal guide for doctors and lawyers*. New York and Albany, 1887.—FIELTZ, F. G. H. *Archiv der gerichtlichen Arzneiwissenschaft*. Luckau, 1812.—FILIPPI, A. *Principii di medicina legale per gli studenti legge ed i giurisperiti*. Firenze, 1889; *Essegesi medico-legale sul methodus testificandi di Giovan. Battista Codronchi*. Firenze, 1883; With SEVERI, A. and MONTALTI, A. *Manuale di medicina legale conforme al nuovo codice penale per medici e giuristi*. Milano, 1889.—FILOMUSI-GUELFI, G. *L'indole delle ricerche medico-legali*. Pavia, 1887; also edits the *Giornale de medicina legale*. Lanciano.—FINCK, J. B. D. *De officio medici circa torturam*. Erfordie, 1760.—FIRGAU, F. *Gifte und stark wirkende Arzneimitteln in gerichtlicher, hygienischer und gewerblicher Beziehung*. Berl., 1901.—FISCHER, J. A. *Responsa practica et forensia selecta ab anno MDCCII*. Francofurti et Lipsiæ, 1719.—FLANDIN, C. *Traité des poisons, ou toxicologie appliquée à la médecine légale, à la physiologie et à la thérapeutique*. Par., 1847.—\*FLEISCHMANN. *Médecine légale*.—FLEISCHMANN, G. *Anleitung zur forensischen und polizeilichen Untersuchung der Menschen und Thierleichen für Vorlesungen entworfen*. Erlangen, 1811.—FLEMMING, A. *Bloodstains in criminal trials*. Pittsburg, 1861.—FLEURET, L. P. *Médecine légale pratique*. Annecy, 1842.—FLITTNER, C. G. *Schutz und Rettung in Todesgefahr*. Berl., 1825.—FODERÉ, F. E. *Les lois éclairées par les sciences physiques*. Par., 1799; *Traité de médecine légale et d'hygiène publique*. Par., 1813.—FORMEY, J. L. *Medizinische Eph. c. meriden von Berlin*. Berl., 1800.—FORNASINI, L. *Studi di medicina legale*. Brescia, 1877.—FORSYTH, J. S. *A synopsis of modern medical jurisprudence*. Lond., 1829.—\*FORTUNATUS, F. *De relationibus medicorum libri quatuor, in quibus ea omnia quæ in forensibus ac publicis causis medici referre solent, plenissime tractantur*. Palerme, 1802.—FRANK, J. P. *System einer vollständigen medicinischen Polizey*. Frank-



- enthal, 1791.—\*FRENZEL, J. S. T. *Gerichtl.-polizeiliche Arzneiwissenschaft*. Leipz., 1791.—FRESCHI, F. *Manuale teorico-pratico di medicina legale*. Milano, 1846.—FRIEDBERG, H. *Gerichtsärztliche Gutachten*. Braunschweig, 1875; *Gerichtsärztliche Praxis*. Wien u. Leipz., 1881.—FRIEDREICH, J. B. *Handbuch der gerichtl.-ärztlichen Praxis mit Einschluss der gerichtlichen Veterinärkunde*. Regensburg, 1844; *Memoranda der gerichtlichen Anatomie, Physiologie und Pathologie*. Wurtzburg, 1857; *System der gerichtlichen Psychologie*. Regensburg, 1842.—FUENTES, M. A. *Manual práctico de medicina legal*. Lima, 1869.—\*FURST, Ueber den Tod durch giftige Gase. Berl., N. d.—\*FÜRSTENAU, J. H. *Medicina forensis contracta*. Rinteln, 1752.
- \*GADELIUS, E. *Manual of medical jurisprudence*. Stockholm, 1804.—GADOW, G. *Rechts-Vademecum für den praktischen Arzt*. Breslau, 1892.—GALLARD, T. *De l'avortement au point de vue médico-légal*. Par., 1878; *Notes et observations de médecine légale et d'hygiène*. Par., 1875; *Deux faits de médecine légale*. Par., 1877.—GALLIER, A. *Médecine légale vétérinaire*. Par., 1895.—GALTIER, C. P. *Traité de toxicologie médicale, chimique et légale*. Par., 1855.—GALTIER, V. *Traité de législation commerciale et de médecine légale vétérinaires*. Par., 1897.—GALVAO, J. P. F. *Tratado elementar de medicina legal*. Porto, 1855.—GANDOLFI, G. *Fondamenti di medicina forense analitica ad uso del medico e del legale*. Bologna and Modena, 1854.—GARIBALDI, G. A. G. *Manuale di medicina legale ad uso degli studenti di legge, con note sui termini tecnici*. Genova, 1876.—GATTA, R. *Compendio di medicina legale*. Napoli, 1892.—GERICHTLICH-medizinische Fälle und Abhandlungen. Berl. and Newied, 1888.—GEREGTELIJKE geneeskunde, Emoi, 1883.—GEREGTELIJK-geneeskundig Vademecum. Leeuwarden, 1847.—GERLACH, A. C. *Handbuch der gerichtlichen Thierheilkunde*. Berl., 1872.—\*GIANELLI, G. L. *Trattato di medicina publica diviso in re parti*. Padova, 1836; *L' uomo ed i codici nel nuovo regno italico*. Milano, 1860.—GIANQUINTO, J. DE G. *Programme du traité médico-légal sur le secret*. Florence, 1867.—GIPFEN, G. H. *Student's manual of medical jurisprudence and public health*. Edinb., 1901.—GILMAN, C. R. *Syllabus of medical jurisprudence*. N. p., N. d. *The relations of the medical to the legal profession*. New York, 1856.—GINÉ Y PARTAGAS, J. *El código penal y la frenopatología*. Barcelona, 1888.—GIUSTINI, A. *Quaestione medico-legale*. Viterbo, 1840.—GLAISTER, J. *Medical jurisprudence, toxicology, and public health*. (The best textbook on legal medicine in the English language.) Edinb., 1902.—GLASER, J. E. *Diss.-commentationem medicam in aedilitium edictum*. Halæ, 1738.—\*GOELICKE, A. O. *Medicina forensis demonstrativa methodo tradita cui praemisse est introductis in historiam litterariam scriptorum medicinarum forensium illustration*. Francfort sur l' Oder, 1723, 1725.—GOLEBINSKI, E. *Atlas and epitome of diseases caused by accidents*. Phila., 1900.—\*GOODWIN, E. *Dissert. de morbo morte quae submersorum investigandis*. Edinb., 1786.—\*GOSSE, H. *Des taches en médecine légale*.—GOTSCHALK, R. *Grundriss der gerichtlichen Medicin für Aerzte und Juristen*. Leipz., 1903; *Krimineller Abort und Kindesmord*. *Klin. Jahrb.*, v. XI, 1903.—GRIBBLE, J. D. B. AND HEHIR, P. *Outlines of medical jurisprudence for India*. Madras, 1898.—GRINNELL, C. E. *Points of law for lawyers and general readers suggested by Guiteau's case*. Rost., 1881.—GROSZ, L. *A manual of medical jurisprudence for Hungary*. Budapest, 1873.—AF GRUBBENS, N. W. *Observations in pathological anatomy at medico-legal autopsies*. Malmö, 1849.—GRÜNER, C. G. *Insunt pandecta medicae*. Jena, 1800.—GUDER, P. M. F. *Compendium der gerichtlichen Medicin*. Leipz., 1887; *Gerichtliche Medicin*. Leipz., 1900.—GUEORGUEFF, S. G. *Quelques notions sur l'histoire de la médecine légale en Russie*. Geneve, 1890.—GUERNSEY, R. S. *Medico-legal sciences*. New York, 1882.—GUERRIER, L. AND ROTUREAU, L. *Manuel pratique de jurisprudence médicale*. Par., 1890.—GUETERBOCK, C. E. *H. de Bracton, quo tempore et qua ratione librum de jure anglicano composueret*. Regimonti, 1860.—GUILLERY, *Vademecum du médecin légiste*. Bruxelles, 1892.—GÜNTNER, F. X. *Handbuch der gerichtlichen Medicin*. Regensburg, 1851.—\*GURNBILL, R. J. *The morals of suicide*. Lond., n. d.—GUY, W. A. *The factors of the unsound mind, with special reference to the plea of insanity in criminal cases and the amendment of the law*. Lond., 1881; *Principles of forensic medicine*. Lond., 1844; Bost., 1868; With FERRIER, D. *Principles of forensic medicine*. Lond., 1874.—GVOZDEFF, I. M. *Systematic list of medico-legal data, chiefly as practically applied to public needs*. Kazan, 1892; *Medico-legal data in the hands of the jurists*. Kazan, 1889.

HAASE, J. G. *De praecipuis momentis quorum ratio a medico forensi est*



habenda officio suo honeste functuro exposuit. Lipsiæ, 1798.—HAINES, W. S. *Textbook of legal medicine and toxicology*. Phila., 1904.—HAGENBACH, F. *Schema des Criminal-Rechtes*. Basel, 1870.—HALL, A. G. *Institutes of medical jurisprudence*. Bost., 1860.—HALL, J. C. *Medical evidence in railway accidents*. Lond., 1868.—VON HALLER, A. *Vorlesungen über die gerichtlichen Arznei- und Wundheilkunde*. Berne, 1784.—HAMER, J. H. *Diss. sistens medicinam renunciatorem sine crisin, qua juris consultis et iudicibus forensibus respondere tenetur medicus*. Erfordiae, 1692.—HAMILTON, A. McL. *A manual of medical jurisprudence with special reference to diseases and injuries of the nervous system*. New York, 1885; With GODKIN, L. *A system of legal medicine*. New York, 1894.—HANSBACH, J. *Handbuch der im Königreich Sachsen gelten den medicinal Polizeigesetze*. Leipzig, 1837.—HANSEN, S. *Handbook of medical jurisprudence*. Kjøbenhavn, 1898.—HASENSTEIN, J. G. *Der medicinische Richter*. Onolzbach, 1759.—HASLAM, J. *Medical jurisprudence, as it relates to insanity, according to law of England*. Cooper's tracts on medical jurisprudence. Phila., 1819.—HASS, S. *Ueber die Instanzenzug gerichtl.-medizinischer Berichte*. Biel, 1890.—VAN HASSELT, A. W. M. *Handbuch der Giftelehre für Chemiker, Aerzte, Apotheker, und Richtspersonen*. Braunschweig, 1862.—HAUSKA, F. *Compendium der gerichtlichen Arzneikunde*. Wien, 1869.—HEBENSTREIT, J. E. *Anthropologia forensis*. 1753; *De officio medici forensis spec. II*. Lipsiæ, 1748.—\*HEINROTH, J. C. A. *System der gerichtlich-psychischen Medizin*. Leipz., 1825.—\*HEINROTH, J. C. A., AND HAYNEL, C. A. *De facinoræ aperto ad medicorum iudicium non deferendo*. Lipsiæ, 1830.—HILST, L. C. *Compendium medicina renunciatoria*. Frankfurt und Leipzig, 1713.—HEKTOEN, L. *Two interesting medico-legal cases*. Chicago, 1891.—HEMMING, W. D. *Forensic medicine and toxicology*. New York, 1889; *Aids to forensic medicine and toxicology*. Lond., 1877.—\*HENCKEL, *Gerichtliche Medizin*. N. p. N. 3.—\*HENKE, A. *Medicina legalis. Abhandlungen aus dem Gebiete der gerichtlichen Medizin*. Leipz., 1823; *Lehrbuch der gerichtlichen Medizin*. Stuttgart, 1832.—HEROLD, J. *A manual of legal medicine*. Phila., 1902.—HERVOUT, G. C. F. *Tuberculisme et tuberculose pulmonaire au point de vue médico-légal*. Nantes, 1903.—HEUR, F. D. *Diss. de sugillatione quatenus infanticidii indicium*. Schlegel's *Collectio opusculorum*. Lipsiæ, 1785.—HIDALGO Y CARPIO, L. *Introducción al estudio de la medicina legal mexicana*. México, 1869; With RUIZ Y SANDOVAL, G. *Compendio de medicina legal*. México, 1877.—HILST, P. F. *Considerations sur les prolégomènes de la médecine légale*. Par., 1823.—HINTZE, W. *Mitteilungen aus dem Gebiete der gerichtlichen Medizin zur Beurteilung der Kurfürstlichen und des Geheimmittelwesens*. Berl., 1897.—HOFFBAUER, J. C. *Die Physiologie in ihrer Hauptanwendung auf die Rechtspflege nach den allgemeinen Gesichtspunkten der Gesetzgebung*. Halle, 1823.—\*HOFFMANN, FR. *Dissert. continens observationes medico-forenses selectas*. Halæ, 1728.—VON HOFMANN, E. *Atlas-manuel de médecine légale*. Par., 1899; *Nouveaux éléments de médecine légale*. Par., 1881; *Lehrbuch der gerichtlichen Medizin*. Wien und Leipz., 1891; *Atlas of legal medicine*, translated by KELLY, A. O. J.; With KOLISKO, A., 1902. (This is the best book on legal medicine published.)—HOFMANN, J. *Aus dem Gerichtssaale*. Erlangen, 1864; *Traité de médecine légale*. Par., 1898; *Die gerichtsarztliche Sprache*. München, 1860.—\*Der preussische Gerichtsarzt.—HOTCHKISS, C. *Criminalité et médecine judiciaire dans l'Inde anglaise*. Lyon, 1893.—HOWARD, H. *Outlines of medical jurisprudence*. Charlottesville, 1845; *A synopsis of medical jurisprudence*. Charlottesville, 1851.—HUARD, *Considérations médico-légales sur deux articles du titre second du code pénal*. Par., 1819.—\*HÜHNFELD, F. L. *Die Chemie der Rechtspflege oder Lehrbuch der polizeil-gerichtlichen Chemie*. Berl., 1832.—HUSBAND, J. A. *Students' hand-book of forensic medicine and public health*. Edinb. and Lond., 1889; *Students' handbook of forensic medicine and medical police*. Edinb., 1874; *Forensic medicine, toxicology, and public health*, edited by Buchanan and Hope. Edinb., 1904.—\*HUTCHINSON, W. *Infanticide*. Lond., 1820.—INSTRUCTIONS pour les médecins de campagne et chirurgiens. Par., 1789.—INSTRUCTION für die öffentlich angestellten Aerzte und Wundärzte in den k. k. Österreichischen Staaten, wie sie sich bei gerichtlich Leichenschauen zu benehmen haben. Wien, 1814.—INSTRUKTION für das Verfahren der Aerzte im Königreiche Bayern bei den gerichtlichen Untersuchungen menschlicher Leichen. München, 1886.—ISTRUZIONI per uso de' medici e chirurghi incaricati di pubblico servizio negli imperiali e reali stati austriaci riguardanti le visite giudiziali dei cadaveri. Venezia, 1819.



- JÄGER, J. *De vinculo juris inter medicum et agrotantem*. Amstelodami, 1850.  
 —JAUZE, F. *Médecine légale hippiatrice abrégé de la pratique vétérinaire*. Par., 1844.—JOHN, J. D. *Die Polizei und gerichtliche Arzneiwissenschaft in der k. k. Erbländern*. Prag, 1798.—JOHNSON, C. *An essay on the signs of murder in new-born children*. Montreal M. J., 1893-1894, v. XXII, p. 169.—JOHNSTON, W., AND VILLENEUVE, G. *Revue statistique des enquêtes tenues par la cour du coroner du district de Montréal pendant l'année 1893*. Montreal, 1893.—JUGLER, J. H. *Repertorium für das neueste aus der Staatsarznei-Wissenschaft*. Braunschweig, 1801; *Medizinische Merkwürdigkeiten für Criminalrichter, Aerzte und Prediger*. Leipz., 1803.  
 KALISCH, M. *Medicinisch-gerichtliche Gutachten der königlichen preussischen wissenschaftlichen Deputation für das medicinälwesen aus den Jahren 1840-1850*. Leipz., 1859.—KANNEGIESER, G. H. *Institutiones medicinae legalis in usum auditorum concinnatae cum praefatione*. Kilise Holsatorium, 1777.—KATAYANA KUNIKI. *Medical jurisprudence*. Tokio, 1824.—KATONE, S. *Horitsu gakusha no danso igaku ni zukete*. Tokyo, 1894.—\*KAUSCH, J. J. *Fragmente der Militärstaatsarzneikunde*. Leipz., 1806; *Memoralbilien der Heilkunde-Staatsarzneiwissenschaft und Thierheilkunst*. Züllichau, 1819; *Über die neuen Theorien des Criminalrechts und der gerichtl. Medizin mit Vorschlägen zur Verbesserung bei der Disciplinen*. Züllichau, 1810.—KECK, J. E. *Abhandlungen und Beobachtungen aus der praktischen und gerichtlichen Arzneiwissenschaft*. Berl., 1787.—KIEN, C. J. *De jure medicorum*. Traj. ad Rhenum, 1757.—KEYSOR, W. W. *A medico-legal manual*. Omaha, 1901.—KIERNANDER, J. *Utkast til medicinal Lagfarenheten Domare til Uplysning Läkare til Hjelpreda och Barnmorskor til Undervisning i Amnen, som röra Manniskokroppen*. Stockholm, 1776.—KIMMICH, D. J. *Sur l'importance de la médecine dans un état policé, ou sur la médecine légale*. Strasbourg, 1803.—\*KIPPENBERGER, C. *Aufgaben einer wissensch.-gerichtl. Chemie*. Berl.—v. KLEIN, C. C. *Beiträge zu der gerichtlichen Arzneiwissenschaft*. Tübingen, 1825.—KLEIN, J. *Elemente der forensisch chemischen Ausmittlung der Gifte*. Hamburg, 1902.—VON KLEIN, C. H. *A manual of medical jurisprudence*. Hamilton, Ohio, 1882.—\*KLEIN, C. F. *Annalen, die Gesetzgebung und Rechtsgelahrtheit in den preuss. Staaten betreffend*. Berl., 1804.—KLOSE, W. F. W. *System der gerichtlichen Physik*. Breslau, 1814; *\*Beiträge zur gerichtlichen Arzneikunde*. Breslau u. Leipz., 1811.—KLOSE, C. L. *Beiträge zur Klinik und Staatsarzneiwissenschaft*. Leipz., 1823.—KLUG. *Auswahl medicinisch gerichtlicher Gutachten der königl. wissenschaftlichen Deputation für das medicinal wesen*. Berl., 1828.—KNAGES, S. AND KNAPE, C. *Kritische Annalen der Staatsarzneikunde*. Berl., 1804; With HECKER, A. F. *Kritische Annalen der Staatsarzneikunde*. Berl., 1806.—\*KNEBEL, J. G. *Grundlage zu einem vollständigen Handbuch der Literatur für die gesammte Staatsarzneikunde bis zu Ende des achtzehnten Jahrhunderts*. Gortitz, 1806; *\*Grundriss der polizeilich gerichtlichen Entbindungskunde*. Breslau, 1801-1803.—KOBERT, R. *Compendium der praktischen Toxikologie*. Stuttgart, 1903; *Lehrbuch der Intoxikationen*. Stuttgart, 1902. (These two works contain much valuable information.)—KOCKEL, R. *Die gegenwärtige Bedeutung der gerichtlichen Medizin*. Leipz., 1898.—KOCHER, A. *De la criminalité chez les Arabes au point de vue de la pratique medico-judiciaire en Algérie*. Lyon, 1883.—KOMORAU, J. *Visa reperta zum praktischen Gebrauche für Aerzte und Wundärzte*. Wien, 1869.—KORNFELD, H. *Handbuch der gerichtlichen Medizin*. Stuttgart, 1884.—KOSOROTOFF, D. P. *Osnovniya pravila sostavleniya sudebno-meditsinskikh aktov o vskritiyakh myortvikh tel*. St. Petersburg, 1900.—KOST, J. *A textbook on medical jurisprudence*. Cincinnati, 1885.—KRAHMER, L. *Handbuch der gerichtlichen Medizin*. Halle, 1851.—KRAMER, G. F. *Repertorium der älteren und neuesten Gesetze über die medicinal Verfassung in Königreiche Bayern*. Augsburg, 1832.—KRAUS, L. G., AND PICHLER, W. *Encyclopädisches Wörterbuch der Staatsarzneikunde*. Erlangen, 1877.—VON KROMBHOlz, J. V. *Gerichtlich medicinische Untersuchungen nebst Gutachten*. Prag, 1841.—KRUGELSTEIN, F. C. C. *Promptuarium medicinae forensis*. Erfurt and Gotha, 1822.—KUBIN, A. *Diss. sistens medicinae veterinaria forensis primas lineas*. Giessen, 1810.—KUHN, J. G. *Sammlung medicinischer Gutachten*. Breslau, 1796.—\*KUHN, O. B. *Practische Anweisung die in gerichtlichen Fällen vorkommenden Untersuchungen anzustellen*. Leipz., 1829.—KURZES *Repertorium der gerichtlichen Medizin, als Vademecum für die Praxis und für die Prüfungen*. Leipz. und Wien, 1892.—KÜHNHOLTZ, H. *Coup d'œil sur l'ensemble systématique de la médecine judiciaire*. Montpellier,



1834.—KUTTER, C. *Kurzgefasste gerichtlich-medicinische Darstellung der häufigsten zur Aburtheilung kommenden Criminalfälle*. Stuttgart, 1860.

LACASSAGNE, A. *Précis de médecine judiciaire*. Par., 1886; *Le vade-mécum du médecin expert*. Par., 1900; *Feuilles d'examen médico-légal et d'autopsie*. Par. and Lyon, 1892.—LAND, A. L. *Handleiding bij het peregte lijk geneeskundig onderzoeken van lijken*. Leeuwarden, 1838.—LONDON, L. *Contributions à l'étude médico-légale des méningites*. Par., 1903.—LARCHER, J. F. *Études physiologiques et médicales sur quelques lois de l'organisme, avec applications à la médecine légale*. Par., 1868.—LARROQUE, F. T. *Quelques propositions générales de médecine, de chirurgie, de médecine légale et d'accouchement*. Par., 1821.—LAURENT, B. *Prodromus einer polizeilich-gerichtlichen Thierarzneikunde*. München, 1812.—LAURA, S. *Trattato di medicina legale*. Torino, 1874.—LAZZARETTI, G. *Corso teorico-pratico di medicina-legale*. Padova, 1880.—LEBLANC, U. *Recherches relatives à la détermination de l'âge des lésions des plèvres et des poumons du cheval—au point de vue médico-légal*. Par., 1841.—LECIEUX, A., et al. *Médecine légale ou considérations sur l'infanticide*. Par., 1819.—LE CLERC, N. *Essai médico-légal sur l'empoisonnement et sur les moyens que l'on doit employer pour les constater*. Par., 1803.—LEE, J. G. *Handbook for coroners*. Phila., 1881.—LEHNE, P. *Aide-mémoire de médecine légale pour la préparation du quatrième examen*. Par., 1903.—LEGLUDIC, H. *Notes et observations de médecine légale*. Par., 1896.—LEGRAND DU SAULLE, H., BERRYER, G., AND POUCHET, G. *Traité de médecine légale, de jurisprudence et de toxicologie*. Par., 1886.—LEPOUTRE, A., AND PIERRE, L. *Traité de jurisprudence médicale*. Lille, 1841.—LESSER, A. *Atlas der gerichtlichen Medicin*. Breslau, 1890. (This is the best illustrated medicolegal atlas in existence.) *Stereoskopischer gerichtsarztlicher Atlas*. Bresl., 1903.—LESUEUR, D. *Recherches médico-légales pouvant servir à déterminer même longtemps après la mort s'il y a eu empoisonnement*.—LEVILLAIN, T. T. *Considérations médico-légales sur les visites et rapports en justice*. Par., 1814.—LIBERATORE, G. *Definizione medicolegali penali*. Napoli, 1841.—LINDES, *Practische Anleitung zu den gerichtlichen gerichtlich-chemischen und sanitäts polizeilichen Untersuchungen*. Berl., 1849.—LION, A. *Taschenbuch der gerichtlichen Medicin nach dem neuesten Standpunkt der Wissenschaft und der Gesetzgebungen Deutschlands*. Erlangen, 1861.—LOBSTEIN, J. F. *Plan raisonné d'un cours de médecine légale*. Strasbourg, 1814.—LOCARD, E. *Le xvi<sup>e</sup> siècle médico-judiciaire*. Lyon and Par., 1902.—LODER, J. C. *Methodum ad medicinam forensensem spectantium*. Jenæ, 1783.—LOEFFLER, F. B. *Das preussische Physikalexamen*. Berl., 1878.—LOMBROSO, C. *Lezioni di medicina legale*. Roma, Torino, and Firenze, 1900; *Troppo foresto*. Torino, 1888.—LORENZ, L. *Criminalité et médecine judiciaire en Cochinchine*. Lyon, 1887.—LOUIS, M. *Lettres sur la certitude des signes de la mort; avec des observations et des expériences sur les noyés*. Par., 1752.—LÖW, J. F. *Theatrum medico-juridicum Norimbergæ*, 1725.—LÖWENHARDT, S. E. *Untersuchungen im Gebiete der gerichtlichen Arzneiwissenschaft für Aerzte und Criminalisten*. Berl., 1848.—LUDWIG, E. *Medicinische Chemie in Anwendung auf gerichtliche Untersuchungen*.—LUDWIG, C. F. *Adversaria ad medicam publicam*. N. p. 1814.—LUDWIG, G. G. *Instructiones medicina forensis praelectionibus academicis accommodatae*. Lipsia, 1788.—LUFF, A. P. *A text-book of forensic medicine and toxicology*. London and New York, 1895.—LUTAUD, A. J. *Manuel de médecine légale et de jurisprudence médicale*. Par., 1886.—LYON, I. B. *A text-book of medical jurisprudence for India*. Calcutta, 1889; Revised edition by L. A. Waddell, 1904.

MACKELROY, R. G. *Estudio médico*. México, 1882.—MACKENZIE, S. C. *Medico-legal experiences in Calcutta*. Edinb., 1891.—MADIA, E. *Compendio di medicina legale*. Napoli, 1902; *Fogli di esame medico-legale e di autopsia*. Napoli, 1901; *Dizionario di medicina legale*. Napoli, 1902.—MAGGIORANI, C. *Proleg. allo stud. di med. politico-legale*. Roma, 1854; *Rettificazione di un errore di storia intorno i primordi della medicina legale*. Roma, 1863.—MAHIER, E. *Les questions médico-légales de Paul Zacchias*. Par., 1872.—MAHON, P. A. O. *Médecine légale*. Par., 1811.—MAILLOT, F. C., AND PUEL, J. A. A. *Aide-mémoire médico-légale de l'officier de santé de l'armée de terre*. Par., 1842.—MAIR, J. *Handbibliothek der praktischen Chirurgie in Monographien nach den einzelnen klinischen Abtheilungen derselben, mit besunder Berücksichtigung der chirurgischen Anatomie und gerichtlichen Medizin bearbeitet*. Ansbach and München, 1855-61; *Juristisch-medizinischer Commentar der neuen kgl. hayerischen, kgl. preussischen und kais. kgl. österreich. Strafgesetzgebung*. Augsburg, 1862; *Gerichtlich-medicinische Casu-*



- istik der Körperverletzungen und Tödtungen durch mechanisch-physische Gewalt. Ingolstadt, 1881. Handbuch des ärztlichen Dienstes bei den Gerichten und Verwaltungsbehörden. Würzburg, 1869; 1879; Die medicinischen Fachbeziehungen der neuen k. bayerischen Straf Gesetzgebung über Verbrechen und Vergehen. Kempten, 1857.—MALE, G. E. Elements of juridical or forensic medicine. Lond., 1818; An epitome of juridical or forensic medicine. Lond., 1816; Exposition historique et appréciation des secours empruntés par la médecine légale à la physique et à la chimie. Strasbourg, 1838; Des attributions particulières de la médecine légale comme science et comme art. Strasb., 1840.—MANDL. Recherches medico-legales sur le sang. Thèse de Paris, 1842.—MANN, J. D. Forensic medicine and toxicology. Phila., 1893; Lond., 1898.—MARC. Manuel d'autopsie cadaverique médico-légale. Par., 1808.—MARCHANT, G. Esquisse historique sur la médecine légale. Toulouse, 1856.—MARTINI, LORENZO. Introduzione alla medicina legale. Turin, 1825; Manuale di medicina legale. Milano, 1829.—MASCHKA, J. Handbuch der gerichtlichen Medicin. Tübingen, 1881-82; Sammlung Gerichtsarztlicher Gutachten der Prager medicinischen Facultät. Prag, 1853.—\*MASIUS, G. H. System der gerichtlichen Arzneiwissenschaft. Rostock, 1810; Lehrbuch der gerichtlichen Arzneikunde. Altona, 1812; Handbuch der gerichtlichen Arzneiwissenschaft. Stendal, 1821-1832; Erörterungen aus dem Civil und Criminal Rechte vom Standpunkte der gerichtlichen Medicin. Rostock, 1821.—MASSON, C. Essai sur l'histoire et le développement de la médecine légale. Lyon, 1884.—MATA, P. Tratado de medicina y cirugía legal teórico y práctico. Madrid, 1875.—MATTHYSENS, F. J. Précis élémentaire de médecine légale. Anvers, 1838.—MAZZINI, G. Giurisprudenza veterinaria. Milano, 1903. (A work of 530 pages.) MAZZIONI, G. B. Sulla origine e progressi della medicina legale. Firenze, 1828.—MCCALLIN, W. Introduction to medical jurisprudence. Lond., 1901.—MCCLELLAND, A. Civil malpractice: A treatise on surgical jurisprudence. New York, 1877.—\*MCGRATH, J. J. Jurisprudence. Textbook of legal medicine and toxicology.—MCLEOD, K. Medicolegal experience in the Bengal Presidency. Calcutta, 1875.—MECKEL, A. Einige Gegenstände der gerichtlichen Medicin. Halle, 1820; Lehrbuch der gerichtlichen Medicin. Halle, 1821.—MEDICAL jurisprudence or code of ethics and institutions adopted by the profession of physicians and surgeons. P. V., 1174.—MEDICINA legale. MS. N. p., 1820.—MEDICO-LEGAL JOURNAL. N. Y. Established in 1883.—MEDICO-LEGAL SOCIETY of New York. Papers for its organization. New York. N. d.; Prize essays, New York, 1890.—MEISTER, C. F. Urtheile und Gutachten in peinlichen und andern Straffällen. Frankf. a. d. Oder, 1808.—MELSHEIMER, J. Ueber die Lehre von Affect in gerichtlich-medicinischer Beziehung. Giessen, 1862.—MENDE, L. J. Editor of the Beobacht u. Bemerk a. d. Geburtshülfe u. gerichtlichen Medicin. Göttingen, 1824; Ausführliches Handbuch der gerichtlichen Medicin. Leipz., 1819.—METZGER, J. D. Gerichtlich-medicinische Beobachtungen. Königsberg, 1786; Systema medicina forensis succinctum. Stendaliae, 1794; Kurz-gefasstes System der gerichtlichen Arzneiwissenschaft. Königsberg und Leipz., 1820; With ELSNER, C. F. Medicinisch-gerichtliche Bibliothek. Königsberg, 1785.—MINOVICI, M. Étude comparative des réformes à apporter à l'enseignement et exercice de la médecine légale en Roumanie. Par., 1889.—MOLL, A. Leerboek der geregtelijke geneeskunde voor geneesken regtskundigen. Arnhem, 1825-1826.—MONACO, G. Manuale di medicina legale. Napoli, 1887.—DE MONTGARNY, H. Essai de toxicologie et specialment avec la jurisprudence médicale. Par., 1878.—DE MONTMAHOUX, E. S. Manuel médico-légale des poisons. Par., 1824.—MORACHE, G. A. La médecine légale. Par., 1890.—MOST, G. F. Ausführliche Encyclopädie der gesammten Staatsarzneikunde. Leipz., 1838.—MOUZIN LIZYS, R. Étude médico-légale sur la responsabilité professionnelle des médecins, chirurgiens, accoucheurs. Par., 1899.—MOYE, G. C. M. Les expertises médicales devant les tribunaux civils. Bordeaux, 1898.—MUCH, H. Über Todesursachen bei Neugeborenen mit besonderer Berücksichtigung ihrer forensischen Bedeutung. Würz., 1903.—\*MULERT. Schemata und Notizen für gerichtliche Obductionen. Waren.—MÜLLER, F. Compendium der Staatsarzneikunde. München, 1855.—MÜLLER, J. V. Entwurf der gerichtlichen Arzneiwissenschaft. Frankf. a. M., 1796.—MURCELL, W. Aids to forensic medicine and toxicology. Lond., 1903.—NAQUET, A. Précis de chimie légale. Par., 1873.—NARDELLI, R. Tre perizie medico-legali. 1884.—NASSALL, L. L. De sectione legali. Wirceburgi, 1798.—NAVARRO Y ORTIZ, E. Elementos de medicina legal, militar y naval. Madrid,

1804.—\*NEISSER, A. *Stereoscopischer medicinischer Atlas gerichtliche Medicin*. Leipz., 1808.—\*NEUMANN, A. C. *Handbuch der gerichtlichen Anatomie für Rechtsgelehrte, Polizeibeamte und Studierende*. Berl., 1852.—NEUMANN, M. *Formis de renuntiationibus*. Leodii, 1826.—VON NEY, F. *Systematisches Handbuch der gerichtsarzneilichen Wissenschaft*. Wien, 1845; *Die gerichtliche Arzneikunde in ihrem Verhältnisse zur Rechtspflege*. Wien, 1847.—NICOLAI, A. H. *Die Medicinal- und Veterinair-Polizei*. Berl., 1838; *Handbuch der gerichtlichen Medicin*. Berl., 1814.—NIEMANN, J. F. *Handbuch der Staats-Arzneiwissenschaft und staatsärztlichen Veterinärkunde*. Leipz., 1813; *Taschenbuch der Civil-Medicinal-Polizei*. Leipz., 1828; *Taschenbuch der gerichtlichen Arzneiwissenschaft*. Leipz., 1837.—NOEL, J. *Médecine légale*. Strasbourg, N. d.—NOGENT, St. L. *Cow d'assises de la Seine*. Par., 1855.—VON NUSSBAUM, J. N. *Einfluss der Antiseptik auf die gerichtliche Medicin*. München, 1880.

OBOLONSKI, N. A. *Aid in the medico-legal examination of the corpse and a search of material proof*. St. Petersburg, 1894.—\*OECHY, J. *Instruction pour l'examen médico-légal*. Prag., 1802.—OETINGER, W. L. *De causa miraculorum in medico foro disputatione*. Tübingæ, 1723.—OGSTON, F. *Lectures on medical jurisprudence*. Phila., 1878.—OLAECHÉA, M. A. *Cuestiones prácticas de higiene y medicina legal*. Barcelona, 1893.—OPPENHEIMER, A. *Juristisches Handbuch für praktische Aerzte*. Würzburg, 1896.—ORDONNAUX, J. *The jurisprudence of medicine*. Phila., 1869.—ORFILA, P. M. J. B. *Rettungsverfahren bei Vergiftungen und im Scheintode*. Berl., 1819; *Mémoires sur plusieurs questions médico-légales*. Par., 1839; *Traité de médecine légale*. (Valuable references.) Par., 1847; *Leçons de médecine légale*. Par., 1823-1828; *Traité des exhumations juridiques et considérations sur les changemens physiques que les cadavres éprouvent en se pourrissant*. Par., 1831; *Leçons faisant partie du cours de médecine légale*. Par., 1821. (Orfila was a prolific writer. See Index Catalogue for other works.)—ORLAMUNDT, C. W. *Haandbog i den legale Medicin*. Kjøbenhavn, 1843.—OSOLINI, G. *Alcune note pratiche di medicina forense*. Lucca, 1881.—OTTO, F. J. *Anleitung zur Ausmittelung der Gifte; so wie die Erkennung der Blutflecken*. Braunschweig, 1856.—OUTLINE of lectures on medical jurisprudence in the University of Aberdeen, 1776.

PACINI, F. *Di alcuni pregiudizi in medicina legale*. Firenze, 1876.—PARÉ, A. *De renuntiationibus et cadaverum embamatibus tractatus*. Amstelredam, 1649.—PARIS, J. A., AND FONBLANQUE, J. S. M. *Medical jurisprudence*. Lond., 1823.—PARMENIONE, L. *Sammlung verschiedener Casuum medico-chirurgico-forensium, mit nützlichen Anmerkungen an das Licht gegeben*. Ulm, 1742.—PAULIER, A. B., AND HÉTET, F. *Traité élémentaire de médecine légale, de jurisprudence médicale et de toxicologie*. Par., 1881.—PEGO MENDEZ, F., AND GALVEZ DURAN, F. *Elementos de medicina legal militar*. Vitoria, 1901.—DE PEIRO, P. M., AND RODRIGO, J. *Elementos de medicina y cirugía legal*. Zaragoza, 1844; Adición, 1849.—PEIXOTO DI REGO, M. J. *Um capitulo de medicina legal*. Porto, 1882.—PELIKAN, E. V. *Beiträge zur gerichtlichen Medizin, Toxikologie und Pharmakodynamik*. Würzburg, 1858.—PELLACANI, P. *Lezioni di medicina legale*. Bologna, 1893.—PERES FURTADO GALVAO, J. *Tratado elementar de medicina legal*. Porto and Coimbra, 1858.—\*PERRONE. *Trattato elementare de med. leg.* Naples, 1840.—PETERMANN, A. *Casuum medico-legalium*. Leipz., 1708-1709.—PETERSON, F., AND HAINES, W. S. *Textbook of legal medicine and toxicology*. Phila., 1903 and 1904.—PFAFF, E. R. *Anleitung zur Vernehmung gerichtsarztlicher Blut-Untersuchungen*. Dresden, 1860.—VAN DER PFORDTEN, G. A. *Beiträge zur Geschichte der gerichtlichen Medizin aus den Justinianischen Rechtssammlungen*. Würzburg, 1838.—PRENGER, A. *Zur gerichtlichen Arzneiwissenschaft*. Coburg and Leipz., 1836.—PHELPS, E. J. *Lectures on topics connected with medical jurisprudence*. Burlington, 1881.—PIANTIERI, F. *Elementi di medicina legale*. Napoli, 1863.—PICHLER, W. *Die gerichtliche Medizin*. Wien, 1861.—PEIRO, P. M., AND RODRIGO, J. *Elementos de medicina y cirugía legal*. Madrid, 1839.—\*PITZNER, F. *Leitfaden bei gerichtlichen Leichenöffnungen*. Landshut, 1833; *Darstellung des zergliederten Gehirnes*. Landshut, 1835.—PHELPS, E. J. *Lectures on topics connected with medical jurisprudence*. Burlington, 1881.—\*PLACEK. *Jahresbericht der Unfallheilkunde, gerichtlichen Medizin und öffentlichen Gesundheitspflege*. Leipz.—PLASSELL, J. *Gerichtlich-medizinische Memoranda aus dem k. k. österr. Strafgesetze*. Innsbruck, 1854; *Einiges über die ärztlichen Sachverständigen in der neuen österr. Strafprozessordnung*. Innsbruck, 1850.—PLATNER, E. *Unter-*



suchungen über einige Hauptcapitel der gerichtlichen Arzneiwissenschaft. Leipz., 1820; *Quæstiones medicina forensis*. Lipsiæ, 1800.—PLAZ, A. G. *De medicina supra jurisprudentiam æstimanda*. Lipsiæ, 1782.—VON PLENCK, J. J. *Elementa medicina et chirurgia forensis*. Viennæ, 1786.—PLOUCQUET, W. G. *Commentarius medicus in processus criminales super homicidio, infanticidio, et embryocytia*. Argentorati, 1786.—POELCHAU, G. F. *Gerichtsärztliche Gutachten*. Riga, 1868.—POILLOUX, J. *Médecine légale criminelle*. Par., 1837.—\*POLSTORFF. *Leitfaden der qualit. Analyse und der gerichtl.-chemischen Analyse*. Leipz., N. d.—POMA, A. *Dizionario anatomico-medico-legale*. Padova, 1834.—POORE, G. V. *A treatise on medical jurisprudence*. New York, 1901.—POTGIETER, A. *Bijdrage tot de geregtelijke geneeskunde*. Leiden, 1834.—POUSSIELGUE, A. L. *Propositions sur divers sujets de médecine légale*. Par., 1804.—PRUNELLE, C. F. V. G. *De la médecine politique*. Montp., 1814.—PUCCINOTTI, F. *Lezioni di medicina legale*. Napoli, 1858.—PUGA BORNE, F. *Recopilacion de las leyes chilenas que se relacionan con la medicina legal*. Santiago, 1884.—PURLAS, I. P. 1870.—PURRINGTON, W. A. *A review of recent legal decisions affecting physicians, dentists, druggists*. New York, 1899.—PUTMAN, C. E. *Tratado práctico de medicina legal en relación con la legislación penal y procedimental del país*. Bogotá, 1896.—PUNEDDU, G. *Programma per il corso di medicina legale e di tossicologia d'igiene e di polizia medica*. Cagliari, 1858.—\*PYL, J. T. Editor of a number of legal journals. 1783-1793.

QUINZIO, C. *Manuale pratico di medicina legale militare*. Firenze, 1899.

RABUTEAU. *Éléments de toxicologie et de médecine légale appliquée à l'empoisonnement*. Par., 1873.—RAFFAELE, A. *Clinica di medicina legale*. Napoli, 1896; 1900; *Guida pratica alle perizie medico-legali civili e penali*. Napoli, 1895.—RAMON Y VEGA, A. *Compendio de práctica medico-forense aplicada á la actual legislación y arreglada á las ultimas disposiciones de la ciencia medico-legal*. Madrid, 1888.—RANBIHAN, JR., E. *De corpore delicti medicæ indagando*. Lipsiæ, 1783.—RASSIER, F. M. *De la valeur du témoignage des enfants en justice*. Lyon, 1892.—RAVAGLIA, G. *Della condizione dei periti medici in Italia*. Bologna, 1881.—RAVARY, C. *Les expertises médico-légales*. Par., 1901.—REBSAMEN, F. X. *Decas observationum medico-forensium epicrisibus physiologicis ornatum*. Viennæ, 1780.—REESE, J. J. *Text-book on medical jurisprudence and toxicology*. Phila., 1902; *The post-mortem imbibition of poisons more especially in its medico-legal relations*. Phila., 1890.—RÈGLEMENT du conseil médical pour les affaires civiles. Constantinople, 1870.—REINESIUS, T. *Schola jura-consultorum medica*. Lipsiæ, 1679.—REINSBERG, J. *The science of legal medicine*. Prazæ, 1896.—REMARKS on medical jurisprudence. Dublin, 1793.—REMER, W. H. G. *Lehrbuch der polizeilich-gerichtlichen Chemie*. Helmstadt, 1827.—RENARD, L. M. *La manière de procéder à l'ouverture des cadavres, et spécialement dans les cas de visite judiciaires*. Par., 1817; Also in *Médecine légale*, Par., 1819.—REY, A. *Traité de jurisprudence vétérinaire*. Par., 1865.—REYES, M. O. *La medicina legal y el médico legista*. México, 1896.—RICHTER, C. H. *Ausgewählte Abhandlungen und Gutachten aus dem Gebiete der gerichtlichen Medicin*. Stuttgart, 1838.—RICHTER, E. E. *Digesta medica, seu decisiones medico-forensis*. Lipsiæ, 1731.—RISTELHUEBER, J. *Rapports et consultations de médecine légale*. Par., 1821.—ROCKINGER, L. *Ueber einen Ordo judicarius, bisher dem Johannes Andreae zugeschrieben*. München, 1855.—ROLFFS, J. C. F. *Taschenbuch zu gerichtlich-medizinischen Untersuchungen*. Köln a. Rh., 1833; 1838. *Praktisches Handbuch zu gerichtlich-medizinischen Untersuchungen und zur Abfassung gerichtlich-medizinischer Berichte*. Berl., 1840.—ROOSE, T. G. A. *Grundriss medizinisch-gerichtlicher Vorlesungen*. Frankf. a. M., 1802; *Beiträge zur öffentl. und gerichtl. Arzneikunde*. Frankfurt, 1802.—ROSE. *Manuel d'autopsie cadavérique médico-légale*. Par., 1808.—ROSENSTENGEL, J. J. *Medicus practicus clinico-forensis*. Frankf. a. M., 1717.—ROSSELL, A. *Manual de medicina legal*. Madrid, 1848.—RUDECK, W. *Medizin und Recht*. Jena, 1899.—RYAN, M. *A manual of medical jurisprudence and state medicine*. Lond., 1836.—RUKHSHTAB, B. G. *On various changes in the legal chemical examination of internal organs*. Farmats., Mosk., 1900, v. VIII, p. 309.

SADUN, B. *Intorno alcuni malintesi che falsano l'indirizzo e la pratica della medicina legale*. Pisa, 1882.—DE SALLES, E. *Traité de médecine légale et jurisprudence médicale*. Par., 1838.—SALZER, J. M. *De inspectione et sectione legali*. Tubingæ, 1736.—SCHACHER, P. F. *De jureconsulto medico*. Lipsiæ, 1737.—SCHAEFFER, A. J. *Sammlung gerichtsarztlicher Gutachten*. Berl., 1852.—SCHALL-



- GRUBER, J. *Abhandlungen im Fache der Gerichtsärzneykunde*. Graz, 1821.—SCHAUENSTEIN, A. *Lehrbuch der gerichtlichen Medicin*. Wien, 1862.—SCHEMATA zum Einzeichnen von Befunden bei gerichtsarztlichen Untersuchungen am Gehirn. Tübingen, 1883.—SCHEMATA zum Einzeichnen von Befunden bei Untersuchung der Brust- und Bauch-Organe mit Skeletteinzeichnung. Tübingen, 1883.—SCHEMATA zum Einzeichnen von Befunden bei gerichtsarztlichen Untersuchungen am Schädel. Tübingen, 1883.—SCHEMA zur forensischen Obduction. Braunschweig, 1880.—SCHIRMER, G. J. *Observationes medico-forensis*. Francof. et Lipsiæ, 1779.—SCHLEGEL, J. C. T. *Collectio opusculorum selectorum ad medicinam forensis spectantium*. Lipsiæ, 1785.—SCHLEGEL, J. H. G. *Materialien für Staatsarzneiwissenschaft und praktische Heilkunde*. Meiningen, 1819.—SCHLOCKOW, Der preussische Physikus. Berl., 1886.—SCHMALZ, K. G. *Gerichtsarztliche Diagnostik*. Leipz., 1840.—SCHMIDTMÜLLER, J. A. *Handbuch der Staatsarzneikunde*. Landshut, 1804.—SCHMITT, G. *A brief of necroscopy*. New York and Lond., 1901.—SCHMITT, W. J., BACHMANN, C. L., KÜTTINGER, J. F. *Einige auserlesene medicinisch-gerichtliche Abhandlungen*. Nürnberg, 1813.—SCHNEIDER, P. J. *Ueber die Gifte in medicinisch-gerichtlicher und medicinisch-polizeylicher Rücksicht*. Tübingen, 1821; *Ueber die von Gerichtsärzten zu erstattenden Gutachten*. Freiburg i. Br., 1851.—SCHNEIDER, F. C. *Die gerichtliche Chemie*. Wien, 1852.—SCHÖMAN *Gerichtliche Medicin*. Jena, 1861.—\*SCHRAND, F. *Elementa medicinae forensis*. Pesth, 1802.—SCHUCHARDT, B. *Vergiftungen*. Tübingen, 1882.—SCHÜRMAYER, L. H. *Gerichtlich-medicinische Klinik*. Karlsruhe, 1844; *Lehrbuch der gerichtlichen Medicin*. Erlangen, 1850.—SCHVOB, A. *Sur la médecine légale chez les libéraux*. Strasbourg, 1861.—SCHWAAB. *Beitrag zur sanitäts-polizeilichen Obduction*. Münch. med. Wehnschr., 1898, v. XLV, p. 1437.—SCHWABE, C. W. *Das Leichenhaus in Weimer*. Leipz., 1834.—SCHWEICKHARD, C. L. *Medicinisch-gerichtliche Beobachtungen*. Strassburg, 1789.—SÉDILLOT, C. E. *Manuel complet de médecine légale*. Par., 1836.—SEIPP, J. P. M. B. *Valentini novellæ medico-legalis*. Gissæ Hassorum, 1704.—SEMPLE, C. E. A. *Essentials of forensic medicine, toxicology, and hygiene*. Phila., 1892.—SEN, S. C. *Bhais hajya Bichar*. Dacca, 1894.—SEYDEL, K. J. *Leit faden der gerichtlichen Medicin*. Berl., 1895.—SHORT, A. *A view of the extent and importance of medical jurisprudence, considered as a branch of education*. N. p., N. d.—SHTOLTS, V. *Handbook of medical jurisprudence*. St. Petersburg, 1890.—SIEBENHAAR, F. J. *Enzyklopädisches Handbuch der gerichtlichen Arzneikunde*. Leipz., 1838.—VON SIEBOLD, E. C. J. *Lehrbuch der gerichtlichen Medicin*. Berl., 1847; *Commentatio nexum jurisprudentiam inter et medicinam exhibens*. Marburgi, 1831.—SIEGFRIED, C. G. *Diss. caudales nonnullas medico-politicas in praxi clinico-forensi observandas*. Halæ, 1726.—SIKORA, M. M. *Conspectus medicinae legalis, legibus Austriaco-provinciislibus accommodatus*. Præge, 1780.—SILVESTRI, S. *Guida alle autopsie per i scope clinico e medico-legale*. Napoli, 1887.—SMITH, F. J. *Lectures on med. jurisprudence and toxicology*. Lond., 1900.—SMITH, J. G. *An analysis of medical evidences*. Lond., 1825; *Hints for the examination of medical witnesses*. Lond., 1829; *Principles of forensic medicine*. Lond., 1827; *Claims of forensic medicine*. Lond., 1829; *The principles of forensic medicine systematically arranged and applied to British practice*. Lond., 1824.—SMITH, W. R. *Introductory address to a course of lectures on forensic medicine*. Lond., 1869.—SNETIWEY, K. *Sammlung auserlesener gerichtlich-medicinischer Untersuchungen*. Prag, 1846.—SONDERMANN, J. G. *Diss. sistens quaestiones medico-forenses problematicas*. Jenæ, 1798.—SONNENSCHNEN, F. L. *Handbuch der gerichtlichen Medicin*. Berl., 1856.—DE SOUZA LIMA, A. J. *Manual de chimica legal*. Rio de Janeiro, 1892.—SPERANZA, C. *Sulla dignità della medicina legale*. Parma, 1833.—\*SPORER, G. M. *Grundrisse eines vollständigen Systems der Staatsarzneikunde*. Klagenfurt, 1837.—\*SPRENGEL, K. *Institutiones medicinae forensis*. Leipz., 1816.—STARCK, J. M. *De medicina utilitate in jurisprudentia*. Helmæstadii, 1730.—STEININGER, F. v. P. *Staatsarzneiwissenschaft*. Wien, 1793.—STEREOSCOPISCHER medicinischer Atlas *Gerichtliche Medicin*. Leipz., 1898.—STERNE, S. *Legal responsibility and accountability*. New York, 1874.—STOLL, J. *Staatswissenschaftliche Untersuchungen und Erfahrungen*. Zürich, 1812.—STOLTZ, V. *Manual for the study of legal medicine*. St. Petersburg, 1885.—STOREK, D. H. *An address on medical jurisprudence*. Bost., 1851.—STRASSMAN, F. *Lehrbuch der gerichtlichen Medicin*. Stuttgart, 1895.—STREIT, E. F. G. *Observationes quasdam et quaestiones ad medicinam forensis spectantes*. Lipsiæ, 1810.—STURM, J. S. *De privilegiis agrotorum, tot*



denen Freyheiten der Krancken. Vitembergæ, 1744.—SUCHOW, E. H. *Die gerichtlich medicinische Beurtheilung des Leichen befunds*. Jena, 1849.—SUE, P. *Aperçu général, appuyé de quelques faits sur la médecine légale*. Par., 1800.

TAPIA, A. A. *Reflexiones sobre algunos artículos del título ii del código penal*. México, 1877.—TARDIEU, A. *Etude médico-légale et clinique sur l'empoisonnement*. Par., 1867; *Etude médico-légale sur la strangulation*. Par., 1859; *Etude médico-légale sur les attentats aux mœurs*. Par., 1862; *Etude médico-légale sur l'avortement*. Par., 1864; *Question médico-légale de la pendaison*. Par., 1865; *Etude médico-légale sur l'infanticide*. Par., 1868; *Etude médico-légale sur la pendaison, la strangulation et la suffocation*. Par., 1870; *Etude médico-légale sur les maladies produites accidentellement ou involontairement par imprudence, etc.* Par., 1879; With ROUSSIN, Z. *Etude sur l'empoisonnement*. Par., 1867.—TAYLOR, A. S. *Elements of medical jurisprudence*. Lond., 1836; *Manual of medical jurisprudence*. Edited by T. STEVENSON. Phila., 1897; (Taylor's book has long been a justly standard work); *On poisons in relation to medical jurisprudence and medicine*. Phila., 1875; *The law in its relation to physicians*. N. Y., 1900.—TEICHMEYER, H. F. *Institutiones medicinae legalis*. Jena, 1767; *De cadaveris sectione legali*. Jena, 1742.—TEIXEIRA DE MATTOS, I. *Nonnullas observaciones et questiones medico-legales*. Amsterd., 1846.—TELGEMANN, R. F. *Dissertatio juridica de frustranea cadaveris inspectione in homicidio*. Helmstadii, 1731.—THIERRY, H. *De la responsabilité atténuée; étude médico-légale*. Par., 1891.—THOMSON, A. T. *Lecture, introductory to the course of medical jurisprudence*. Lond., 1831; *Syllabus of lectures on medical jurisprudence*. Lond., 1848; With AMOS, A. *Syllabus of lectures on medical jurisprudence*. Lond., 1830; *Vorlesungen über gerichtliche Arzneiwissenschaft*. Leipz., 1840.—TIDY, C. M. *Legal medicine*. (A well recognized authority.) Lond., 1882.—TIEGEL. *Lectures on medical jurisprudence*. Tokio, 1880.—TORTOSA, G. *Instituzioni di medicina forense*. Firenze, 1829.—TOSCANI, D. *Introduzione allo studio della medicina legale*. Roma, 1880.—TOURDES, G. *Des cas rares en médecine légale*. Strasbourg, 1840; With MEIZGUER, E. *Traité de médecine légale théorique et pratique*. Par., 1896.—TOWNSEND, J. F. *Relations between the professions of law and medicine*. Albany, 1854.—TRABAJOS de la Comisión de medicina legal é higiene pública de la real Academia de ciencias médicas. Habana, 1872; 1874.—TRAILL, T. S. *Outlines of a course of lectures on medical jurisprudence*. Edinb., 1841.—TRATTATO di medicina legale conforme al diritto italiano costituito. Milano, 1901.—TRÉBUCHET, A. *Jurisprudence de la médecine, de la chirurgie, et de la pharmacie*. Bruxelles, 1838.—TRILLER, C. F. *Exercitatio de vinculo quo medicina cum jurisprudentia cohaeret*. Vitebergæ, 1778.—TRINQUIER, F. V. *Plan raisonné d'un cours de médecine légale*. Strasbourg, 1840; *\*Système complet de médecine légale*. Montp., 1836-1838; *Exposition historique et appréciation des secours empruntés par la médecine légale à la chirurgie*. Strasbourg, 1838.—TROPPANNEGER, C. G. *Decisiones medico-forenses*. Dresden and Neustadt, 1733.—TSCHOUJIN, G. F. *Gerichtliche Thierarzneykunde*. Karlsruhe and Baden, 1822.—TYRMAN, J. *Die Verletzungen des Ohres*. Vienna, 1903.

UDEN, K. F. *Ueber die Glaubwürdigkeit der Medizinalberichte in peinlichen Rechtshändeln*. Berl., 1780.—USHAKOFF, F. S. (Cadavers for the chair of medical jurisprudence of the Imperial Army Medical Academy for fifteen years.) St. Petersburg, 1900.

VALENTINI, M. B. *Corpus juris medico-legale*. Francf. a. M., 1722; *Pandectæ medico-legales*. Francf. a. M., 1701; *Novellæ medico-legales*. Francf. a. M., 1711.—VALENTI Y VIVO, I. *Curso elemental de medicina legal*. Barcelona, 1873.—VALLADA, D. *Elementi di giurisprudenza medico-veterinaria*. Torino, 1883.—\*VALLE DEL, J. F. *Chirurgia forensis*. Madrid, 1796.—VARSHAVSKI, V. *Review course of legal medicine*. Kiev, 1891.—VERDIER, J. *La jurisprudence de la médecine en France*. Alençon, 1763; *La jurisprudence particulière de la chirurgie en France*. Par., 1764.—VIBERT, C. *Précis de médecine légale*. 1903 (An excellent work, giving an account of the French laws bearing upon this subject); *Précis de toxicologie clinique et médico-légale*. Par., 1900.—VIDAL, L. *Les certificats médico-légaux usuels*. Par., 1900.—\*VIESS, F. B. *Vorlesungen über gerichtliche Arzneikunde*. Vienne, 1817.—VILLEBRUN, E. *Des ongles, leur importance en médecine judiciaire*. Lyon, 1883.—VAN VISVLIET, E. P. *Specimen juridicum sistens casus quosdam in quibus scientia medica jurisconsultis est necessaria*. Lugd. Bat., 1760.—VOLPRECHT, C. *Diss. acad., qua disquiritur an medicis*

*impune occidere liceat.* Kiliz, 1704.—VORSCHRIFT für die Vornahme der gerichtlichen Todtenbeschau von 28 Jänner, 1855. Amtliche Handausgabe. Wien, 1861.—\*VOY. *Introduction du systema jurisprudentia medic. d' Alberti et les elemens de jurisprudent. medicale de T. R. Buk.* Lond., 1825.

WACHHOLZ, L. *Manual of medical jurisprudence.* Kraków, 1899.—WAGNER, P. P. V. *Anleitung zur gerichtlichen Arzneikunde.* Wien, 1833.—WALBAUM. *Erfahrungen aus dem Gebiete der Hygiene, Sanitätspolizei und gerichtlichen Medicin.* Leipz., 1871.—WALD, H. *Gerichtliche Medicin.* Leipz., 1858.—WARREN, R. B. *A familiar forensic view of man and law.* Columbus, 1860.—WEBSTER, J. *An essay on medical jurisprudence.* Phila., 1824.—WEIL, G. D. *De l'exercice illégal de la médecine et de la pharmacie.* Par., 1886.—WEIZ, F. A. *Vermischte Beiträge zur gerichtlichen Arzneigelahrtheit in verschiedenen vorgekommenen Fällen.* Frankenthal, 1787.—WENDLER, C. A. *De ambigua corporis delicti notione.* Lipsiæ, 1845.—WHARTON, F., AND STILLÉ, M. *A treatise on medical jurisprudence.* Phila., 1855.—WIENER, D. *Sammlung gerichtlich-medicinischer Obergutachten.* Berl., 1891; *Methodik Diagnostik und Technik bei gerichtsarztlichen Obductionen menschlicher Leichen.* Stuttgart, 1882.—WILBRAND, F. J. J. *Über das Wesen des Menschen vom gerichtlich-medicinischen Standpunkte aus.* Giessen, 1882.—WILL, J. R. *Ueber das Formelle bei gerichtlich-chemischen Untersuchungen.* Cassel, 1857.—WILDBERG, C. F. L. *Handbuch der gerichtlichen Arzneiwissenschaft.* Berl. 1812; *Bibliotheca medicina publica.* Berl., 1819; *Lehrbuch der medizinischen Rechtsgelahrtheit zum Unterricht für Rechtsgelehrte.* Leipz., 1826; *Rhapsodien aus der gerichtlichen Arzneiwissenschaft.* Leipz., 1822; *Codex medico-forensis.* Berl., 1842; *Anleitung zu den polizeilich medizinischen Geschäften der Physiker.* Erfurt, 1833; *Mag. für die gerichtl. Arzneiwissenschaft.* Berl., 1831-1834; *Jahrbücher der Gesamten Staatsarzneikunde.* Leipz., 1835; *Die Geschäftsführung der Physiker als Polizei und Gerichts-Aerzte in Beyspielen.* Erfurt, 1824; *Von den gerichtlich-medicinischen Geschäften der Physiker.* Erfurt, 1822; *Anweisung zur gerichtlichen Zergliederung menschlichen Leichname.* Berl., 1817.—WILHELM, M. *De l'aspect extérieur du cadavre au point de vue médico-légal.* Par., 1878.—WILLIAMS, P. C. *An examination of Prof. Reese's review of the trial of Mrs. Wharton for the murder of General Ketchum.* Balt., 1872.—WILLIAMS, S. W. *A catechism of medical jurisprudence.* Northampton, 1835.—\*WINSLOW. *Medico-legal evidence.*—WIPPERMANN, O. M. *Diss. juridica.* Rintellii, 1720.—WISTRAND, A. T. AND A. H. *Handbok i rätts medicinen.* Stockholm, 1857.—WISTRAND, A. T. *Handbok i forensiska medicinen.* Stockholm, 1838; *Tabeller-för medicolegala obductioner.* Sigtuna, 1838; *Rättsmedicinska afhandlingar.* Stockholm, 1863; *Granskning af lagberedningens förslag till straffbalk sa vidt den rör rättsmedicinen.* Stockholm, 1846.—WISTRAND, H. *Rättsmedicinsk kasuistik.* Stockholm, 1855.—WITTHAUS, R. A. *On the post-mortem imbibition of poisons, and the chemico-legal aspect of embalming. Researches of the Loomis Laboratory.* N. Y., 1890, v. I; With BECKER, T. C. *Medical jurisprudence, forensic medicine, and toxicology.* N. Y., 1894; (A new edition of this work is in preparation).—WOODMANN, W. B., AND TIDY, C. M. *A handy book of forensic medicine and toxicology.* Lond., 1877.—WORMLEY, T. G. *Micro-chemistry of poisons.* New York, 1885.—WUNDERBAR, R. J. *Biblisch.-talmudische Medicin.* Riga and Leipz., 1865.

YAMAMOTO, Y. *Forensic medicine.* Tokio, 1886.—YANEZ, T. *Elementos de medicina legal y toxicologia.* Madrid, 1884.—YOSHIYAMA, S. *Collection of medico-legal cases in Japan.* Tokio, 1892-1893.

ZABITZEANO, K. 1850.—ZACCHIAS, P. *Questiones medico-legales.* Lipsiæ, 1650.—ZIEGLER, C. J. A. *Beobachtungen aus der Arznei-wissenschaft Chirurgi- und gerichtliche Arznei-kunde.* Leipz., 1787.—ZUINO, G. *Compendio di medicina legale e giurisprudenza medica.* Milano, 1890.—ZITTMANN, J. F. *Medicina forensis.* Franckf. a. M., 1706.

(References marked with a \* have not been verified.)



## GLOSSARY INDEX



(The Roman numeral letters refer to chapters; the Arabic numerical figures to pages. Cross-references are frequent, and explanatory notes will be found inserted here and there.)

### A

- Abdomen, enlargement of veins of, 61  
 injecting embalming fluid through, 286  
 regional landmarks of; epigastric, umbilical, hypogastric, right and left hypochondriac, right and left lumbar, and right and left iliac.  
 striæ of, 61  
 Abdominal cavity, abnormal contents of, 20, 85, 86, 88, 89  
 bile in, 86  
 dissection of, in animals, 383, 393  
 diseases of organs of, x, p. 159; xi, p. 199; xii, p. 213; xviii, p. 275  
 examination of, 88, 90, 91, 161, 162, 167, 275, 438, 439, 440, 442, 443  
 exposure of, in animals, 375  
 exudates in, 86  
 fluid in, 85  
 foreign bodies in, 88  
 free bodies in, 20, 88  
 in anthrax, 292  
 incision, closure of, 285  
 organs of, critical examination of, x, p. 159  
 removal of organs of, 91, 442  
     from animals, 378, 391, 393  
 superficial examination of organs of, vi, p. 79; xxvii, p. 440  
 surgical instruments in, 88  
 technic of opening, vi, p. 79; xviii, p. 275; xxvii, p. 440  
 topographic examination of organs in, vi, p. 79; xviii, p. 275; xxvii, p. 440  
 transudates in, 86  
 hernia, 90  
 incision, method of closing, 285  
 muscles, post-mortem rigidity of, 51  
 organs, condition of, in cholera Asiatica, 293  
 pregnancy, see Pregnancy, extra-uterine.  
 sympathetics, 14  
 tuberculosis of, 85, 312  
 typhus, see Typhoid fever.
- Abnormalities, see Malformations and the organs themselves.  
 Abortion, causes of, 303, 406  
     criminal, 406  
     abscess of lung following, 147  
     history of a case of, 279  
 Abrachius, upper limbs absent, lower limbs well formed.  
 Abrasions in skin of hands, how detected, 38  
 Abrine, toxicology of, 421  
 Abscess, actinomycotic, 290  
     amœbic, of liver, 174  
     appendiceal, 171  
     as a cause of death, 408  
     cold, 311  
     embolic, of lung, 147  
     metastatic, of heart, 130  
     multiple, of liver, 219  
     of bones, 268  
     brain, 246  
     liver, 159, 213, 219  
     lungs, 147  
     multiple, 147  
     solitary, 147  
     œsophagus, 190  
     prostate, 212  
     psoas muscles, 198  
     testicles, 211  
     perinephritic, 203  
     periurethral, 297  
     retropharyngeal, 292  
     subdiaphragmatic, 159  
     subphrenic, 159  
     vaginal, 211  
 Absence of anus, 69  
     clitoris, 69  
     hymen, 69  
     kidneys, 199  
     nose, 68  
     penis, 68  
     sternum, 68  
     urethra, 69  
 Absorbent cotton, use of, in preservation of body, 283  
 Absorption in pneumonia, 153  
     of deleterious gases, 463  
 Acardia, 113  
 Acarus scabiei, 320

- Accessory adrenals, 179  
 kidneys, 199  
 livers, 204, 367  
 lung-tissue, 91  
 pancreases, 220  
 sinuses, exposure of, in animals, 389  
 spleens, 162  
 thyroid glands, 96
- Accidental submersion, 463  
 traumatism, 463
- Accidents of pregnancy, 247, 460  
 puerperal, 460
- Accephalus, 67
- Acetic acid as a fixing fluid, 329  
 toxicology of, 427
- Acetone, in diabetic urine, 222
- Acholia, deficiency or absence of the secretion of bile.
- Achorion Schönleini, Plate V, No. 9, facing page 350
- Acid, acetic, as a fixing fluid, 329  
 toxicology of, 427  
 beta-oxybutyric, in diabetes, 222  
 toxicology of, 421  
 boric, toxicology of, 429  
 carbolic, effect of, on skin, 65  
 on urine, 425  
 toxicology of, 421, 422, 423, 425, 426  
 chromic, as a fixing fluid, 330  
 chromo-acetic, as a fixing fluid, 331  
 chromo-aceto-osmic, as a fixing fluid, 331  
 chromo-formic, as a fixing fluid, 331  
 chromo-nitric, as a fixing fluid, 331  
 chromo-osmic, as a fixing fluid, 331  
 corrosive, toxicology of, 421, 423, 424, 425  
 diacetic, in diabetes, 222  
 filicic, toxicology of, 421  
 helvellac, toxicology of, 423, 425  
 hydrochloric, toxicology of, 422, 427  
 hydrocyanic, toxicology of, 50, 421, 425, 430  
 hydrofluoric, toxicology of, 422  
 nitric, as a fixative of tissue, 336  
 toxicology of, 422, 423, 427  
 osmic, as a fixative of tissue, 336  
 toxicology of, 422  
 oxalic, toxicology of, 424, 426, 427  
 oxybutyric, toxicology of, 421  
 parabanic, toxicology of, 424  
 picric, as a fixative, 337  
 toxicology of, 423, 424  
 picro-acetic, as a fixative, 337  
 pyrogallac, in study of anaërobic bacteria, 353  
 salts of metals, toxicology of, 424  
 sulphuric, toxicology of, 426  
 sulphurous, toxicology of, 422  
 uric, in calculi, 207
- Acids, corrosive, toxicology of, 421, 423  
 effect of, on skin, 65, 426  
 mineral, toxicology of, 424, 426
- Acids, organic, employed as fixing fluids, 329  
 vegetable, toxicology of, 426
- Acne, 62  
 of vulva, 69
- Aconite, toxicology of, 411, 427
- Aconitine, toxicology of, 421, 422
- Acquired anomalies and deformities, 19, 72
- Acrodynia, 351
- Acromegaly, 223, 246  
 cause of, 70  
 thymus gland present in, 95
- Actinomycosis, 290, 373  
 abscesses in, 290  
 in peritoneum, 159  
 of alimentary tract, 290  
 appendix, 171  
 brain, 246, 255  
 heart, 134, 290  
 lungs, 290  
 skin, 290  
 thorax, 290  
 primary lesion of, 290  
 synonyms of, 449
- Acute, see the diseases themselves.
- Addison's disease, 16, 180  
 cause of, 181  
 complications of, 452
- Adenitis, mediastinal, simple, 146  
 suppurative, 146  
 tuberculous, 146
- Adenocystomata of kidney, 220  
 liver, 220
- Adenoids in chronic nasal catarrh, 141
- Adenoma of bladder, 205  
 ciliary body, 75  
 intestines, 171  
 liver, 220  
 lungs, 155  
 ovary, 209
- Adhesions, 20, 87, 195
- Adhesive pericarditis, 144
- Adipocere, 46, 49
- Adiposis dolorosa, 59
- Administration of poisons, symptoms observed after, 421
- Adonis, toxicology of, 424
- Adrenals, 175  
 aberrant, 204, 220  
 accessory, 179  
 atrophy of, 180  
 color of, 180  
 dimensions of, 369  
 disease of, 16, 60, 180, 181, 324  
 examination of, 14, 15, 175, 179, 180, 276, 442  
 fixatives for, 340  
 measurements of, 369  
 miliary tubercles in, 180  
 removal of, 15, 176, 180, 280  
 in animals, 381  
 situation of, 179

- Adrenals, tumors of, 180**  
     weight of, 369  
**Adynamic fever, see Typhoid fever.**  
**Æstivo-autumnal malaria, 322**  
**Affections, see the various diseases.**  
**Agaricus bulbosus, toxicology of, 423**  
**Age, 55, 215**  
     apparent, 55  
     estimation of, 55, 58  
     real, 55  
**Agglutinative reaction, 118**  
**Ague, see Fever, malarial.**  
**Air emboli, method of discovering, in thoracic cavity, 100**  
     embolism of lungs, 148  
**Air-cells, hemorrhage into, 148**  
**Air-passages, diseases of, ix, p. 141**  
     exposure of, 256  
     examination of, xvi, p. 256  
     foreign bodies in, 146  
**Albinos, hair of, 76**  
**Albuminous degeneration, fixative for, 338**  
     periostitis, 271  
**Albuminuria in erysipelas, 295**  
     pregnancy, 448  
     scarlatina, 448  
     smallpox, 448  
**Alcohol as a fixing fluid, 330, 337, 338**  
     preserving fluid, 327, 328, 342, 343  
     toxicology of, 421, 423, 427  
**Alcoholism, 427**  
     as a cause of abortion, 407  
     cirrhosis of liver due to, 216, 427  
     delays post-mortem rigidity, 51  
     gastritis due to, 427  
     gastro-intestinal tract in, 427  
     genito-urinary tract in, 427  
     nervous system in, 427  
     synonyms of, 452  
     toxicology in, 421  
     vascular system in, 427  
**Alimentary tract, measurements of, 366**  
     suspicious undissolved foreign bodies in, 418  
     tuberculosis of, 312  
**Alkalies, toxicology of, 421, 424, 427**  
**Alkaline carbonates, effect on urine, 424**  
**Alkalinity of gastric contents, 418**  
**Alkaloidal poisons, 421**  
**Alkaptonuria, 223**  
**Alum hæmatoxylin, 341**  
**Alveolar process, absent, 68**  
     deformities of, 68  
**Amanita phalloides, 418, 424**  
**Amanuensis at postmortem, 22**  
     duties of, 22  
**American Anthropometric Society, 4**  
     insurance standard of weight, 358  
**Ammonia, to remove iodine stains, 286, 421**  
     toxicology of, 19, 21, 411, 421, 422, 423  
     use of, by injection, as a sign of death, 47  
**Ammoniaemia, effect on urine, 424**  
**Ammonium urate, 207**  
**Amœba coli, method of fixing, 338**  
**Amœbic abscess of liver, 174**  
     dysentery, 173, 219  
**Amputation, complications of, 462**  
     muscular twitchings in limbs after, 47  
     of hand in spreading gangrene, 45  
     synonyms of, 461  
**Amussat's hammer, 243**  
**Amyl nitrite, toxicology of, 21, 423, 425**  
**Amyloid degeneration, 129, 307, 418**  
     fixatives for, 338  
     of arteries, 135  
     kidney, 199  
     liver, 214  
     spleen, 164  
     reaction, 179  
**Anæmia, blood changes in, 122**  
     causes of, 116  
     cerebri, 247  
     definition of, 116  
     essential, 116  
     idiopathic, 116  
     von Jaksch's, 124  
     of brain, 247  
     lungs, 148  
     pernicious, 121, 195  
     primary, 116  
     progressive pernicious, 121  
     secondary, 116  
     simple, 116  
     splenic, 123  
     symptomatic, 116  
     synonyms of, 452  
     varieties of, 116  
**Anæmic infarcts of kidney, 201**  
     spleen, 163  
**Anaërobic growth, Wallis's method for, 349**  
     methods, 352  
     organisms, 349  
**Anæsthetic form of leprosy, 299**  
**Anatomical board, 5**  
     warts, 318  
**Anatomy, morbid, see diseases of, under the various organs.**  
     study of, at postmortems, 3  
**Anchylostomum duodenale, 121, 321**  
**Anencephalus, 67**  
**Aneurism, 137**  
     anastomotic, 138  
     arteriovenous, 139  
     by distention, 138  
     erosion, 138  
     rupture, 138  
     cirroid, 138, 140  
     conditions associated with, 138  
     congenital, 139  
     contour of blood-vessels in, 18  
     cylindrical, 138  
     death from, 407, 408  
     deformity due to, 72  
     dissecting, 138



- Aneurism, endothelial lining of, 130**  
 etiology of, 139  
 false, 137  
 femoral, mistaken for hernia, 139  
 fibrous clot of, 138  
 fusiform, 137  
 gluteal, mistaken for abscess, 139  
 hernial, 137  
 infective, 138  
 miliary, 138  
 mycotic, 138  
 of animals, 383  
   aorta, method of removing, 100  
   bones, 274  
   carotid mistaken for sarcoma, 138  
   cerebral arteries, 247  
   heart, 130  
     location of, 130  
     sinus of Valsalva, 131, 138  
 pathology of, 138  
 rupture of, as a cause of death, 138, 139  
 saccular, 138  
   treated by wiring and electrolysis, 139  
 seats of, 139  
 spurious, 137  
 syphilitic, 308  
 traction, 138  
 traumatic, 138  
 true, 137  
 varieties of, 137
- Angina pectoris, 129, 455**
- Angioma of arteries, 137**  
 bone, 273  
 brain, 255  
 iris, 75  
 kidney, 204  
 liver, 220  
 spleen, 164  
 tonsils, 143
- Angiosarcoma of nasal passages, 141**
- Angular method of removing the calvarium, 226, 227**
- Anhydræmia, 116**
- Anideus, 67**
- Anilin blue as a stain, 341**  
 colors, effect on urine, 425, 426  
 toxicology of, 421, 422, 423, 425, 426  
 yellow, toxicology of, 422
- Animal poisons, 421**
- Animals, aneurism in, 383**  
 domestic, contagious diseases from, 373  
 inoculation of, 353, 354  
 intraperitoneal inoculation of, 354  
 intravenous inoculation of, 354  
 irritant poisons, 421  
 lower, value of necropsies upon, 373  
 method of cutting ribs of, 377  
 methods of killing for post-mortem examinations, 374  
 most used for inoculation, 353
- Animals, post-mortem examination of, 355, xxv, p. 373**  
 preparation of, for inoculation, 353  
 removal of extremities of, 375  
   hide of, 375  
 site of inoculation in, 354  
 subcutaneous inoculation of, 354  
 taking temperature of, 353  
 toxic poisons, 421  
 weight of, 353  
 where and how to be killed for a necropsy, 374
- Animation, suspended, 46**
- Ankylosing arthritis, 264**
- Ankylosis, 71, 263, 264, 267**  
 congenital, 267  
 false, 267  
 mistaken for post-mortem rigidity, 51  
 of cricoarytenoid joint, 143  
 traumatic, 71
- Ankylostoma americana, 170**  
 caninum, 170  
 duodenale, 121
- Anomalies, 17, 19, 70**  
 congenital, of brain, 249  
   cord, 249  
   heart, 113  
   liver, 220  
   nails, 77  
   pancreas, 220
- Anopheles mosquitoes, as cause of malaria, 322**
- Anterior poliomyelitis, 351**
- Anthracosis, 16, 155, 216**
- Anthrax, 291, 373, 449**  
 of abdomen, 292  
   brain, 292  
   kidneys, 292  
   spinal cord, 292  
   thorax, 291  
   spores of, long lived, 356
- Anthropometric Society, American, 4**
- Antifebrin, toxicology of, 423, 425**
- Antimony, toxicology of, 418, 421, 422, 423, 428**
- Antipyretics, 421, 422, 425**
- Antipyrin, toxicology of, 422, 425**
- Antiseptic dressings for post-mortem wounds, 43**
- Antiseptics, synthetical, 421**
- Antiserum test for diagnosis of human blood, 118**
- Antrum of Highmore, empyema of, 141**
- Anus, absence of, 69**  
 artificial, 20  
 atrophy of, 69  
 diseases of, 69, 70, 307, 457  
 examination of, 14, 15, 438  
 fissures of, 69  
 hypertrophy of, 69  
 tumors of, 69
- Aorta, abdominal, 4, 139, 197, 365, 442**

- Aorta**, diseases of, 42, 72, 100, 111, 133, 134, 139, 156, 307  
 endocardial vegetations of arch of, 100  
 examination of, 14, 15, 105, 111, 197  
   in animals, 383  
 hypoplasia of, 136  
 removal of, 15, 100, 111, 197  
   in animals, 381, 383  
 thoracic, 14, 135, 139, 365  
**Aortic valves**, diseases of, 129, 133, 135, 307  
 examination of, 103, 104  
 measurements of, 103, 365  
**Aortitis**, 137  
**Aparathyroidism**, 97  
**Aphthæ of vulva**, 69  
**Apnoea**, 21  
**Apodia**, 70  
**Apomorphine**, toxicology of, 424  
**Apoplexia neonatorum**, 247  
**Apoplexy**, cerebral, *see also* Brain, hemorrhage of.  
   condition of eye in, 48  
   mistaken for alcoholism, 21  
   pancreatic, 221  
   pulmonary, 456  
**Appendicitis**, 171, 172  
   catarrhal, 171, 172  
   follicular, 171  
   gangrenous, 172  
   infective, 172  
   obliterative, 172  
   organisms of, 171  
   parasites in, 171  
   suppurative, 171  
**Appendix vermiformis**, 171  
   articles found in, 171  
   diseases of, 165, 171, 290, 299, 411  
   examination of, 15, 89, 276, 443  
   measurements of, 171, 366  
   removal of, 16, 64  
   tumors of, 172  
   Virchow's dictum concerning, 89  
   weight of, 366  
   worms in, 171  
**Arachnoid**, 231, 233, 241, 251, 252, 313  
   weight of pia and, 362  
**Arecoline**, toxicology of, 422, 423, 424  
**Argyria**, 422  
**Arsenic**, 418  
   detection of, by X-rays, 429  
   toxicology of, 411, 421, 422, 423, 424, 425, 426, 428, 429  
**Arsenical melanosis**, 427  
   poisoning, accidental, 428  
   localization of, 429  
   varieties of, 428  
**Arsenism**, 422  
**Arseniuretted hydrogen**, toxicology of, 423, 425  
**Arterial embalming**, 286  
   sclerosis, 223  
**Arteries**, 129  
   amyloid degeneration of, 135  
   aneurismal, in tuberculosis, 316  
   aneurisms of, 137  
   arteriosclerosis of, 134  
   atheroma of, 20, viii  
   atrophy of, 135  
   calcareous infiltration of, 135  
   cerebral, aneurism of, 247  
   condition of, after death, 47  
   coronary, 129  
   diseases of, 18, 20, 21, 128, viii, p. 134;  
     221, 223, 249, 253, 306, 308, 316, 407,  
     411, 431  
   examination of, 439, 440  
     in animals, 383  
   fatty degeneration of, 135  
   hyaline degeneration of, 136  
   hypertrophy of, 136  
   hypoplasia of, 136  
   iliac, curvatures in, 20  
   inflammations of, 136  
   morbid changes in, 134  
   pipe-stem, 129  
   pulmonary, how distinguished from  
     veins, 110  
   syphilis of, 137  
   tuberculosis of, 137  
   tumors of, 137  
**Arteriosclerosis**, 134  
   death from, 408  
   diffuse, 135  
   elastic tissue in, 134  
   in aortic incompetency, 133  
   nodular, 135  
   of penis, 68  
   pathology of, 134  
   seat of, 134  
   senile, 135  
   special forms of, 135  
   stages of, 134  
   syphilitic, 307  
**Arteriovenous aneurism**, 138  
**Arteritis obliterans**, 136  
**Arthritis**, 153, 261, 461  
   acute, 261  
   ankylosing, 261, 264  
   deformans, 136, 262, 265, 266  
   fibrinous, 261  
   gonorrhœal, 263, 297  
   gouty, 261, 263  
   infective, 261  
   neurogenous, 264  
   obliterative, 250  
   pneumococcal, 264  
   purulent, 261, 262  
   rheumatic, 262, 263, 266  
   scarlatinal, 305  
   serofibrinous, 261  
   serous, 261  
   spinal, 264  
   syphilitic, 261, 264  
   toxic, 264

- Anterior interscapular*, 261, 264, 265  
*ulcerative*, 262  
*Antropogathes*, 246, 261  
*Anthropomyia*, 261  
*Asplenosis*, 73, 143  
*Ascaris lumbricoides*, 321  
*Ascites*, associated conditions of, 85  
     detection of, 66  
*Asporous*, a spore lying within a special  
     sporecase or ascus, Plate v. No. 12, 350  
*Asiatic cholera*, synonyms of, 449  
*Asmata*, 67  
*Asparagus*, effect on urine, 424  
*Aspergillus*, 151, 152  
*Asphyxia*, 21, 415  
     suicide by, 462  
*Asphyxiation*, distention of right heart  
     after, 100  
*Assistants*, 6, 37  
*Asthma*, 144, 456  
*Atavism*, congenital characteristics de-  
     rived from remote ancestors  
*Ataxia*, hereditary, 248  
     locomotor, 248  
*Ataxic fever*, see Typhoid fever  
*Atelectasis of lungs*, 147  
     fetal, 147  
*Atelectatic bronchiectasis*, 144  
*Atheroma*, 20, 134, 307  
*Atheromatous kidneys*, 179  
*Atony of gastric walls*, 191  
*Atresia of glans penis*, 68  
     urethra, 69  
     uterus, 209  
     oris, 68  
*Atrophic bronchitis*, 145  
     changes in pancreas, 196  
     chronic nasal catarrh, 141  
     cirrhosis of liver, 216  
     shape of ensiform appendix in, 94  
     emphysema of lungs, 150  
     endometritis, 210  
     gastritis, 192  
*Atrophy*, 70  
     acute yellow, of liver, 18, 213  
     and hypertrophy in same part, 18  
     brown, 129  
     cause of, 70  
     congenital, 70  
     local, 70  
     location of, 70  
     of adrenals, 180  
     anus, 69  
     arteries, 135  
     bone, 264  
     clitoris, 69  
     fingers, 68  
     glans penis, 68  
     limbs, 56  
     liver, acute yellow, 425  
     marrow, 267  
     ovary, 209  
     scrotum, 69  
*Atrophy of testicles*, 69, 211, 419  
     testes, 68  
     tongue, 68  
     ureters, 209  
     vulva, 69  
     progressive muscular, 253  
     senile, 129  
*Atropine*, effect on pupil after death, 4  
     toxicology of, 421, 422, 423, 424, 425  
     426, 429  
*Aurania*, toxicology of, 422  
*Autodigestion*, 88  
*Auto-intoxication*, 413, 419  
*Autopsy*, 87  
*Autopsy*, see Postmortem  
     first use of word, 2  
*Autistic monster*, 67  
*Average height*, 358  
     weight, 358  
*Avoirdupois weight*, 357  
*Axes*, visual deviation of, 73, 74  
*Axillary cellulitis*, treatment of, 44  
     glands, how exposed, 81  
*Axis-cylinders*, fixatives for, 339  
*Ayer Clinical Laboratory of the Penn-  
     sylvania Hospital*, 10  
*Azygos veins*, 15
- B**
- Bacilli*, how distinguished from bacteria  
     349  
*Bacillus anthracis*, 290, 350  
     diphtheria, see *Corynebacterium diph-  
         theria*  
     lepra, see *Mycobacterium lepra*  
     mallei, see *Corynebacterium mallei*  
     oedematis maligni, 350  
     tetani, 350  
*Bacteria*, cultures of, how prepared for  
     inoculation, 353  
     effect on urine, 424  
     how distinguished from bacilli, 349  
     in the body, 207  
     phosphorescent, 404  
*Bacterial spores*, longevity of, 356  
     toxic poisons, 421  
*Bacteriologic examination*, care of instru-  
     ments in, 346  
     of bile-ducts, 194  
     gall-bladder, 194  
     investigations, 14, xxiii, p. 346  
*Bacterium aerogenes capsulatum*, 350  
     coli commune, 86, 160, 350  
     dysenteria, 350  
     enteritidis, 350  
     influenza, 350  
     paratyphoid, 350  
     pestis, 301, 350  
     pneumonia, 151, 350  
     proteum, see *B. vulgare*, 161, 350  
     pyocyanum, 41, 42, 161, 350  
     rhinoscleromatis, 350



- Bacterium* septicæmiæ hæmorrhagicum, 350  
     *typhi murium*, 350  
     *typhosum*, 350  
     *vulgare*, 161, 350  
 Balsam of copaiba, toxicology of, 422  
 Barium salts, toxicology of, 424  
 Bark, cinchona, toxicology of, 422  
 Barlow's disease, 126, 271  
 Bartholinian glands, enlargement of, 69  
 Baryta, toxicology of, 424  
 Basal ganglia, fixatives for, 339  
 Basedow's disease, 97, 147, 258  
 Base of skull, examination of, 233  
 Basins, enamelled, 33  
 Battery fluid, toxicology of, 8  
 Beaded ribs of rhachitis, 72  
 Beale's Prussian blue, 345  
 Béclard's sign, 278  
 Bedbug, 319  
 Bed-sores, location of, 66  
 Belladonna, toxicology of, 421, 422, 424  
 Bellows for inflating viscera, 34  
 Bensley's solution for fixing tissues, 335  
 Benzaldehyd, toxicology of, 425  
 Benzokoll, toxicology of, 425  
 Beriberi, 292, 351  
 Bertillon classification of causes of death, xxviii, p. 448  
     system, 53  
 Beta-oxybutyric acid, 222, 421  
 Bichlorid of mercury as a fixative, 333  
     effect on urine, 425, 426  
 Bichlorid-tablet solution for preserving tissues, 326  
 Bichromate of potassium, 8, 234, 327, 331  
 Bile, 214  
     collection of, 193  
     composition of, 214  
     in abdominal cavity, 86  
         blood, 120  
         gall-bladder, 195  
     pigments, 214  
     specific gravity of, 214  
 Bile-ducts, bacteriologic examination of, 194  
     congenital absence of common, 215  
     diseases of, 194, 195  
     examination of, 162, 187, 195, 442  
         in animals, 391, 393  
 Bilharzia hæmatobia, 202, 207  
 Biliary calculi, see Gall-stones.  
 Bilirubin crystals, 120  
     infarcts, 201  
 Binocalate of potassium, effect on urine, 424, 426  
 Biologic blood-test, 118  
 Biondi-Heidenhain stain, 341  
 Birds, post-mortem examination of, 394  
 Bishop, post-mortem on, 7  
 Bismarck brown, toxicology of, 422  
 Bismuth, toxicology of, 423  
 Bisulphid, carbon, toxicology of, 421, 422  
 Black urine, causes of, 206  
 Bladder, adenoma of, 205  
     *Bilharzia hæmatobia* of, 207  
     carcinoma of, 205  
     *cysticercus* of, 207  
     dimensions of, 368  
     diseases of, 68, 126, 127, 205, 206, 224, 313, 427, 459  
     *Distoma hæmatobium* of, 207  
     *echinococci* of, 207  
     *Eustrongylus gigas* of, 207  
     examination of, 14, 15, 68, 181  
     extroversion of, 68  
     fibroma of, 205  
     filaria of, 207  
     gangrene of, 205  
     inflammation of, see Cystitis.  
     inversion of, 206  
     measurements of, 368  
     mixed tumors of, 206  
     myoma of, 205  
     parasites in, 207  
     pentastoma of, 207  
     pockets in walls of, 206  
     removal of, 184, 442  
         in animals, 386  
     rupture of, 205  
     sarcoma of, 205  
     tuberculosis of, 205, 314  
     tumors of, 205  
     weight of, 368  
 Bleeding, see Hemorrhage.  
     use of the word in court instead of "hemorrhage," 401  
 Blenorrhagia of adult, synonyms of, 450  
     complications of, 450  
 Blister on dead skin, 48  
     location of, to be noted, 66  
 Blocks of wood as head-rests, 33  
 Blood, abnormal constituents of, 120  
     appearance of, in poisoning, 419  
         cyanid poisoning, 419  
     bile in, 120  
     bilirubin crystals in, 120  
     blackish pigment in, 120  
     character of, in membranes of brain, 231  
     Chareot-Leyden crystals in, 121  
     coagulation of, at postmortem, 115  
     color of, 113, 195  
     cryoscopic index of, 114  
     diabetic, reaction of, 224  
     diseases of, viii, p. 113; 172, 323, 427, 432, 433, 434  
     effect of air on, 119  
     examination of, viii, p. 113; 228, 419, 431, 438, 444  
     freezing-point of, 114  
         in drowning, 417  
     gas-bubbles in, 121  
     glycogen in, 120  
     hæmatoidin crystals in, 120  
     in death by electricity, 413

- Blood in sunstroke, 414  
 lesions of, 113  
 macroscopic changes in, 113  
 method of fixing, 338  
 on body, location of, to be noted, 65  
 organisms in, 121  
 pathologic conditions of, 115  
 poisons in, 121  
 removal of, for bacteriologic examination, 444  
 specific gravity of, 114  
 spectroscopic picture of, 418  
 tests for, 118  
 tumor-cells in, 120  
 uric acid in, in gout, 128
- Blood-blisters, caused by costotome, 31
- Blood-color scale, postmortem, 113
- Blood-diseases, viii, p. 121
- Blood-flukes, 321
- Blood-plates, 122
- Blood-stains, 117
- Blood-tests, 113
- Blood-vessels, diseases of, 18, 20, viii, p. 134; 221, 224, 247, 307, 308, 314, 408, 427  
 examination of, 14, 441, 442, 443  
 tumors of, 139, 140
- Blow-pipe, 32
- Board, Anatomical, 5
- Boards of Health, 4
- Boards to stand on while making post-mortem, 33
- Bodies, frozen, 289, 437  
 intercarotid, 112, 260  
 loose, in joint, 265  
 rice, 261
- Body, see Cadaver.
- Bone, abscesses in, 268  
 age of, how indicated, 58  
 aneurisms of, 274  
 angiomas of, 273  
 ankylosis of, 71, 262, 263, 264, 267  
 atrophy of, 264  
 chondromata of, 273  
 cysts of, 274  
 decalcification of, 345  
 deformities in dwarfism, 56  
 diseases of, 70, 71, 72, 122, 123, 124, 127, 246, xvii, p. 261; 303, 306, 308, 317, 432, 461  
 erosion of, 268  
 examination of, 14, 58, xvii, p. 261; 439, 440  
     in animals, 389  
 fixatives for, 338  
 forceps, 31  
 fractures, 70  
 hæmatoma of, 274  
 inflammation of, 262  
 injuries to, 267  
 lipomata of, 273  
 marrow of, 267  
 measurements of, 57
- Bone, myxoma of, 274  
 necrotic changes in, 267, 268  
 osteomyelitis of, 270  
 parasites of, 273  
 rhachitic, 71  
 sarcoma of, 274  
 syphilis of, 264, 306, 307  
 tuberculosis of, 264  
 tumors of, 273
- Bony formations in muscle, 83
- Borders of organs, 16, 18
- Boric acid as preservative for food, 45  
 toxicology of, 429
- Bothriocephalus latus, 121, 320
- Bovine tuberculosis, 373
- Bowel, see Intestine.
- Bow-leg, 71
- Box, refrigerator, 10
- Brachycephalic skull, 361
- Brain abscess, 246  
 actinomycosis of, 255  
 anemia of, 247  
 angioma of, 255  
 anthrax of, 292  
 cavity, injecting embalming fluid into, 288  
 charts of, 17  
 child's, removal of, 276  
 color of, method of preserving, 21  
 congenital anomalies of, 249  
 diameter of, 363  
 diseases of, 14, 67, 68, 117, 132, 15, 138, 149, 151, 233, xv, p. 246; 290, 291, 299, 302, 307, 308, 313, 411, 432, 433  
 dissection of, in animals, 388  
 embolism of, 16  
 encephalitis of, 250  
 examination of, 14, 402, xii, p. 225  
     Dejerine's method, 238  
     Giacomini's method, 241  
     Hamilton's method, 239  
     Kaiserling's method, 343  
     Meynert's method, 237  
     Pitres's method, 237  
     Virchow's method, 235  
 fibro-endothelioma of, 255  
 fibroma of, 255  
 glioma of, 255  
 gliosarcoma of, 255  
 granulomata of, 255  
 hardening of, 9, 241  
     advantages of, 234  
     with zinc chlorid, 241  
 horizontal section of brain, 237, 240  
 internal examination of, 234  
 length of, 363  
 lipoma of, 255  
 measurements of, 363  
 method of fixation, 338  
 methods of sectioning, 234  
 myxoma of, 255  
 neurofibroma of, 255

- Brain of animals, examination of, 388  
 osteophytes of, 255  
 parasites of, 255  
 pathology of, xv, p. 246  
 perithelioma of, 255  
 psammosarcoma of, 255  
 removal of, 7, 14, 52, 232, 276, 281, 439  
   in animals, 387, 388, 394  
     child, 276  
 sarcoma of, 255  
 sectioning of, 237  
 specific gravity of, 363  
 stem, weight of, 362  
 syphilis of, 255, 307, 308  
 table of weights of, 362  
 time required for hardening, 338  
 tuberculosis of, 311, 313  
 tumors of, 255, 313  
 vertical transverse incision of, 239  
 volume of, 363  
 weighing of, 232  
 weight of, 360
- Bran, in preservation of body, 283
- Break-bone fever, 293
- Breast, abscess of, 61  
 adult, 372  
 atrophy of, 61  
 cancer of, 61  
 diseases of, 61, 81, 301, 317  
 examination of, 61  
 fluid in, 61  
 hypertrophy of, 61  
 infantile, 279  
 infection of, 61  
 lactation of, 372  
 non-puerperal diseases of, 460  
 pathologic conditions of, 81  
 puerperal diseases of, 460  
 shape of, 61  
 size of, 61  
 supernumerary, 61  
 tumors of, 61  
 weight of, at birth, 372  
 wren's, 106
- Breathing, see Respiration.
- Bremer-Williamson reaction of diabetic blood, 224
- Bright's disease, complications of, 458  
 skin eruptions in, 64  
 synonyms of, 458
- Broad ligament, examination of, 14, 15
- Bromatotoxismus, 418
- Bromid, toxicology of, 421, 422
- Bromin, for disinfecting wounds, 34  
 post-mortem wounds, 43  
 toxicology of, 21, 422, 423
- Bromoforn, 21
- Bronchial glands, diseases of, 145, 151, 153, 155, 291, 301, 313, 316
- Bronchi, carcinoma of, 146  
 diseases of, 116, 144, 145, 146, 147, 153, 246, 291, 297, 316, 319, 408, 427, 435
- Bronchi, examination of, 15, 441, 444  
 fistula in, 144  
 foreign bodies in, 146  
 oedema of, 146  
 primary tumors of, 146  
 secondary tumors of, 146  
 stenosis of, 146  
 tumors of, 146
- Bronchiectasis, 144, 145, 150
- Bronchitis, 144  
 acute, 144  
 atrophic, 145  
 capillary, 427  
 catarrhal, 145  
   synonyms of, 455  
 cheesy, 146  
 chronic, 144  
   hypertrophic, 145  
   synonyms of, 456  
 complications of, 448  
 croupous, 145  
 death due to, 408  
 due to gangrenous material in lungs, 150  
 fibrous, 145  
 gangrenous, 146, 150  
 hypertrophic, 145  
 plastic, 145  
 purulent, in typhoid fever, 319  
 putrid, 144, 145  
 subacute, 144  
 suppurative, 145  
 tuberculous, 145
- Bronchopneumonia, 151, 223  
 complications of, 145, 294, 306, 448  
 synonyms of, 456
- Bronchus, fluid in, examination of, 108  
 left, position of, 108  
 right, position of, 108  
 stenosis of, 146
- Bronzed diabetes, 224, 422
- Bronzing of skin, 16
- Brooks, Phillips, 4
- Brown atrophy of heart, 129
- Bruises, location of, to be noted, 65
- Brunetti's chisels, 243
- Bubonic plague, 301, 373
- Bucket method of opening and cleansing intestines, 167
- Buffy angina, see Diphtheria.
- Buhl's disease, icterus neonatorum.
- Bulbs as a means of securing fluids for bacteriologic study, 347
- Bulk, determined by comparison, 18  
 measurement, 18  
 Virchow's method, 18
- Bullous emphysema of lungs, 150
- Burials, premature, 46
- Burns, 412  
 effects of, 412  
 forms of, 463  
 from corrosive substances, 463
- Bursæ, 272



Bursitis, 273  
 Butter yellow, toxicology of, 422  
 Buttonhole mitral, 133

## C

- Cachectic subjects, post-mortem rigidity of, 51  
 Cachexia, changes of subpericardial fat in, 101  
   malarial, 322, 323  
   pigmentation in, 112  
   purpuric, 126  
 Cadaver, bacteria in, 207  
   breathing, absence of, 47  
   closure of, after postmortem, 445  
   color of skin of, 60  
   contact with surrounding objects, 55, 416  
   cupping of, 47  
   disembowelling of, 15, 276  
   elevation of temperature of, 7  
   embalming of, 286  
   examination of exterior of, v. p. 46  
   examination of, 437  
   exterior of, examination of, v. p. 46  
   frozen, 289  
   identification of, 53  
   lividity of, 49  
   measurements of, 53, 50, 57, 358, 444  
   pecuniary value of, 4  
   place of finding, 55  
   preservation of, xx, p. 286  
   portions of, 5  
   removal of, 54  
   restoration of, xx, p. 283  
   right to dispose of, by will, 4  
   scarification of, 47  
   subcutaneous injections into, 47  
   surroundings of, 55, 416  
   thawing of, 289  
   weighing of, 10  
   weight of, 54, 358, 360, 444  
 Cecum, diseases of, 308, 312  
   examination of, 15, 100  
   measurements of, 300  
   removal of, 15, 105  
 Carsson disease, 220  
 Calcareous deposits, 120, 272  
   infarcts in kidneys, 200  
   infiltration, 120  
   of arteries, 135  
 Calciosis, 135  
 Calcification in tuberculosis, 312  
 Calculated bodies in abdominal cavity, 20  
 Calculi, biliary, see Gall-stones.  
   hydronephrotic, 200  
   in pancreas, 100  
   metamorphosed, 207  
   renal, 200  
   salivary, 200  
   vesical, 200  
 Calculous pyelitis, 203, 204  
 Calipers, graduated, 32  
 Calomel, toxicology of, 148, 424  
 Calvarium, clamps for holding, 32  
   diseases of, 230  
   examination of, 229, 230, 439  
   method of loosening, 227  
   methods of removing, 226, 408  
     in animals, 387, 391  
 Campecia wood, effect on urine, 425  
 Camphor, toxicology of, 421  
 Canal of Wirsung, opening of, 188  
   spinal, xiv, p. 242  
 Cancer, see Carcinoma.  
 Cancerous stomach, removal of, during life, 190  
 Cannabinone, toxicology of, 421  
 Cannabis indica, toxicology of, 421  
 Cannula, 32  
 Cantharide, potassium, effect on urine, 426  
 Cantharides, toxicology of, 203  
 Cantharidinate, potassium, effect on urine, 424  
 Cantharidine, toxicology of, 423, 424, 426  
 Capsule of kidney, examination of, 178  
   liver, 195  
   organs, 16, 18  
   Tenon, 257  
 Caput medusæ, 140  
 Carbohydrate metabolism, 222  
 Carbol-fuchsin for staining tubercle bacillus, 310  
 Carbolic acid in treatment of post-mortem wounds, 43  
   toxicology of, 411, 421, 422, 425, 426  
 Carbonates, alkaline, effect on urine, 42  
 Carbon bisulphid, toxicology of, 421, 42  
   dissolved as a freezing agent, 12  
   toxicology of, 42, 415, 421  
 Carbonic acid, toxicology of, 421  
 Carbon monoxid, toxicology of, 419, 42, 425, 424  
 Carbonyl chlorid, toxicology of, 422  
 Carcinoma, 408, 451  
   bodies, fixation of, 357  
   colloid, of stomach, 100  
   of bladder, 205  
   breast, 451  
   bronchi, 140  
   choroid, 140  
   ciliary body, 75  
   cisterns, 60  
   heart, 134  
   intestines, 171  
   kidney, 204  
   larynx, 143  
   liver, 100, 214, 215, 220  
   lungs, 155  
   lymph-vessels, 140  
   mediastinum, 140

- Carcinoma of nasal passages, 141**  
     navel, 66  
     œsophagus, 190  
     penis, 68  
     pericardium, 99  
     peritoneum, 160  
     pleura, 156  
     spleen, 164  
     stomach, 190  
     vulva, 69  
     sarcomatodes of kidney, 204  
**Cardiac plexus, dissection of, 276**  
**Cardol, toxicology of, 422**  
**Care of hands, iv, p. 38**  
     clothes removed from body, 54  
**Caries, 268**  
     of orbit, 76  
**Carmin stains, 333**  
**Carnification of lung, 153**  
**Carnoy's fluids, as fixing fluids, 330**  
**Carotid arteries, 80, 112, 138, 232, 304, 442**  
     bodies, 112, 260  
     tumors of, 112  
**Cartilage, degenerations of, 264**  
     fixatives for, 338  
**Caruncle of urethra, 70**  
**Cascarilla, to remove odor, 10**  
**Caseous nodules in tuberculosis, 316**  
     pneumonia, 311  
     tuberculosis, 311  
**Catarrh, acute nasal, 141**  
     atrophic, 141  
     chronic nasal, 141  
         adenoids in, 141  
         fibrinous, 141  
         hypertrophic,  
         membranous, 141  
**Catarrhal and croupous pneumonia, table**  
     showing difference between, 154  
     appendicitis, 171, 172  
     bronchitis, 145  
     dysentery, 173  
     laryngitis, 142  
     pneumonia in typhoid fever, 318  
     pyelitis, 203  
     synovitis, 262  
**Catheter, metal, 32**  
**Cat, postmortem of, 394**  
**Causes of death, 16, xxviii, p. 448**  
     complications of, xxviii, p. 448  
     in 799 coroner's cases, 411  
     nomenclature of, xxviii, p. 448  
     synonyms of, xxviii, p. 448  
     usual, xxviii, p. 448  
**Caustic salts, toxicology of, 423, 427**  
     soda for absorption of oxygen, 353  
**Caustics, effect of, on dead skin, 48**  
     toxicology of, 423  
     used in treating post-mortem wounds,  
     43  
**Cautery, for post-mortem wounds, 43**  
**Cavity, abdominal, exposure of, vi, p. 79**  
     of animals, 375  
     abdominal, exposure of, in dog,  
     393  
         topographic examination of, vi,  
         p. 79  
     cranial, of animals, 386  
         of cow, after removal of bony  
         vault, 390  
     embalming, 288  
     oral, of animals, 384  
     thoracic, critical examination of, vii,  
     p. 92  
         exposure of, vii, p. 92  
         of animals, 376  
**Cells, ganglion, fixatives for, 339**  
**Cellular tissue, diseases of, 461**  
**Cellulitis, axillary, treatment of, 44**  
     of hand, treatment of, 45  
     orbit, 258  
**Central lesion of brain, 110, 240, 302**  
     nervous system, actions of poisons on,  
     419  
         fixatives for, 338  
**Centre governing development, 66**  
**Centrum ovale, methods of studying, 235,**  
**237**  
**Cephæline, toxicology of, 423**  
**Cephalanthine, effect on urine, 425**  
**Cephalic cavities of a horse, lines for saw-**  
**ing, 387**  
     inspection of, xiii, p. 225  
     method of opening, 387  
     index, 360  
         types distinguished by, 361  
**Cephalonic skull, 361**  
**Cerebellum, average weight of, 362**  
     removal of, 236  
     tuberculosis of, 313  
**Cerebral arteries, aneurism of, 247**  
     hemorrhage, 250  
         as cause of death, 408  
     hyperæmia, 251  
     softening, complications of, 453  
**Cerebrospinal leptomenigitis, acute, 251**  
     menigitis, acute, 252  
**Cerebrum, average weight of, 362**  
**Cervical organs in animals, dissection of,**  
**384**  
     ribs, 68  
**Cervico-thoracic abdominal cavity, exami-**  
**nation of, 275**  
**Cervix uteri, malignant neoplasms of, 211**  
**Cestodes, 320**  
     intestinal, 320  
**Chain hooks, 32**  
     saw, 259  
**Chalky deposits in gout, 127**  
**Chancre, 307**  
     of vulva, 69  
**Chancroid, 351**  
     of vulva, 69  
**Chantemasse, hacillus of, 172**  
**Charcot-Leyden crystals, 121, 144**  
**Charcot's joint, 71, 248**

- Charts, outline, how used, 17  
 English gummed outline, 17  
 Cheek, sucking gland in, 66  
 Cheeks, fissures, 68  
 Cheesy bronchitis, 146  
 Cheiragra, 263  
 Cheiranthus, toxicology of, 424  
 Chemic, inorganic poisons, 420  
 organic poisons, 421  
 poisoning, 16, 420, 421  
 Chemicals detected by odors coming from  
 body, 418  
 Chenopodium, effect on urine, 425  
 Chiara's raspatory, 31  
 Chicken-fat clots, 115  
 Chicken-pox, 351  
 Child, full-term, healthy, dimensions of,  
 359  
 in postmortem, whether it is viable,  
 403, 444  
 preservation of body of, 275  
 Chisels, 31  
 guarded, 31  
 hatchet-shaped, 31  
 other kinds of, 31  
 spinal, 31  
 straight, 31  
 T-shaped, 31  
 Chloral hydrate, effect often difficult to  
 determine, 429  
 toxicology of, 421, 422, 425, 429  
 Chlorate, potassium, toxicology of, 203,  
 423, 425  
 Chlorin, toxicology of, 21, 422  
 Chloroform and ether poisoning, deter-  
 mined by odor, 430  
 fright as an element in,  
 429  
 in evidence, but death  
 due to other causes, 430  
 toxicology of, 21, 411, 421,  
 425, 429, 430  
 and its derivatives, toxicology of, 421  
 narcosis, death from, 198  
 Chlorosis, 122, 123, 452  
 blood changes in, 122  
 due to hypoplasia of aorta, 136  
 Chlorphenol, odor of, 21  
 toxicology of, 21  
 Cholangitis, suppurative, 219  
 Cholecystitis, acute infectious, 215  
 Cholelithiasis, 215, 220  
 Cholera Asiatica, 292  
 appearance of stools in, 293  
 coeliac ganglion in, 293  
 heart in, 293  
 kidney in, 293  
 liver in, 293  
 lungs in, 293  
 mucous membrane in, 293  
 odor in, 170  
 post-mortem rigidity of abdomi-  
 nal muscles in, 51  
 Cholera morbus, 449  
 nostras, synonyms of, 449  
 Chondritis, 265  
 Chondroma, due to misplaced carti-  
 nous tissues, 273  
 of lungs, 155  
 nasal passages, 141  
 pleura, 156  
 vulva, 69  
 Chordæ tendineæ, 129  
 Choreia, 249, 351, 453  
 Choroiditis, syphilitic, 306  
 Choroid plexus, 236, 439  
 Christmas-tree, needles from, as evidence,  
 55  
 Chromates, toxicology of, 422, 423, 425  
 Chromatolysis of cells of central nervous  
 system in uræmia, 117  
 Chromic acid as a fixing fluid, 330  
 toxicology of, 423  
 Chromo-acetic acid, as a fixing fluid, 331  
 Chromo-aceto-osmic acid, as a fixing fluid,  
 331  
 Chromo-formic acid, as a fixing fluid, 331  
 Chromo-nitric acid, as a fixing fluid, 331  
 Chromo-osmic acid, as a fixing fluid, 331  
 Chrysarobin, effect on urine, 422, 425  
 Chrysoidin, toxicology of, 422  
 Chyle duct, examination of, 197  
 Chylothorax, 157  
 Cicutoxin, toxicology of, 421  
 Cimex lectularius, 319  
 Cinchona bark, toxicology of, 422  
 Circular method of removing calvarium,  
 226  
 Circulation, tests for, 47, 48  
 Circulatory apparatus, diseases of, 454  
 disturbances of kidney, 200  
 lungs, 148  
 system, syphilis of, 307, 308  
 tuberculosis of, 314  
 Cirrhosis, atrophic, of liver, 216  
 Hanot's hypertrophic, 115  
 of liver, 216, 224  
 complications of, 458  
 due to alcoholism, 427  
 Cirroid aneurism, 137  
 of vein, 140  
 Clamps for holding calvarium, 32  
 Classification, Bertillon, of diseases, xxviii  
 p. 448  
 of poisons, 420  
 Clavicle and ribs, protection from injury  
 from, 94  
 Cleft scrotum, 69  
 urethra, 69  
 Clinoccephalic skull, 361  
 Clitoris, absence of, 69  
 atrophy of, 69  
 carcinoma of, 69  
 hypertrophy of, 69  
 Closure of body after postmortem, xx  
 p. 283



- Clothing, bullet-holes in, 54  
 care of, 54  
 examination of, 54  
 receipt for delivery of, 54  
 stains on, 54
- Clots, chicken-fat, 115  
 currant-jelly, 115  
 fibrous, of aneurism, 138  
 of pernicious anæmia distinguished from those of cancer, 122
- Clubbed fingers, 68, 269  
 foot, 269
- Coal dust in hands, 16  
 lungs, 16  
 spleen, 163
- Cocaine, toxicology of, 421, 422, 425, 426, 430
- Codeine, toxicology of, 422
- Code of Hammurabi, 1
- Coeliac ganglion, appearance of, in cholera Asiatica, 181, 293  
 hemorrhage of, 293  
 situation of, 181
- Coffee, burning of, to remove odor at postmortem, 10
- Colchicine, toxicology of, 421, 422, 423
- Cold, death from, 414
- Colitis, 174  
 chronic, 175  
 croupous, 153  
 membranous, 174  
 simple, 174  
 ulcerative, 175
- Collection of material, 5, 326, 346
- Colloid cancer of lungs, 156  
 stomach, 190  
 material, formalin of Orth for fixing, 340
- Coloboma, 75
- Colocynthis, toxicology of, 423
- Colon, dilatation of, 175  
 diseases of, 153, 160, 172, 173, 174, 175, 202, 312  
 examination of, 15, 89, 165  
 malignant disease of, 175  
 measurements of, 366  
 of animals, removal of, 379  
 removal of, 15, 165  
 tumors of, 171, 175
- Color changed by washing, 19  
 of eyes, peculiarities of, 74  
 racial peculiarities of, 74  
 feces, 170  
 kidney after removal of capsule, 178  
 organs, 16, 18, 109, 195  
 values by kromskop, 18
- Colors, toxicology of anilin, 422, 426
- Colostrum, examination of, 81
- Coma, fatal, 21
- Comma bacillus of Koch, 292
- Common bile duct, see Ductus choledochus communis.
- Comparative postmortems, xxv, p. 373
- Compensation in practice of medicine, 398
- Complications of causes of death, xxviii, p. 448
- Compound fractures, 266
- Concentric atrophy of bone, 264  
 hypertrophy of heart, 106, 133
- Conception, performing on cadaver operation for prevention of, 3
- Concretions in kidneys, 200
- Condyloma of urethra, 70
- Cones, graduated, 33
- Congenital abnormalities of uterus, 209  
 aneurism, 139  
 anomalies, 17, 19, 249  
 cystic kidneys, 200  
 defects in kidney, 199  
 deformities, 66  
 dislocation of hip, 266  
 diverticula of duodenum, 188  
 luxation of sternum, 266  
 redundancy of sigmoid flexure, 160  
 rickets, 272  
 syphilis, 307  
 as cause of death, 408  
 syphilitic pancreatitis, 221
- Congestion, hypostatic, 49, 149  
 of kidney, 199  
 liver, 218  
 lungs, 149  
 passive, of lungs, 148
- Coniine, toxicology of, 422
- Conium, toxicology of, 421
- Conjunctiva, 19, 74  
 palpebral, hemorrhages of, 132  
 in endocarditis, 132  
 post-mortem changes in, 48
- Conjunctivitis, 74, 153  
 gonorrhæal, 297
- Connective tissue, fixation of, 340
- Considerations, general, i, p. 1
- Consistency of organs, how affected and how determined, 17, 19, 195
- Constipation, 169
- Contagious diseases derived from domestic animals, 373
- Contents, abdominal, in animals, dissection of, 383  
 removal of, of a dog, 393  
 in swine, 391  
 of cavities, 20  
 thoracic, removal of, in animals, 384
- Continued fever, see Typhoid fever.
- Contour of organs, 14, 18
- Contraction of bowels, 165  
 pupils after death, 48  
 scrotum, 69
- Contracts between physician and patient, 397
- Contractures, 71, 269  
 of muscles, disappearance of post-mortem, 71
- Contrecoup, 230

- Convallaria, toxicology of, 424  
 Conviction of murder by poisoning, 420  
 Convulsions, 448, 453  
 Copaiba, balsam of, toxicology of, 422  
 Copper sulphate, purification of drinking water by, 430  
   toxicology of, 423, 430  
 Cord, diseases of, xv, p. 246  
   examination of, xiv, p. 242  
   opening of, 244  
   preservation of, 244  
   removal of, 242, 277  
     in animals, 389  
   spinal, see also Spinal cord.  
   syphilis of, 307, 308  
   tuberculosis of, 313  
   tumors of, 255  
   weight and dimensions of, 364  
 Cords, vocal, examination of, 111  
 Cornea, appearance of, after death, 257  
   post-mortem changes in, 48, 74  
 Corneal abscess, 74  
 Cornell head-rest, 225  
 Cornutine, toxicology of, 421, 423  
 Coronary arteries, 105, 129, 131, 135, 308  
   obstruction of, cause of death, 408  
   "pipe-stem" character of, 101  
 Coroner, abolition of, 1  
   authority of, 1, 5  
   creation of, 1  
   duties of, 1  
   office of, establishment of, 1  
 Coroner's cases, 1, 411  
 Coronilla, toxicology of, 424  
 Corpora Arantii, 129  
   quadrigenina, 237, 439  
 Corpse, see Cadaver.  
 Corpulency, causes for, 50  
*Corpus delicti*, 55, 58, 401  
 Corpus striatum, examination of, 439  
 Corrosion, by poison, 426  
 Corrosive acids, toxicology of, 421, 423  
   alkalies, effect on urine, 424  
   toxicology of, 421  
   metallic salts, toxicology of, 423  
   poisons, toxicology of, 423, 425  
   sublimate, 418  
     for fixing tissues, 333  
 Cortex of kidney, normal relation to medulla, 179  
 Cortical lesion, 237  
 Cor villosum, 99  
 Corynebacterium diphtheriæ, 151, 350  
   mallei, 291, 350  
   xerosis, 350  
 Coryza, 141  
 Costotome, 31, 93  
 Cotton, absorbent, in the preservation of the body, 283  
   plugs, 352, 353  
   wool, 34  
 Court and expert, 399  
   testimony in, 17  
 Court, weights and measures used in, 17  
 Cover-slips, preparation of, 348  
 Cow-pox, 351  
 Cranial cavity, examination of, xiii, 43  
   in animals, exenteration of, 38  
   of cow, after removal of base vault, 390  
   dog, exposure of, 393  
   horse, lines for opening, 38  
   postmortem of, xiii, p. 225; 43  
 Craniopagus, 67  
 Cranioschisis, 249  
 Craniotabes, 72  
 Cranium, dimensions of, 363  
   examination of, xiii, p. 225; 439  
   sectioning of, in animals, 386  
   vault of, substitute for, 284  
 Credé's ointment, 44  
 Creosote, toxicology of, 21, 411, 425  
 Crepitation of lungs, how elicited, 109  
 Cresol, toxicology of, 425  
 Cretinism, 96, 249  
 Cricothyroid joint, ankylosis of, 143  
 Criminal abortion, 406  
 Critical examination of organs of abdominal cavity, x, p. 159  
   thoracic cavity, vii, p. 92  
 Croton oil, toxicology of, 422, 423, 424  
 Croup, 293, 448  
 Croupous and catarrhal pneumonia, table showing differences between, 154  
   bronchitis, 145  
   colitis, 153  
   gastritis, 153  
   pneumonia, 151, 152, 154, 223  
     gray hepatization in, 153  
     hyperæmia in, 152  
     red hepatization in, 152, 153  
 Crura of diaphragm, severing of, 236  
 Crushing injuries to joints, 267  
   suicide by, 463  
 Cryer's electrical surgical engine, 30  
 Cryoscopic index, 114  
 Cryoscopy, 114, 416  
 Crystalline lens, 75  
 Crystals, Charcot-Leyden, 121  
   hæmatoidin, 120  
 Csoker's method of removing thoracic and abdominal contents, in animals, 382  
 Cubebene, toxicology of, 422  
 Cultivation of organisms, 352  
 Culture-media, inoculating, 351  
   list of, 350, 351  
   selection of, 349  
 Cultures of anthrax, 356  
   preparation of, for inoculation, 353  
     from solid organs, 352  
   smear, 352  
   stab, 352  
   stroke, 352  
   used in inoculation, 353  
 Cupping on dead subject, 47  
 Curarine, toxicology of, 422, 425

Curcas oil, toxicology of, 422  
 Currant-jelly clot, 115  
 Curschmann's spirals, 144  
 Curvatures of iliac arteries, 20  
   spinal, 269  
 Cutaneous diphtheria, 294  
   eruptions in syphilis, 307  
   form of tuberculosis, 318  
 Cutting instruments, suicide by, 462  
 Cyanid of cacodyl, toxicology of, 428  
   potassium, toxicology of, 411, 419, 430  
 Cyanosis, 133, 216  
   chronic, 128  
   of skin in cholera Asiatica, 293  
 Cyanotic cirrhosis of liver, 217  
   induration, 133  
 Cyclamine, toxicology of, 425  
 Cyclocephalus, 67  
 Cylindrical aneurism, 137  
 Cystadenoma, papillary, 146  
 Cystic disease of kidney, 200  
   duct, 188, 383  
   formation in prostate, 212  
   growths in nasal passages, 141  
   kidney, incising of, 178  
 Cysticercus cellulosæ, 151, 320  
   of bladder, 207  
     bone, 274  
     brain, 255  
     heart, 134  
     kidney, 202  
     pericardium, 99  
 Cystinuria, 424  
 Cystitis, 205  
   diphtheritic, 205  
   due to alcohol, 427  
     colon bacillus, 205  
   effect on urine, 424  
 Cysts, dermoid, in mediastinum, 146  
   occlusion, 66  
   of bones, 273  
     Fallopian tube, 208  
     iris, 75  
     kidney, 200  
     nasal passages, 141  
     ovary, 209, 459  
     pancreas, 196  
     peritoneum, 159  
     ureters, 204  
 Cytisine, toxicology of, 421, 432  
 Cytoalbumin increased in hæmophilia, 125  
 Cytology, 417  
 Cytolysins, 421

D

Dactylitis, syphilitic, 306  
 Daylight, substitute for, at postmortem, 8  
 Dead body, see Cadaver.  
 Dead-house, arrangement of, 10  
   library for, 12  
   museum for, 12

Death, ascertainment of cause of, 16  
   by cold, 414  
     drowning, 415  
     electricity, 413  
     hanging, 415  
     heat, 414  
     poisons, xxvi, p. 417  
     starvation, 414  
     suffocation, 415  
     throttling, 415  
     traumatism with but slight external injury, 409  
     violence, xxvi  
   cause of, 16, xxviii, p. 448  
   causes of, in 799 coroner's cases, 411  
   circumstances of, 53, 437  
   complications of causes of, xxviii, p. 448  
   cooling of body after, 49  
   due to bolus of food in larynx, 143  
     piece of lead-pencil in larynx, 143  
   ill-defined or unspecified causes of, 464  
   in cases of abortion, usual causes of, 407  
   mask, 289  
   microscopic changes in, 413  
   molecular, 326  
   nomenclature of causes of, xxviii, p. 448  
   of illegitimate child, concealment of, 55  
   pupils in, 46, 413  
   rattle, 46  
   signs of, 46  
     cessation of all tissue vitality, 46  
     complete loss of vital warmth, 48  
     continuous cessation of circulation, 47  
       respiration, 47  
     earlier, 46  
     inability to move, 46  
     insensibility, 46  
     loss of muscular irritability, 47  
       nervous irritability, 47  
       reflexes, 46  
     negative, 46  
     positive, 47  
     putrefaction, most positive, 49  
     vary under different conditions, 46  
   spermatozoa motile after, 46  
   stiffening or post-mortem rigidity, 50  
   sudden, due to anæmic infarct of heart, 130  
     aortic stenosis, 133  
     in infants from pressure on thymus, 96  
   synonyms of causes of, xxviii, p. 448  
   three proximate causes of, 21  
   time after, to make postmortem, 7, 436



- Death, usual causes of, 407, xxviii, p. 448  
 violent, 403, 408
- Decapsulation of kidney, 3, 179
- Decomposition, 50, 52  
 delayed, 52  
 discoloration of, 52  
   Hofmann's treatment of, 52  
 greenish color of, 50  
 post-mortem, 52  
 rapid, 52, 304, 414
- Deformities, 66, 71  
 acquired, 66, 72  
 asymmetrical, 66  
 congenital, 66  
 discovered by comparison, 71  
 diseases associated with, 71, 72  
 due to dislocations, 71  
   embryologic defects, 66  
   fractures, 71  
   injury at birth, 66  
   pathologic intra-uterine causes, 66  
   traumatism, 66  
 laryngeal, 142  
 multiple, 66  
 muscular, 72  
 orthopædic, 268  
 pathologic, 71  
 single, 66  
 symmetrical, 66
- Degenerated nerve-fibres, fixatives for, 339
- Degeneration, 129  
 albuminous, fixatives for, 338  
 amyloid, 129, 135, 418  
   fixatives for, 338  
   of liver, 214  
 cardiac, 129  
 fatty, 129, 218, 223, 418  
   of arteries, 135  
   heart, 129  
   liver, 218  
 hyaline, 129  
   of arteries, 136  
   pancreas, 222  
 lardaceous, 144  
 myocardial, 129  
 pancreatic, 196  
 signs of, 70
- Dejerine's method of examining brain, 238
- Delirium, acute, 250
- Dementia, paralytic, 248
- Dengue, 293, 373
- Dental engines, 29  
 tumors, 78
- Deodorants, 34
- Dermatomyositis, 63
- Dermoid cysts of lungs, 155  
 mediastinum, 146  
 orbit, 75  
 ovary, 209  
 uterus, 209
- Dermopathies, 248
- Determination of viability of child, 403, 444
- Development, stunted, cause of, 70
- De Vilbiss's electrical surgical engine, 30
- Dextrocardia, 113
- Diabetes, bronzed, 224, 422  
 complications of, 452  
 insipidus, 127  
   urinary changes in, 127  
 islands of Langerhans in, 223, 224  
 mellitus, 222, 223, 224, 452
- Diabetic blood, reaction of, 224
- Diacetic acid, 222
- Diagnosis, hypothetical or tentative, 16
- Diaphanous test, 47
- Diaphragm, cause of depression of, 85  
 elevation of, 85  
 diseases of, 148, 149, 197  
 examination of, 13, 14, 15, 85, 197, 406, 440, 444  
   in animals, 376, 391  
 height and location of, 17, 84, 85  
 removal of, 275, 444  
   in animals, 381
- Diaphragmatic pleurisy, 158
- Diarrhœa, summer, cause of, 172
- Diastasis, 266
- Dicephalus, 67
- Differentiation, optical, in preservation of tissues, 329
- Digestion, post-mortem, 87
- Digestive apparatus, diseases of, x, p. 150: 458
- Digitalis, toxicology of, 101, 421, 423, 424
- Dilatation of colon, 175  
 heart, 133  
 lymph-vessels, 140  
 pupils after death, 48  
 right heart, 144  
 stomach, 190
- Dimensions, xxiv, p. 357  
 how expressed, 17, 357  
 of adrenals, 369  
   babes, 358, 359  
   bladder, 368  
   children, 358  
   cranium, 363  
   embryos, 359  
   gall-bladder, 367  
   hair, 359  
   heart, 364  
   kidney, 368  
   large intestine, 366  
   liver, 367  
   lungs, 366  
   men, 358  
   œsophagus, 366  
   orifices of the heart, 365  
   ovaries, 371  
   pancreas, 369  
   pituitary gland, 363  
   prostate, 372  
   seminal vesicles, 372

- Dimensions of skull, 360  
 small intestine, 366  
 spinal cord, 364  
 spleen, 369  
 stomach, 366  
 suprarenals, 369  
 testes, 370  
 thymus gland, 370  
 thyroid gland, 370  
 umbilical cord, 359  
 ureters, 368  
 urethra, 368  
 uterus, 371  
 vermiform appendix, 366  
 vesicles, seminal, 372  
 women, 358
- Dioxid, nitrogen, toxicology of, 422
- Diphtheria, 293, 448  
 bacillus of, 151  
 complications of, 448, 449  
 corynebacterium of, Plate V, No. 6.  
 p. 350  
 cutaneous, 294  
 endometritic, 210  
 false membrane of, 294  
 gastritic, 191  
 growths associated with, 294  
 laryngeal, 294  
 mucous membranes in, 295  
 nasal, 294  
 of bladder, 206  
 vulva, 69  
 pharyngeal, 294  
 pseudobacillus of, 293  
 retropharyngeal abscess in, 294
- Diplococcus erysipclatis, 295  
 pneumoniz, 151, 152  
 varieties of, 349
- Diploe in animals, 390
- Diprosopus, 67
- Dipygus, 67  
 parasiticus, 67
- Directors, grooved and curved, 32
- Dissection means division of.  
 of abdominal contents of animals, 383
- Discoloration, by poison, 426, 427
- Disease, Addison's, 16  
 Barlow's, 126  
 Basedow's, 97, 147  
 Duke's, 300  
 Glenard's, splanchnotosis or abdominal ptosis.  
 Graves's, 97, 147  
 Hodgkin's, 124  
 von Jaksch's, 124  
 of various organs and parts will be  
 found indexed under their names.  
 Osler's, 128  
 Paget's, of nipple, 324  
 Raynaud's, 254  
 von Recklinghausen's, 254
- Diseases, Bertillon classification of, xxviii,  
 p. 448
- Diseases, blood, 121  
 contagious, and domestic animals, 373  
 contracted at postmortems, 41  
 diagnosed by agglutinative reaction,  
 115  
 due to hæmatozoa, xxi, p. 322  
 micro-organisms, xxi, p. 290  
 parasites, xxi, p. 319  
 specific organisms or proto-  
 zoa not yet isolated, 351  
 echinococcic, 322  
 epidemic, 449  
 following post-mortem wounds, 41  
 general, 452  
 hereditary, as cause of death, 408  
 mortal, cause of sudden death, 407  
 non-puerperal, of the breast, 460  
 non-venereal, of the genital organs of  
 the male, 459
- Disinfectants, 34
- Disinfection of hands after postmortem,  
 40
- Dislocations, 20, 71, 263, 266, 286  
 congenital, 71  
 of hip, 66, 266  
 of hyoid bone, 71  
 neck, 66  
 semilunar cartilages, 266  
 pathologic, 71  
 source of deformity, 71  
 spontaneous, 71  
 traumatic, 71
- Dissecting aneurism, 138  
 wounds, 41
- Dissection of brain in animals, 388  
 cardiac plexus, 276  
 cervical organs, 384  
 heart in animals, 385  
 lungs in animals, 385  
 œsophagus in animals, 384  
 thoracic organs in animals, 384  
 trachea in animals, 385
- Disseminated fat necrosis, 221, 222  
 sclerosis, 254
- Distention of bowels, 165
- Distoma hæmatobium, 151, 202, 207, 220  
 hepaticum, 159  
 Westermanii, 151
- Distomiasis, 321
- Distribution of tubercles in body, 311
- Diverticula, of bladder, 206  
 œsophagus, 190
- Dogs, postmortems on, 392
- Dolichocephalic skull, 361
- Dothienenteritis, see Typhoid fever.
- Double hearts, 113  
 monsters, 67  
 parasites, 67  
 uterus, 209
- Dourine, 324
- Drains, 11
- Drawing up of protocol of postmortem,  
 445

- "Drill" bones, 83  
 Dropsy, chronic articular, 262  
   general, 156, 463  
 Drowning, 415, 463  
   appearance of skin after, 66  
   freezing-point of blood in, 417  
   in salt and fresh water, difference of  
   appearance in, 416  
   suicide by, 462  
 Ducts, liver, diseases of, xii, p. 213  
   method of examining, 188  
   pancreatic, diseases of, xii, p. 220  
 Ductus arteriosus (Botalli), 102, 276, 404  
   choledochus communis, 15, 187, 188,  
   194, 195, 196, 215, 216  
 Duke's fourth disease, 300  
 Duodenal ulcer, 165  
 Duodenum and its ducts, 187  
   diseases of, 196, 220, 221, 428  
   examination of, 14, 15, 187, 442, 443,  
   444  
   in animals, 383  
   length of, 187, 266  
   removal of, 15, 187, 443  
   in animals, 379  
 Dupuytren's contracture, 269  
 Dura mater, adherent, loosening of, 227  
   diseases of, 227, 230, 233, 244  
   examination of, 230, 233, 439, 440  
   French method of opening, 228  
   hemorrhage in, 230, 233  
   in animals, incising of, 387  
   method of dividing, 231  
   of spinal cord, 243  
   removal of, 230, 440  
   in animals, 387  
   tumors in, 230  
 Dura-tongs, 32  
 Dusts, irritating, toxicology of, 422  
 Duties of obducent during postmortem,  
   437  
 Dwarf, microsome, 70  
 Dwarfism, 56  
 Dysentery, 172  
   acute, 172  
   catarrhal, 173  
   amoebic, 173  
   chronic, 172, 173  
   complications in, 174  
   epidemic, 449  
   gangrenous, 173  
   malignant diphtheritic, 173  
   odor in, 170  
   synonyms of, 449
- E**
- Ears, absence of helix, 68  
   diseases of, 67, 70, 73, 128, 246, 252,  
   253, 295, 301, 304, 306  
   examination of, 14, 70, 233, xvi, p.  
   258: 405, 439, 444  
   hæmatoma of, 16, 68  
   Ears, middle, examination of, 158  
   tumors of, 16  
 Eccentric atrophy of bone, 264  
   hypertrophy of heart, 106, 133  
 Ecchymoses, 101, 171  
 Echinococci, 322, 373  
   of bladder, 207  
   bone, 274  
   brain, 255  
   heart, 134  
   kidney, 202  
   liver, 220  
   navel, 66  
   pericardium, 99  
   spleen, 164  
 Eclampsia, 278, 279, 448, 453, 460  
 Ectopia of testicles, 69  
 Ectopic pregnancy, 207  
 Ectromelus, 67  
 Eczema of vulva, 69  
 Elastic fibres, agents for fixing, 340  
   tissue in arteriosclerosis, 134  
   of spleen, 163  
 Electric contractility in body after death,  
   47  
   disturbances, 463  
   fan, 11  
   light over post-mortem table, 11  
   öse, 34  
 Electricity, death by, 413  
   appearance of blood in those  
   dying from, 413  
   heart in, 413  
   pupils in, 413  
   entrance of current in, 65  
   indications of, 413  
   microscopic changes in, 413  
 Electrocutation, 7, 413  
 Electrolysis, aneurism treated by, 139  
 Elephantiasis, 69  
 Elongation of penis, 68  
 Emaciation, 59, 414  
   in atrophic cirrhosis, 57  
   cancer of upper digestive tract, 59  
   muscular atrophy, 59  
   phthisis, 59  
 Embalmed bodies, 286  
 Embalming, arterial, 288  
   cavity, 286  
   fluid, 9  
   effect of, on muscles, 82  
   method of injecting into abdomen,  
   286  
   brain cavity, 287  
   thorax, 287  
   Nauwerck's, 287  
 Emboli of carotid, 112  
 Embolism, 112, 132, 137, 151, 246, 408, 455  
   air, of lungs, 148  
   fat, of lungs, 148  
   of brain, 16, 246, 250  
   pulmonary, 102, 148  
   in chlorosis, 123



- Embryos, fixation of, 337
  - method of determining age of, from length of, 359
- Emetine, toxicology of, 423
- Emphysema, atrophic, of lungs, 150
  - bullous, of lungs, 150
  - local, complicating croupous bronchitis, 145
  - of liver, 218
    - lungs, 149, 150, 151
  - post-mortem, 95
  - vesicular, of lungs, 150
- Empyema, frontal, 258
  - of antrum of Highmore, 141
    - joint, 261
    - pleura, 156
- Encephalitis, 250, 452
- Encephalocele, 68, 249
- Encephaloid cancer of lungs, 156
- Encephalon, maximum weight of, 361
  - relative weight of, to body, 363
- Encysted pleurisy, 158
- Endarteritis, acute, 136
  - chronic, 136
  - obliterans, 135
  - obstructive, 136
  - proliferative, 136
- Endemic hepatitis, 219
- Endocardial vegetations of arch of aorta, 100
- Endocarditis, 131
  - benign, 132
  - cause of hemorrhage into palpebral conjunctiva, 132
  - complications of, 153, 210, 303, 304, 448
  - diphtheritic, 132
  - fetal, 132
  - fibrous, 132
  - gonorrhœal, 207
  - malignant, 16, 132
  - metastatic inflammations associated with, 132
  - mycotic, 132
  - organisms of, 132
  - rheumatic, 132
  - septic, 132
  - syphilitic, 132
  - ulcerative, 132
  - ulcerosa in erysipelas, 295
  - varicose, 132
  - vegetative, 132
- Endocardium, diseases of, 131, 132
- Endocyma, 68
- Endometritis, acute, 209
  - as a cause of spontaneous abortion, 407
  - atrophic, 210
  - chronic hypertrophic, 210
  - diphtheritic, 210
  - gangrenous, 210
  - hemorrhagic, 210
  - tuberculous, 210
- Endophlebitis, fibrous, 140
- Endothelioma of lungs, 156
  - peritoneum, 159
  - pleura, 156
- Engines, dental and trephining, 29
- Enlargement, see Hypertrophy.
- Enophthalmus, 258
- Enostosis of bone, 273
- Ensiform appendix, hook-shaped, 94
- Enteric fever, see Typhoid fever.
- Enteritis and diarrhœa, synonyms of, 457
  - as cause of death, 408
  - nodularis in diphtheria, 295
- Enterorrhagia, 128
- Enterotome, 31
- Enzymes, toxicology of, 421
- Eosin and methylene blue, 341
- Eosinophilia, 116
- Ephedrine, toxicology of, 422
- Epidemic affections, 449
  - cerebrospinal meningitis, synonyms of, 452
  - erythema, 351
- Epiglottitis, 15, 73, 96, 110, 142, 256
- Epilepsy, 453
- Epiphyseal fractures, 127
- Epispadias, 68
- Epistaxis, 128, 142
- Epithelial nephritis, 199
- Epithelioma of larynx, 143
  - lungs, 156
  - nasal passages, 141
  - tonsils, 143
- Epithelium, œsophageal, 189
- Epulis, 78
- Equipment for bacteriologic investigations, 346
- Equivalents of kilogramme and litre, 357
- Erectile tumor of arteries, 137
- Erection of penis, post-mortem, 68
- Ergotism, 418, 421, 423, 424, 430
- Erlicki's solution for fixation of tissues, 333
- Erosion of bone, 268
- Erosions of vagina, 211
- Eruption, cutaneous, in syphilis, 307
  - herpetic, 141
- Erysipelas, 295
  - ambulans, 295
  - of larynx, 142
  - phlegmonous, 295
  - simplex, 295
  - synonyms of, 449
- Erythema, epidemic, 351
- Erythromelalgia, 250
- Essential anæmias, 116
- Ether, as a freezing agent, 12
  - toxicology of, 21, 421, 429, 430
- Ethereal oil of mustard, toxicology of, 422
  - oils, odor of, 21
  - toxicology of, 422, 425
- Ethyl chlorid, 12
- Eustrongylus gigas, 207

- Eversion of eyelids, 73  
 Evidences of development of infant, 405, 406  
 Evisceration of new-born, 275  
 Exalgin, toxicology of, 423  
 Examination, critical, of abdominal cavity, x, p. 159  
     thoracic cavity, vii, p. 92  
     external, 52, 438  
         of lower animals, scheme for recording, 394  
     in cases of criminal abortion, 406  
     internal, in postmortem, 439  
         of lower animals, scheme for recording, 394  
     microscopic, in postmortem, 438  
     of adrenals, 14, 179  
         aorta, 14, 197, 441  
             of animals, 383  
         appendix, 15, 166, 171, 443  
         base of skull, 233  
         bladder, 14, 15, 181, 442  
         blood, viii, p. 113; 444  
         bones, xvii, p. 261  
             of face, 281  
         bowel in animals, 383  
         brain, xiii, p. 225  
             in animals, 389  
         capsule of kidney, 178  
         child, 275  
         chyle-duct, 197  
         contents of stomach, 189  
         cord, xiv, p. 242  
         diaphragm, 14, 15, 197  
         dura mater, 230  
         ears, 14, xvi, p. 258  
         exocranial sinus, 260  
         exterior of body, v, p. 46  
         external surface of brain, 233  
         eyes, 14, xvi, p. 256  
         female genital tract, 207  
         gall-bladder, 14, 15, 100, 194, 441  
         inner table of skull, 230  
         intestines except the duodenum, 164  
         joints, xvii, p. 261  
         kidney, 14, 178  
         liver, 103, 195  
         lungs, 14, 15, 97, 109, 441  
         macroscopic specimens, 341  
         male genital tract, 211  
         mastoid process, 260  
         mesenteries, 159  
         middle ear, 258  
         nasopharynx, xvi, p. 256  
         œsophagus, 14, 15, 189  
         omentum, 150  
         oral cavity, 281  
         ovary, 14, 15, 183, 442  
         pancreas, 14, 15, 196, 442  
         pericardium, 13, 14, 15, 198, 441  
             in animals, 384  
         peritoneum, 14, 15, 159  
     Examination of prostate, 212  
         retina, 257  
         retroperitoneal lymph-glands, 196  
         salivary glands, 260  
         scalp, 225  
         skull, xiii, p. 225  
         spermatic cord, 212  
         spinal canal, 242  
             cord, xiv, p. 242  
         spleen, 14, 15, 163  
         stomach, 14, 15, 188, 443  
             in animals, 383  
         thoracic duct, 14, 15, 197  
         uterus, 14, 15, 184, 209  
         vagina, 15, 210  
         vena cava, 197  
     order of, at postmortem, ii, p. 13  
     post-mortem, of inoculated animals, 355  
         birds, 394  
         lower animals, 374  
         new-born, xviii, p. 275  
         restricted, xix, p. 280  
     topographic, of parts contained in abdominal cavity, vi, p. 79  
 Examinations, bacteriologic, 14, xxiii, p. 346  
     mediocolegal, post-mortem form of report in, 403, 447  
 Examiners, medical, 1  
 Exanthematous typhus, synonyms of, 448  
 Excision of liver of animals, 379  
 Excitomotor alkaloidal poisons, 421  
 Excrescences, calcareous, of trachea, 146  
     papillomatous, of trachea, 146  
 Exencephalus, 67  
 Exenteration in dorsal position, 381  
     of animals, Vienna method of, 382  
         cranial cavity in animals, 386  
         liver in animals, 381  
         pancreas in animals, 381  
         pelvis in animals, 385  
         spleen in animals, 381, 382  
         stomach, 381  
 Exfoliation of cartilages of larynx, 143  
 Exophthalmic goitre, 147, 452  
     presence of thymus gland in, 95  
 Exophthalmus, causes of, 258  
 Exostosis, 273  
 Expert and court, 399  
     medical man as an, in medicolegal cases, 400  
     testimony, 398, 399, 401  
         sure ground to be adhered to, 401  
         unsafe ground to be acknowledged, 401  
 Exploratory puncture, fatal, 65  
 Exposure of abdominal cavity of animal, 375  
     accessory sinuses in animals, 389  
     thoracic cavity of animals, 376  
         technic of, vii, p. 92  
 Exterior of body, examination of, v, p. 46

- External anthrax, 291  
   developmental period of infant, 405  
   examination of body, v, p. 46; 438  
     lower animals, scheme for recording, 394  
   method of examining testicles, 186  
 Extract of male fern, toxicology of, 422  
 Extrameningeal hemorrhage, 251  
 Extra-uterine pregnancy, 207  
 Extremities, deformity of, in rhachitis, 71  
   of an animal, removal of, 375  
 Exudates, 20, 86  
   acute inflammatory, fixative for, 338  
   chylous, 86  
   distinguished from transudate, 86  
   encapsulated peritonitic, at navel, 66  
   fatty, 86  
   how distinguished, 86  
   odor of suppurative, 86  
   varieties of, 86  
 Eyelids, diseases of, 73, 128, 291, 295, 416  
   movements of, when head has been severed, 47  
   puffiness of, 73  
     in arsenical poisoning, 73  
       Bright's disease, 73  
       cardiac affections, 73  
       traumatism, 73  
 Eyes, 73, 257  
   abnormalities of, 73  
   appearance of, in apoplexy, 48  
     carbonic acid poisoning, 48  
     hydrocyanic acid poisoning, 48  
   artificial, 73  
   color of, 74  
   conjugate deviation of, in apoplexy, 74  
   conjunctiva of, 74, 297  
   consistency of, 258  
   cornea of, 74  
   crystalline lens of, coloboma of, 75  
   diseases of, 67, 73, 74, 75, 76, 253, 257, 295, 297, 301, 306, 308, 432  
   examination of, 14, 47, 48, 73, 75, xvi, p. 257; 438, 444  
   expression of, at death, 73  
   gonorrhœal infection of, 297  
   growths in, 75  
   macroscopic examination of, 257  
   measurements of, 74  
   microscopic examination of, 257  
   optic nerve of, 75  
   orbital injuries of, 76  
   position of, 73, 258  
   post-mortem changes in, 48  
   protrusion of, 258  
   pupils of, 74  
   removal of, 257, 282  
   retraction of, 258  
   sarcoma of, 16  
   sclera of, 75  
   syphilis of, 306  
   tumors of, 16, 74  
   Face, appearance of, after death, 47  
     examination of, 260, 439  
     in death by electricity, 413  
   Facial angle of Camper, 361  
   Facies hypocratica, 47  
   Fæces, color of, 170  
     in partakers of opium, 169  
   Fallopian tubes, 207  
     anomalies of, 207  
     cysts of, 208  
     diseases of, 207, 208, 297, 315  
     examinations of, 14, 15, 185, 442  
     hæmatosalpinx, 208  
     hemorrhages of, 208  
     hydrosalpinx of, 208  
     pregnancy in, 207  
     pyosalpinx of, 208  
     removal of, 183, 184  
     tumors of, 208  
   False aneurism, 137  
     joint, 266  
     membrane of diphtheria, 294  
     sand in intestines, 170  
   Falx, detachment of, 231, 277  
   Fan, electric, 11  
   Farcy, 296, 449  
   Fat, color of, 80  
     embolism of lungs, 148  
     necrosis, 196, 221, 222  
   Fatty changes in liver, 218  
     methods of fixing, 340  
   cirrhosis of liver, 217  
   degeneration, 129, 223, 418  
     of arteries, 135  
     liver, 218  
     marrow, 267  
   infiltration, 129  
     of liver, 18, 219  
     pancreas, 221  
   Favus, 192  
   Fecal fistulas, 457  
   Feet, diseases of, 296, 299, 438  
   Felon, 268  
   Female genital tract, 207  
   Femoral hernia, contents of, 20, 73  
   Fever, break-bone, 293  
     cerebrospinal, 252  
     dengue, 293  
     exanthematous, synonyms of, 448  
     glandular, 295  
     hay, 142  
     intermittent, synonyms of, 448  
     malarial, 322  
     mosquitoes in, 322  
     malta, 300  
     Mediterranean, 300  
     miliary, 351, 449  
     paratyphoid, 169  
     recurrent, synonyms of, 448  
     relapsing, 302  
     rheumatic, 303



- Fever, scarlet, 303, 351  
     distinguished from Duke's fourth  
     disease, 300  
     trypanosoma, 324  
     typhoid, 318, 408, 448  
     typhus, 62, 351, 448  
     yellow, 319, 351, 373, 449  
         mosquito in, 319
- Fevers, infectious, as cause of death, 408  
     abortion from, 407
- Fibrin, agents for fixing, 340
- Fibro-endothelioma, of brain, 255
- Fibroid changes in lung, 153  
     myocarditis, 133  
     phthisis, 311, 315, 316
- Fibroma in nasal passages, 141  
     of bladder, 205  
         brain, 255  
         heart, 134  
         intestines, 171  
         kidney, 204  
         liver, 220  
         lungs, 155  
         ovaries, 209  
         peritoneum, 159  
         pleura, 156  
         tonsils, 143  
         urethra, 70  
         uterus, 209  
         vulva, 69
- Fibromyxoma in nasal passages, 141
- Fibrosis of aortic valves, 129
- Fibrous bronchitis, 145  
     clot of aneurism, 138  
     endophlebitis in veins, 140  
     heart, 129  
     nephritis, 199  
     polyps of heart, 134  
     tumors in nasal passages, 141
- Filaria sanguinis hominis, 159, 202, 207, 321
- Filicic acid, toxicology of, 421
- Fimbriated hymen, 69
- Final report of postmortem, 445
- Fingers, absence of, 68  
     atrophied, 68  
     clubbed, 68  
     hypertrophied, 68  
     supernumerary, 68  
     webbed, 68
- Finger-stalls, rubber, 40
- Firearms, suicide by, 462
- Fish-bone in omentum, 20
- Fish poisoning, toxicology of, 422, 423
- Fissures, of anus, 69  
     cheeks, 68  
     penis, 68  
     sternum, 68
- Fistula in ano, 313
- Fistulae of bronchi, 144  
     rectum following abscesses, 69  
     trachea, 144
- Fixatives, 327, 328, 341  
     pathologic conditions suggest-  
     ing certain, 338  
     penetration as a characteristic of, 329
- Fixing fluids, 329  
     smear preparations, 348
- Flaxseed poultices for post-mortem  
     wounds, 44
- Flax thread, coarse, 38
- Flea, common, 319
- Flemming's fixative, 327  
     solution, 336
- Florida, phthisis, 311, 315
- Fluid, embalming, 9  
     in peritoneal cavity, color of, 85  
         normal amount of, 85  
     pleural cavity, amount of, 97  
         character of, 97  
     peritoneal, collection of, from inocu-  
     lated animals, 356
- Fluids, fixing, 329, 348  
     preserving, 342
- Fluorescence, as a pathologic force, 404
- Fluorescin, 47
- Fly, Spanish, toxicology of, 422  
     tsetse, 324
- Fœtal atelectasis of lungs, 147
- Fœtus, absence of post-mortem rigidity  
     in, 52  
     immature, 52  
     percentage of water in, 360  
     post-mortem expulsion of, 46, 51
- Follicular appendicitis, 171  
     conjunctivitis, 454
- Fontanels, examination of, 444
- Foot-and-mouth disease, 295, 351
- Foot, perforating ulcer of, 248
- Foramen of Monro, opening of, 236
- Forceps, 31, 32  
     curved, 31  
     dissecting, 31  
     other forms of, 32  
     pointed, 31  
     straight, 31
- Foreign bodies in abdominal cavity, 88  
     air-passages, 146  
     heart, 134  
     joints, 267  
     pericardial cavity, 99  
     uterus, 210
- Form of physician's certificate for return  
     of a death in America, 464  
     report in medicolegal post-mortem  
     examinations, 403
- Formad pocket-case, 36
- Formalin (formaldehyde), for embalm-  
     ing, 9  
     for fixing retina, 339  
         tissues, 335  
     hardening brain, 9, 241  
     preserving fluid, 342  
     in preservation of macroscopic speci-  
     mens, 342  
     toxicology of, 21, 421, 430

- Formates and formic acid, toxicology of, 425
- Formic acid, toxicology of, 425
- Fourth disease, Duke's, 300
- Fractures, 20, 70, 266  
by contrecoup, 230  
compound, 266  
epiphyseal, 127  
forms of, 17, 20, 463  
of hyoid bone, 71  
laryngeal cartilages, 71  
neck, 66  
orbit, 76  
overlooked, 71  
recent, usual signs of, 71
- Frænum of tongue, deformities of, 68
- Fragilitas ossium, 265
- Fragmentation of heart, 129
- Framboesia, 296
- Freezing, 463  
agents, 12  
of bodies to preserve, 289  
point of blood, 114  
in drowning, 417
- French method of opening dura, 228  
skull, 228
- Fresh material for microscopic observation, 347
- Freud's gold stain, 333, 341
- Friedreich's ataxia, 248
- Fright, death from, 198
- Frozen bodies, 289, 414, 437  
sections, 12
- Fuchsin, toxicology of, 425
- Fungi-joint in adiposis dolorosa, 59
- Fungi, poisonous, toxicology of, 424
- Funnel-heart of rhachitis, 72
- Furuncle, 461
- Fusiform aneurism, 137
- G**
- Gabbett's solution for staining tubercle bacillus, 310
- Galactotoxismus, 418
- Gall-bladder, 193  
absence of, 215  
bacteriologic examination of, 194  
dimensions of, 367  
diseases of, 195, xii, p. 215; 220  
examination of, 14, 90, 193, 194  
impaction of, due to gall-stones, 216  
measurements of, 367  
removal of, 193  
tumors of, 42, 195  
weight of, 367
- Gall-ducts, diseases of, xii, p. 215
- Gall-stones, 170, 195, 215  
composition of, 216  
number of, 216  
shape of, 216
- Ganglia, semilunar, 181  
spinal, 243  
study of, 181
- Ganglion, 273  
cells, fixatives for, 339  
Gasserian, 282
- Gangrene of bladder, 205  
hand, 45  
lungs, 150, 153, 456  
circumscribed, 150  
diffuse, 150  
penis, 68
- Gangrenous appendicitis, 172  
bronchitis, 146  
dysentery, 173  
endometritis, 210  
ergotism, toxicology in, 423  
lymphangitis of scrotum, 69  
pancreatitis, 220
- Gas bacterium, 349  
bubbles in blood, 121  
fixtures over post-mortem table, 11  
in abdominal cavity, 84  
odor of, in perforation of stomach, 84
- Gases, deleterious, absorption of, 463  
irrespirable, 420, 421
- Gasserian ganglion, 282
- Gastrectasis, 190
- Gastric cancer, 189, 190  
contents, reaction of, 189, 418  
ulcer, situation of, 189  
walls, atony of, 191
- Gastritis, 191  
acute, 191  
atrophic, 192  
chronic, 191, 192  
croupous, 153  
diphtheritic, 191  
due to alcoholism, 427  
hypertrophic, 192  
mycotic, 192  
phlegmonous, 191  
simple, 191  
suppurative, 191  
toxic, 192
- Gastro-intestinal form of influenza, 299  
tract in alcoholism, 427  
syphilis of, 308
- Gastroptosis, 61
- Gastrorrhagia, 128
- Gelatin of Wharton, 276
- Gelsemine, toxicology of, 421, 422
- General considerations, i, p. 1; 439, 440  
in preservation of tissues, 342
- Genitalia, female, diseases of, 56, 69, 207, 405, 438, 444, 460  
removal of, 183  
in acromegaly, 56  
male, diseases of, 56, 68, 211, 405, 438, 444, 459  
removal of, 185, 442
- Genito-urinary system, diseases of, xi, p. 199; 314, 427

- Genito-urinary system in alcoholism, 427  
tuberculosis of, 314
- Genu extrorsum, 71  
valgum, 71, 268  
varum, 71
- German measles, 303  
how distinguished from Duke's  
fourth disease, 300
- Gersung's method, 3
- Gestation, period of intra-uterine, how determined from examination of fœtus, 358, 405
- Giacomini's method of hardening brain, 241
- Giantism, 56, 70
- Gilson's solution for fixation of tissue, 334
- Glacial acetic acid, for post-mortem wounds, 43
- Glanders, 296, 373, 449
- Gland, pineal, see Pineal gland.  
pituitary, see Pituitary body.  
thymus, see Thymus gland.  
thyroid, see Thyroid gland.
- Glands, alcohol as a fixative for, 340  
Bartholinian, enlargement of, 69  
cervical, enlargement of, 295, 297, 301  
diseases causing enlargement of, 66  
lymphatic, tuberculosis of, 66, 316  
mammary, diseases of, 61, 81  
examination of, 81  
tuberculosis of, 317  
maxillary, weight of, 366  
mesenteric lymphatic, hyperplasia of, in typhoid fever, 318  
peribronchial, caseating, 134  
retroperitoneal lymph, examination of, 196  
salivary, 260  
tuberculous, 16
- Glandular fever, 295
- Glass balls, 32  
cones, graduated, 32  
graduated measuring vessels, 32  
vessels for preserving specimens, 34, 444
- Glaucoma, 258
- Glioma of brain, 255  
retina, 75
- Gliosarcoma of brain, 255
- Glisson's capsule, 224
- Glossary index, 22, xxx, p. 489
- Glossina palpalis, 324
- Glottis, œdema of, 426
- Gloves, rubber, 34, 343
- Glycogenic reaction of blood, 120
- Glycosuria, 222
- Gmelin's test for bile, 120
- Goat, acute meningitis of, 253
- Goitre, 146, 249  
colloid, 96  
congenital, 96  
cystic, 146  
exophthalmic, 147, 452
- Gold stains, 341
- Golgi chrome-silver preparation, 341
- Golgi's stains, 332, 341
- Gonococcus, 350
- Gonorrhœal affections of child, 450  
arthritis, 263, 297  
hydrocele, 417  
infection, 297
- "Goose-flesh," appearance of skin in drowning, 66
- Gout, 127, 452  
chalky deposits in, 127  
chronic, 262
- Gouty arthritis, pathology of, 263  
urates in chronic tenosynovitis, 272
- Graduated cones, 33
- Grains, 357
- Grammes, 357  
when used, 17
- Granular kidneys, 179
- Granulation tissue, methods of fixing, 340
- Granulomata of brain, 255  
iris, 75
- Graves's disease, see Basedow's disease.
- Gray hair, cause of, 76
- Grippe, 298, 449  
complications of, 299  
synonyms of, 449
- Guaiacol, effect on urine, 425
- Guinea-pigs, how affected by X-rays, 261  
inoculation of, 45, xxiii, p. 346
- Gumma, 69, 99, 233, 306, 307, 308, 309
- Gummata of pericardium, 99
- Gunshot wounds, in violent death, 408
- Gyrinus natator, toxicology of, 426

## H

- Hæmatemesis, 128
- Hæmatocele, 69  
vaginal, 211
- Hæmatochromatosis, 224
- Hæmatoidin crystals, 120, 201
- Hæmatoma of bones, 274  
ears, 16, 68  
vulva, 69
- Hæmatomyelia, 250
- Hæmatoxylin, effect on urine, 425
- Hæmatozoa, xxi, p. 322
- Hæmoglobin, diffusion of, into tissue, 50  
percentage of, in blood twenty-four hours after death, 113
- Hæmolysins, 421
- Hæmolysis, diseases present in, 116
- Hæmopericardium, 98, 99
- Hæmophilia, 125, 129
- Hæmoptysis, 128, 148  
parasitic, 321
- Hæmothorax, 157
- Hair balls, 189  
color of, 76  
diseases of, 77  
distribution of, 77



- Hair, examination of, 76, 436, 438, 444
  - growth of, after death, 46
  - injuries and tumors under, 20, 77
  - length of, 76
  - loss of, 77
  - quality of, 77
  - quantity of, 77, 306, 307
- Hallux valgus, 269
- Hamilton's method of examining brain, 239
- Hammers, 31
- Hand-bag for carrying instruments, 34
- Hands, amputation of, in spreading gangrene, 45
  - care of, and treatment of post-mortem wounds of, iv, p. 38
  - cellulitis of, how treated, 45
  - diseases of, 45, 59, 68, 246, 250, 254, 299, 303, 304, 307
  - examination of, 47, 68
  - palms of, color of, in corpse, 47
  - post-mortem position of, 51
  - tumors of, 44, 273
- Hanging, 112, 415, 462
  - and throttling, differentiation of, 415
- Hanot's hypertrophic cirrhosis, 115
- Hardening of tissues, 337
  - brain, advantages of, 234
- Harelip, 68
- Harke's method for examining nasopharynx, 256
- Hay fever, 142
- Head, diseases of, 67
  - examination of, 14, 225, 438, 439, 444
  - measurements of, 56, 444
  - of animals, removal of, 386
  - rests, 11
  - size and shape of, 225
- Health, Boards of, 4
- Heart, abnormalities of, 113
  - abscess, metastatic, of, 130
  - actinomycosis of, 134
  - adiposity of, 101, 106
  - amyloid degeneration of, 129
  - anæmic infarct of, cause of sudden death, 130
  - aneurism of, 130
  - anomalies of, 113
  - appearance of, in cholera Asiatica, 293
  - atrophy of, 106
  - beat, absence of, on palpation and auscultation, 47
  - brown, 106
  - carcinoma of, 134
  - characteristics of, vii, p. 100; viii, p. 113
  - color of, 100, 106
  - complications, 455
  - concentric hypertrophy of, 133
  - consistence of, 100, 106
  - cysticerci of, 134
  - degenerations of, 129
  - dilatation of, 106, 133
  - dimensions of, 364
  - diseases of, viii, p. 113; 135, 138, 144, 147, 200, 272, 290, 291, 292, 293, 294, 299, 302, 303, 304, 305, 311, 316, 318, 323, 408, 411, 419, 427, 431, 433, 441
  - displacement of, how determined, 100
  - double, 113
  - eccentric hypertrophy of, 133
  - ecchymoses of, subpericardial, 101
  - echinococci of, 134
  - enlargement of, causes of, 100
  - examination of, 14, 15, 100, 101, 441, 445
    - in animals, 385
  - exhaustion of, as cause of death, 408
  - fatty degeneration of, 106
  - fibromata of, 134
  - fibrous, 129
    - polyps of, 134
  - foreign bodies in, 134
  - fragmentation of, 129
  - hypertrophy of, 106, 133
    - in tuberculosis, 316
  - in animals, dissection of, 385
    - death by electricity, 413
  - infarcts of, 130
  - infiltrations of, 129
  - injuries to, 99, 129
  - in overdoses of digitalis group, 419
  - lesions, as a cause of death, 408
    - valvular, 303
  - lipomata of, 101, 134
  - lymphosarcomatosis of, 134
  - measurements of, 100, 364
  - measuring and testing valves of, 103
  - melanotic sarcoma of, 134
  - method of opening, 101
  - miliary tubercles of, 134
  - muscle, in diphtheria, 106
    - typhoid, 106
  - soft in sepsis, 106
  - suturing of, 100
  - myomalacia of, 130
  - myxomata of, 134
  - organic diseases of, 454
  - pentastomata of, 134
  - primary incisions of, 102
  - removal of, from body, 15, 95, 102, 280, 441
    - in animals, 382
  - rhabdomyomata of, 134
  - right, dilatation of, 144
  - rudimentary, 113
  - sarcomata of, 134
  - sclerosis of, 131
  - secondary incisions of, 104
  - segmentation of, 129
  - simple hypertrophy of, 133
  - spontaneous rupture of, 129
  - syphilis of, 133, 307, 308
  - thrombosis of, 137
  - traumatism of, 99, 129

- Heart, tuberculosis of, 134, 311  
 tumors of, 99, 101, 133, 134  
 valvular diseases of, 133, 200  
 volume of, 365  
 weight of, 105, 106, 360  
   proportion, to body weight, 364
- Heat, death from, 414  
 decomposition in death from, 414  
 in smear preparations, 348
- Heat-stroke, 414  
 post-mortem rigidity after, 51
- Heberden's nodosities, 263
- Heidenhain-Biondi triple stain, 333, 341
- Height, 56  
 average (European standard), 358  
 methods of determining, 56  
   approximate, 56, 57
- Helix of ears, absence of, 68
- Hellebore, toxicology of, 424
- Helvellac acid, toxicology of, 423, 425
- Hemiplegia, 250
- Hemispheres of brain, examining, 239
- Hemorrhage as a complication of scarlatina, 444  
 causes of, 128  
 cerebral, 250  
   as cause of death, 408  
 complementary, 128  
   synonyms of, 453  
 complications of, 453  
 consecutive, 128  
 due to acute yellow atrophy of liver, 95  
   phosphorous poisoning, 95  
   trauma, 95, 128  
   vicarious menstruations, 128
- extrameningeal, 128, 251  
 fatal, 49  
 from cord, as cause of death, 128  
   laceration of liver, 20  
   stomach, 192  
   wounds, as a therapeutic measure, 43
- intestinal, 171, 448  
 into air-cells and lung tissue, 148  
   dura, 230, 233  
   mediastinum, 95  
   mesentery, 160  
   pancreas, 106  
   spinal membranes, 251  
   sympathetic nerves, 112
- meningeal, 251  
 nasal, 128  
 non-puerperal uterine, synonyms of, 459
- of bowel, 171  
   Fallopian tubes, 208  
   liver, fatal, 65  
   lungs, 148  
   pancreas, 220  
   spinal membranes, 251  
   umbilical cord, 128  
   uterus, 209
- Hemorrhage per diapedesis, 128  
 rhexin, 128  
 petechial, 145  
 puerperal, synonyms of, 460  
 pulmonary, 128  
   after calomel injections for syphilis, 148  
 renal, 203  
 secondary, 128  
 subcutaneous, in plague, 302  
 synonyms of, 455  
 varieties of, 128
- Hemorrhagic, chronic, peritonitis, 161  
 endometritis, 210  
 exudate, 86  
 infarct, 136  
   in mesentery, 160  
   scurvy, 127  
   sinks in water, 110
- pachymeningitis, 162  
 pancreatitis, 220  
 pleurisy, 157  
 pyelitis, 203  
 smallpox, 305
- Hemorrhoids, internal or external, 69, 141
- Henoch's purpura, 126
- Hepatic duct, examination of, 194  
 peritoneum, mechanical irritation of
- Hepatitis, endemic, 219  
 pyæmic, 219  
 suppurative, 219  
 tropical, 219
- Hepatization, red and gray, in croup pneumonia, 152, 153
- Hereditary ataxia, 248  
 syphilis, 306
- Hermann's fixative, 327
- Hermaphrodite, 55  
 sex of, 55
- Hernia, 73  
 complications of, 457  
 examination for, 90  
 femoral, containing appendix, 20  
 frequent varieties of, 90  
 infrequent varieties of, 90  
 inguinal, containing sigmoid flexum, 20  
   of scrotum, 69  
   usual locations of, 73  
   vaginal, 211
- Hernial aneurism, 137  
 opening of omentum, 159
- Herpes progenitalis, 69
- Herpetic eruptions, 141
- Hessee's method for test-tube culture, 67
- Heteraluis, 67
- Heterotyphus, 67
- Hewson's preservative injection, 288
- Hexenmilch, 279
- Hev's saw, 20
- Hide-bound skin, 305
- Hide of an animal, removing of, 375
- Hip, congenital dislocation of, 3, 66.

- Hip, diseases of, 66, 266, 301  
 examination of, 444  
 Hirshberg's method for test-tube culture, 353  
 Historical data on postmortems, 1  
 Hodgkin's disease, 124  
 effect upon mesentery, 160  
 post-mortem findings in, 124, 125  
 Hofmann's treatment of discoloration of decomposition, 52  
 Homicide, 408, 411  
 Horizontal plane, accepted, 361  
 section of brain, 239  
 Horse-flesh, how recognized, 360  
 Hour-glass contraction of stomach, 90, 191  
 Housemaid's knee, 273  
 Human tuberculosis, 373  
 Hume's intravenous injection of silver nitrate solution, 44  
 Hutchinson's teeth, 78  
 Hyaline degeneration, 129  
 fixatives for, 340  
 of arteries, 136  
 pancreas, 222, 224  
 Hydatids of liver, 458  
 lungs, 155  
 pleura, 156  
 Hydræmia, 116  
 Hydrarthrosis, 262  
 Hydrate, chloral, toxicology of, 425  
 Hydrocele, 69  
 gonorrhœal, 417  
 mechanical, 417  
 syphilitic, 417  
 tuberculous, 417  
 Hydrocephalic skull, 361  
 Hydrocephalus, 249, 313  
 Hydrochloric acid, toxicology of, 422, 427  
 Hydrocyanic acid poisoning, condition of eye in, 48  
 post-mortem lividity in, 50  
 toxicology of, 50, 411, 421, 425, 430  
 Hydrofluoric acid, toxicology of, 422, 427  
 Hydrogen, arseniuretted, toxicology of, 423  
 peroxide of, 404  
 sulphid, its action on blood, 19, 50  
 Hydromeningocele, 249  
 Hydronephrosis, 200  
 Hydropericardium, 98  
 Hydrophobia, 297, 373  
 Hydrops bursarum, 273  
 Hydrosalpinx, 208  
 Hydrostatic test for lung, as sign of viable child, 404  
 Hydrothorax, 156  
 Hygroma, acute, 273  
 Hymen, 69  
 Hyoscyamine, toxicology of, 422, 423, 426  
 Hyoscyamus, toxicology of, 422, 424  
 Hyperæmia, cerebral, 251  
 hepatic, 219  
 Hyperæmia in croupous pneumonia, 152  
 of spleen, 163  
 Hyperinosis, 115  
 Hyperplasia of mesenteric lymphatic glands in typhoid fever, 318  
 Hyperplastic osteitis, 270  
 Hypertrophic bronchitis, chronic, 145  
 cirrhosis, Hanot's, 115  
 of liver, 18, 217  
 endometritis, chronic, 210  
 gastritis, 192  
 nasal catarrh, chronic, 141  
 pulmonary osteo-arthritis, 247  
 rhinitis, chronic, 142  
 vegetations, vaginal, 211  
 Hypertrophy and atrophy in same part, 18  
 concentric, of heart, 106, 133  
 congenital, 70  
 eccentric, of heart, 106, 133  
 general, 70  
 of anus, 69  
 arteries, 136  
 clitoris, 69  
 fingers and toes, 68  
 glans penis, 68  
 heart, 106, 133  
 concentric, 106, 133  
 eccentric, 106, 133  
 in tuberculosis, 316  
 most marked in double aortic disease, 106  
 simple, 133  
 marrow cells, 267  
 ovary, 209  
 prostate, 212  
 scrotum, 69  
 testicles, 69  
 tongue, 68  
 uterus, 209  
 vulva, 69  
 Hypinosis, 115  
 Hypnotics, toxicology of, 421  
 Hypodermic syringe in inoculation of animals, 383  
 Hypoplasia, 249  
 of aorta, 136  
 testicles, 69  
 uterus, 209  
 Hypospadias, 68  
 Hypostasis or hypostatic congestion, 49  
 Hypostatic congestion, 49, 149  
 pneumonia in typhoid fever, 318  
 Hypsicephalic skull, 361  
 Hysterectomy, vaginal, 281  
 Hysteria, synonyms of, 453
- I
- Icard's test for possible presence of life, 47  
 Ichthyotoxismus, 418  
 Icterus, grave, synonyms of, 458  
 neonatorum, 458, 462



- Icterus, odor in, 170  
   of yellow fever, 319  
 Ictrogen, toxicology of, 425  
   [lupinotoxin] in animals, toxicology of, 423  
 Identification of body, 53  
   before beginning post-mortem examination of, 53  
   by Bertillon system, 53  
     cast of face, 53  
     of mouth, 53  
   clothing, 54  
   evidence of certain diseases, 53  
   photograph, 53  
   physical defect or peculiarity, 53  
   scars, 53  
   skiagraph of old osseous lesions, 53  
   teeth, 53  
   various aids, 53  
 Idiocy, 249  
 Idiopathic anæmia, 116  
 Ileocæcal valve, 166, 312  
 Ileum of animals, removal of, 378  
 Iliac fossa, phlegmon of, and appendicitis, synonyms of, 458  
 Ill-defined diseases, 463  
 Illuminating gas, toxicology of, 50, 411, 431  
 Illumination at postmortem, 11  
 Immunity to staphylococci and streptococci, 43  
 Impaction of gall-bladder, due to gall-stones, 216  
 Imperforate anus, 279  
   hymen, 69  
   mouth, 68  
   rectum, 279  
   vulva, 69  
 Inanition, 463  
   toxic, 418  
 Incense, to remove odor, 10  
 Incising heart, kidneys, etc., see under the various organs.  
 Incision of organs, 20  
 Incompetency, aortic, arteriosclerosis in, 133  
 Index, see p. 22  
 Induration, cyanotic, 133  
 Infant, autopsies on, 403  
 Infanticide, methods of, 414  
 Infantile scurvy, 126  
   uterus, 209  
 Infarct, anæmic, in spleen, 163  
   bilirubin, 201  
   of heart, cause of sudden death, 130  
   spleen, 163  
   hemorrhagic, 136  
   of heart, 130  
   of kidneys, 200  
   of lungs, 148, 149  
   mesentery, 160  
   scurvy, 127  
   uterus, 209  
   syphilitic, 307  
 Infections, gonorrhœal, 297  
   local and general as cause of death, 408  
   paracolon, 319  
   paratyphoid, 319  
   purulent and septicæmic, synonyms of, 449  
 Infectious angina, see Diphtheria and Croup.  
   diseases contracted at postmortem, 41  
   of liver, 220  
   fevers, 50, 202, 214, 218, 220  
   as cause of death, 408  
   of spontaneous abortion, 407  
   processes, acute, fixative for, 338  
 Infective aneurism, 138  
 Infiltration, calcareous, 129  
   of arteries, 135  
   fatty, 129  
   of liver, 18, 219  
   pancreas, 221  
 Infiltrations of heart, 129  
 Inflammation, acute, of prostate, 212  
   caused by poison, 426  
   chronic, 262  
   interacinous, 223  
   of arteries, 136  
   dura, 233  
   lymphatics in post-mortem wounds, 44  
   muscles, 83  
   testicles, 211  
   urethra, 69  
   syphilitic, of veins, 140  
 Inflammatory changes in joint, 71  
   exudates, acute, fixative for, 338  
 Influenza, 298  
   gastro-intestinal form of, 299  
   nervous form of, 299  
   respiratory form of, 299  
 Inguinal canal, examination of, 20, 73  
 Injected specimens, 344  
 Injecting fluid, 9, 288  
   syringe for, 32  
 Injuries, extent of, 20  
   in violent death, 408  
   location, etc., to be noted, 65  
   minute description of, important, 408  
   of vulva, 69  
   overlooked, 20, 77  
   to bones and joints, 267  
   heart and pericardium, 99, 129  
 Inner table of skull, examination of, 230  
 Innominate vein, care not to injure, 93  
 Inoculated tubes, cultivation of, 352  
 Inoculating culture-media, 351  
 Inoculation of animals, 353  
   intravenous, 354

- Inoculation of animals, introperitoneal, 354  
     materials used for, 353  
     media for, 352  
     negative results from, cause of, 353  
     preventive caution needed in, 356  
     site of, 356  
     subcutaneous, 354  
     tube containing medium, 352  
 Inoculum, avoidance of contamination of, 353  
     preparation of, 353  
 Inorganic chemic poisons, 420  
     irrespirable gases, 420  
     irritant poisons, 420  
 Insolation, 463  
 Inspection of body, 13, v, p. 46  
     special organs in animals, 395, 396  
 Instruments, care of, for bacteriologic examination, 346  
     for bacteriologic investigations, 347  
     post-mortem examinations of large domestic animals, 374  
     private post-mortem work, 35  
     post-mortem, and how to use them, iii, p. 28; 374, 446  
     necessary, 28  
     proper handling of, 36  
     sterilizing of, 352, 356  
     to be taken to postmortem, 35  
     various, 32  
 Insular sclerosis, 254  
 Insurance, American, standard of weight, 358  
     in accident company, 45  
 Interacinous, chronic, inflammation, 223  
 Intercarotid bodies, 112, 260  
 Interlobular pleurisy, 158  
 Interments, premature, 46  
 Intermittent fever and malarial cachexia, synonyms of, 448  
 Internal anthrax, 291  
     evidences of period of development of infant, 405, 406  
     examination in postmortem, 439  
     of brain, 234  
     lower animals, scheme for recording 394  
 International system of nomenclature of diseases and causes of death, xxviii, p. 448  
 Interstitial, chronic, pancreatitis, 221  
     nephritis, 199, 201  
     pancreatitis, 221  
     pneumonia, 223  
 Intestinal adhesions, 85  
     contents, removal of, 286, 443  
     hemorrhage, as a complication of typhoid fever, 448  
     lesions of typhoid fever, 318  
     obstructions and hernias, 457  
 Intestinal parasites, 457  
     perforation, as a complication of typhoid fever, 448  
     plague, 301  
     tract, gall-stones and worms in, 170  
 Intestines, adenoma of, 171  
     agglutinated, 165  
     bucket method of opening and cleansing, 167  
     carcinoma of, 171  
     contraction of, 165  
     diseases of, 126, 127, 151, x, p. 159; 217, 219, 291, 292, 293, 299, 304, 306, 308, 311, 317, 318, 319, 320, 408, 411, 426, 428, 430, 431, 432, 433  
     distention of, 165  
     examination of, 14, 15, 164, 440, 443  
     in animals, 376, 383  
     faeces in, 169  
     fibroma of, 171  
     hernia of, 159  
     inflammation of, 159  
     large, diseases of, 165, 166, 167, 318  
     examination of, 164, 165, 166, 167, 442, 443  
     measurements of, 366  
     removal of, 166, 443  
     in animals, 380, 382  
     tumors of, 171  
     weight of, 366  
     ligation of, for removal, 165  
     lipoma of, 171  
     measurements of, 366  
     method of opening, 166  
     removing from body, 165  
     myoma of, 171  
     opening and cleansing of, 166  
     of, in animals, 383  
     other affections of, 457  
     parasites of, 170  
     perforation of, 164  
     polyps of, 171  
     removal of, 15, 165, 288, 443  
     in animals, 380, 382, 383, 393  
     sarcoma of, 171  
     small, diseases of, 165, 308, 311, 432  
     examination of, 15, 164, 165, 166, 167, 442, 443  
     measurements of, 366  
     removal of, 15, 166, 443  
     in animals, 380, 382, 393  
     tumors of, 171  
     weight of, 366  
     stricture of, 306  
     tuberculosis of, 311, 312  
     tumors of, 171  
     weight of, 366  
     worms in, 170  
 Intima of carotids, torn, in death from strangulation, 112  
 Intoxications, trade or occupation, 452  
 Intraperitoneal inoculation of animals, 354  
 Intra-uterine gestation, period of, 405, 444

Intravenous injections, 44  
 inoculations, 354  
 Intussusception, 172  
 Invagination, 90, 91  
 Inversion of bladder, 206  
   eyelids, 73  
 Investigations, bacteriologic, xxiii, p. 346  
 Iodin stains, how removed, 286  
   therapeutic use of, 44  
   toxicology of, 21, 126, 422, 423, 431  
 Iodoform for post-mortem wounds, 43  
   toxicology of, 21, 421  
 Ipecacuanha, toxicology of, 422  
 Iris, 75  
 Iritis, 297, 306  
 Iron of hæmoglobin decomposed by hydrogen sulphid, 19  
   tripods, 32  
   useful in axillary cellulitis, 44  
 Irrisable gases, organic and inorganic, 420, 421  
 Irritability, muscular and nervous, loss of, how determined, 47  
 Irritant inorganic poisons, 420  
   organic poisons, 421  
   poisons, 420, 421  
 Irritating dusts, toxicology of, 422  
   vapors, toxicology of, 422  
 Ischiopagus, 67  
 Islands of Langerhans in diabetes, 223, 224  
 Itch, 461  
 Itch-mite, 320

## J

von Jaksch's anæmia, 124  
 Jalap, toxicology of, 424  
 Jars, quart, museum, for specimens, 34  
 Jaundice, 153  
   appearance of skin in, 60  
   in eyes, 74  
 Jaw, deformities of, 257  
   examination of articulation of, 260  
   necrosis of, 434  
 Jejunum of animals, removal of, 378  
 Jigger, 319  
 Johnson's mixture, 257  
 Joint, condition of, in gout, 128  
   Charcot's, 284  
   cricoarytenoid, ankylosis of, 143  
   diseases of, 72, 126, 127, 153, xvii, p. 261; 293, 297, 299, 303, 305, 419  
   empyema of, 261  
   examination of, 14, 261, 439  
   false, 266  
   foreign bodies in, 267  
   inflammation of, 261  
   injuries to, 267  
   removal of, 261  
   rheumatism of, 303  
   tumors of, 261, 263  
 Jumbo, weight of heart of, 365  
 Jumping from high places, suicide by, 463

## K

Kairin, effect on urine, 425  
 Kaiserling fluid, 343  
 Karorosis of vulva, 69  
 Karyokinesis, fixation of tissues for, 337  
 Keratitis, 74  
 Kidney, abnormality of, 20, 199  
   absence of, 199  
   adrenal tissue in, 204  
   amyloid changes in, 199, 307  
   angiomata of, 204  
   anthrax of, 292  
   atheromatous, 179  
   calculi of, 204  
   capsule of, examination of, 178  
   carcinoma of, 204  
     sarcomatodes of, 204  
   cholera Asiatica of, 293  
   color of, 178, 179  
   concretions in, 200  
   congenital cystic, 200  
     defects in, 199  
   congestion of, 199, 200  
   cortex of, relation of, to medulla, 179  
   cystic disease of, 200  
   cysticerci of, 202  
   decapsulation of, 3, 179  
   degeneration and regressive processes in, xi, p. 199  
   dimensions of, 368  
   diseases of, 64, 85, 98, 99, 100, 116, 117, 126, 127, 128, 132, 135, 136, 142, 144, 145, 151, 153, xi, p. 166; 217, 220, 291, 293, 294, 295, 299, 300, 303, 304, 307, 311, 313, 323, 407, 408, 410, 411, 418, 426, 427, 428, 431, 434, 435, 459  
   distoma hæmatobium of, 202  
   echinococci of, 202  
   examination of, 14, 178  
   fibromata of, 204  
   fibrous changes of, 199  
   filaria sanguinis of, 202  
   floating, 175  
   granular, 179  
   horseshoe, 199  
   incising of, 178  
   infarcts in, 200  
   in postmortem, 442  
   interstitial nephritis of, 201  
   lesions in, in scarlet fever, 304  
   lipomata of, 204  
   liver-tissue in, 204  
   lobulation of, 199  
   lymphadenomata of, 204  
   lymphomata of, 204  
   measurements of, 368  
   method of opening, 177, 178  
     removing from body, 176  
   movable, 202  
   myomata of, 204  
   myxomata of, 204  
   opening of, 178



- Kidney, parasites in, 202
    - parenchymatous nephritis of, 202
    - pentastoma of, 202
    - poisons especially affecting, 418
    - ratio of weight of, to that of body, 368
    - removal of, 38, 64, 176, 177, 275, 280, 442, 444
      - in animals, 356, 380
    - rhabdomyomata of, 204
    - right and left, distinguishing points between, 368
    - sarcoma of, 204
    - scars of, 179
    - specific gravity of, 368
    - stones in, 204
    - strongylus gigas of, 202
    - syphilis of, 308
    - traumatism of, 199
    - tuberculosis of, 311, 314
    - tumors of, 179, 204
    - vascular changes in, 199
    - weighing of, 179
    - weight of, 177, 179, 360, 367
  - Killing of animals for necropsy, 374
  - Klebs-Löffler bacillus, 145, 293
  - Knives, 28
    - amputation, 29
    - brain, 29
    - curved, probe-pointed bistoury, 29
    - pointed, danger of, 37
    - section or cartilage, 29
    - Valentine, 29
    - Waring's, 29
  - Knock-knee, 71, 268
  - Koch's inoculation syringe, 353
  - Kolisko, 8
  - Kreotoxismus, 418
  - Kromskop, its usefulness, 18
  - Kupffer's cells of liver, 224
  - Kyphosis, 198, 246, 269
- L**
- Labelling tissues preserved for microscopic purposes, 327
  - Labor, normal, 460
  - Lacerations, 17, 20, 69
  - Lack of care as cause of death, 462
  - Ladles, 33
  - Landmarks, regional, 17
  - Lardaceous degeneration, 144
  - Large intestine, see Intestines, large.
  - Laryngeal cavities, exposure of, 256
    - diphtheria, 294
    - stenosis, 143
  - Laryngis pachydermia, 143
  - Laryngitis, acute catarrhal, 142
    - chronic, 142
    - diphtheritic, 142
    - edematous, 142
    - sicca, 143
    - syphilitic, 309
    - tubercular, 315, 449
  - Larynx, carcinoma of, 143
    - deformity of, 142
    - diseases of, 97, 128, 141, 142, 133, 146, 246, 294, 297, 299, 300, 305, 306, 307, 309, 312, 315, 318, 455
    - epithelioma of, 143
    - erysipelas of, 142
    - examination of, 14, 15, 442, 445
    - exfoliation of the cartilages of, 143
    - fibroma tuberosum of, 143
    - lupus of, 318
    - papillary fibroma of, 143
    - papilloma in, 143
    - perichondritis of, 143
    - removal of, 37, 442
    - sarcoma of, 143
    - syphilis of, 307, 309
    - tuberculosis of, 315
    - tumors of, 143
    - ulcerations of, 143
  - Lateral sinus, thrombosis of, 112
  - Lathyrism, 418
  - Lavatories, 12
  - Law permitting post-mortem examinations, 5
  - Lead, toxicology of, 411, 420, 422, 423, 424, 425, 431
    - in animals, 421, 422, 423, 424
  - Length and weight of a full-term sound child, 358
  - Lenhossek's stain, 341
  - Leontiasis ossea, 70, 247
  - Leprosy, 299
    - anæsthetic form of, 299
    - synonyms of, 449
    - tubercular form of, 299
  - Leptocephalic skull, 361
  - Leptomeningitis, 139, 233, 253
    - acute purulent cerebrospinal, 251
    - chronic deep, 253
    - diffuse, 253
  - Lesions, see diseases of the various organs.
    - cortical, 237
    - distribution of, 16
    - suggestive, 16
  - Letulle's method of making postmortem, 15
  - Leucocythæmia, blood changes in, 123
    - complications of, 452
    - effect on mesentery, 160
    - lymphatic, 124
    - medullary, 123
    - mixed, 124
    - pseudo-, 124
    - specific gravity of blood in, 114
    - splenic, 164
  - Leucocytosis, 117
  - Leucopenia, 117
  - Leukæmia, see Leucocythæmia.
  - Library, 12
  - Lice on body, method of quickly removing, 59

- Ligaments, diseases of, 73, 128, 262, 268  
 Light, artificial, 8  
 Lightning, death by, 49, 413  
 Limb, atrophy of, 56  
   examination of, 438  
   shortening of, 56  
 Lime, deposits of, in colloid goitre, 96  
 Lindsay Johnson's mixture, 257  
 Lion-forceps, 32  
 Lipæmia, 117  
 Lipoma, arborescent, 263, 273  
   of bone, 273  
   eye, 75  
   heart, 134  
   intestines, 171  
   kidney, 204  
   peritoneum, 159  
   pleura, 156  
   vulva, 69  
 Lips, diseases of, 68, 141, 295, 296, 307  
 Liquefaction of lungs, 153  
 Literature, xxix, p. 465, and many foot-  
   notes and references in the text  
 Lithography and three-color printing, 18  
 Lithopædion, 208  
 Lithosis, 155  
 Liver, abscess of, 213, 219  
   aberrant adrenal tumors of, 220  
   accessory, 367  
   acidity of, 418  
   acute yellow atrophy of, 213  
   adenocystomata of, 220  
   adenomata of, 220  
   adrenal tissue in, 91, 220  
   after tight lacing, 91  
   amyloid degeneration of, 214, 307  
     fixative for, 340  
   and gall-bladder, 193, 213  
   angioma of, 220  
   anthracotic cirrhosis of, 217  
   appearance of, in cholera Asiatica, 293  
   atrophic cirrhosis of, 216  
   atrophy of, acute yellow, 213  
   bile capillaries of, fixative for, 340  
   cancer of, 214  
   capsular cirrhosis of, 217  
   carcinomata of, 220  
   cirrhosis of, 216, 217, 224, 417, 427, 458  
   color of, 195  
   congestion of, 218  
   consistency of, 195  
   cyanotic cirrhosis of, 217  
   dimensions of, 367  
   discolorization of, in ammonium hy-  
     drate poisoning, 19  
   diseases of, 18, 20, 61, 65, 66, 85, 86,  
     94, 95, 112, 116, 122, 124, 127, 135,  
     140, 144, 145, 148, 151, 153, 161, 174,  
     189, 190, 200, 201, xii, p. 213; 221,  
     222, 223, 246, 280, 293, 295, 300, 302,  
     301, 307, 308, 309, 311, 313, 317, 318,  
     319, 323, 324, 408, 411, 418, 427, 428,  
     431, 433, 434  
   Liver, ducts of, diseases of, xii, p. 213  
     echinococcus of, 322  
     emphysema of, 218  
     examination of, 14, 15, 91, 193, 194,  
       195, 276, 442, 443, 444  
     exenteration of, in animals, 381  
     extra lobes of, 91  
     fatal hemorrhage from, 65  
     fatty changes in, 218  
       cirrhosis of, 217  
       degeneration of, 218  
       infiltration of, 18, 219  
     fibromata of, 220  
     fixatives for, 340  
     flukes, 321  
     hemorrhage from, 20  
     hepatitis of, 219  
     hydatid tumors of, synonyms of, 458  
     hyperæmia of, 219  
     hypertrophic cirrhosis of, 18, 217  
     in cholera Asiatica, 293  
       general tuberculosis, 313  
     infectious diseases of, 220, 293  
     inflammation of, 219  
     injury to, 219  
     in postmortem, 443  
     laceration of, from external violence,  
       20  
     lobules of, 195  
     lymphoma of, as a lesion in typhoid  
       fever, 318  
     malarial cirrhosis of, 217  
     measurements of, 367  
     method of removing from body, 193  
     multiple abscess of, 219  
     of animals, excision of, 379  
     other affections of, 458  
     parasites of, 220  
     passive congestion of, 218  
     pernicious anæmia, fixative for, 340  
     poisons especially affecting, 418  
     position of, 195  
     primary cancer in, 215  
     psorosperms in, 323  
     removal of, 15, 193, 194, 275, 443, 444  
       in animals, 356, 381, 393  
     rhachitic cirrhosis of, 217  
     sarcoma of, 16  
     scarlatinal cirrhosis of, 217  
     secondary cancer in, 214  
     specific gravity of, 367  
     supernumerary, 91  
     syphilis of, 309  
     syphilitic cirrhosis of, 217  
     tumors of, 16, 214, 215  
     volume of, 367  
     weight of, 360, 366, 367  
 Lividity, cadaveric, cause of, 49  
   color of, 49, 50  
   degree of, depends on fluid condition  
     of blood, 50  
   distinguished from beginning decom-  
     position, 50

- Lividity, cadaveric, distinguished from  
bruise, 49, 50  
in sunstroke, 414  
location of, 49  
maximum time of, 49  
post-mortem, 49  
rose-red color of, 50  
time of appearance of, 49
- Lobar pneumonia, 151, 152
- Lobeline, toxicology of, 422, 424
- Lobular pneumonia, 151, 154, 299
- Lobulation of kidney, 199  
pancreas, 221
- Locomotion, diseases of organs of, 461
- Locomotor ataxia, 248, 452
- Longevity of bacterial spores, 356
- Longitudinal sinus, 123, 231, 277
- Loose bodies in joint, 265
- Lordosis, 197, 269
- Lorenz operation, 3
- Louse, 319
- Luer's rhachiotome, 242
- Lugol's solution, 214
- Lumpy jaw, 290
- Lungs, abscesses of, 147  
adenoma of, 155  
air embolism of, 148  
anæmia of, 148  
apoplexy of, 148  
atelectasis of, 147  
brown induration of, 179  
cancer, secondary, of, 155  
carcinoma of, 155  
changes in color of, 97  
chondroma of, 155  
circulatory disturbances of, 148  
cirrhosis of, 155  
color of, at different ages, 97  
cut surface of, 109  
in heart disease, 109  
normal, 109  
congenital, changes in, as blood is  
oxidized, 276  
congestion of, 19, 148  
consolidation of, 109  
contraction of, 97  
dermoid cysts in, 155  
dimensions of, 366  
diseases of, 16, 20, 61, 65, 67, 72, 80,  
85, 94, 95, 97, 98, 99, 100, 109, 110,  
123, 126, 127, 128, 132, 138, 139, 140,  
ix, p. 147; 190, 200, 223, 246, 247,  
252, 261, 269, 270, 272, 290, 293, 294,  
296, 299, 300, 301, 302, 303, 308, 309,  
311, 312, 315, 318, 324, 408, 431, 433,  
434, 435  
emphysema of, 149, 456  
examination of, 14, 15, 97, 108, 110,  
152, 276, 280, 441, 442, 444, 445  
in animals, 385  
expanded, appearance of, 97, 276  
fat embolism of, 148  
fibroma of, 155  
Lungs, gangrene of, 150, 456  
hæmoptysis of, 148  
hemorrhage of, 148  
hydatids in, 155  
hypostatic congestion of, 149  
in animals, dissection of, 385  
incising of, 108, 109  
infarcts of, 148, 149  
malignant growths of, 155  
measurements of, 366  
microscopic examination of, 109  
œdema of, 149  
osseous formations in, 155  
osteoma of, 155  
palpation of, 109  
parasites in, 151  
pneumonic consolidation of, 151  
pneumoconiosis of, 155  
post-mortem appearances of, in deter-  
mining viability of child, 404  
purulent œdema of, 145  
removal of, 15, 95, 107, 280, 441  
in animals, 356, 382  
specific gravity of, 366  
spindle-celled sarcoma of, 156  
superficial examination of, 97  
suppuration of, 153  
syphilis of, 309  
traumatism of, 155  
tuberculosis of, 147, 311, 312, 315  
tumors of, 155  
colloid, 156  
encephaloid, 156  
epitheliomatous, 156  
scirrhus, 156  
unexpanded, appearance of, 276  
weight of, 109, 360, 365  
Lung-tissue, hemorrhage into, 148  
Lupinotoxin, toxicology of, 423  
Lupus vulgaris, 318  
of larynx, 318  
vulva, 69  
Luxations, 266, 463  
Lymph, collections of, mistaken for other  
lesions, 89  
tumors, 140  
vessels, diseases of, viii, p. 140  
Lymphadenitis in scarlet fever, 304  
Lymphadenoma, 124, 204  
Lymphangitis, 69, 153  
Lymphangioma, 140  
Lymphatic glands, diseases of, 16, 44, 66,  
89, 94, 116, 123, 124, 125, 140,  
153, 190, 224, 291, 293, 294, 295,  
297, 301, 304, 306, 307, 316, 455  
examination of, 15, 61, 66  
involvement of, in post-mortem  
wounds, 44  
mediastinal, pigmentation of, 95  
mesenteric, hyperplasia of, in ty-  
phoid fever, 318  
mistaken for melanotic sarcoma,  
95



Lymphatic glands, new growths of, 196  
 of axilla, inflammation of, in post-mortem wounds, 44  
 removal of, in animals, 356  
 retroperitoneal, examination of, 196  
 tuberculosis of, 316  
 tumors of, 41, 140  
 system, affections of, 455  
 vein, great, 15  
 vessels, examination of, 442, 443  
 Lympho-carcinoma in mediastinum, 146  
 Lymphocytosis, diseases present in, 116  
 Lymphoid marrow, 267  
 Lymphoma in mediastinum, 146  
 of kidney, 204  
 liver, a lesion in typhoid fever, 318  
 tonsils, 143  
 Lymphosarcomatosis of heart, 134  
 Lymphotoxæmia in new-born, 279  
 Lymph-vessels, carcinoma of, 140  
 dilatation of, 140  
 endothelioma of, 140  
 lesions of, 140  
 syphilis of, 140  
 tuberculosis of, 140  
 Lysol, effect on urine, 425

## M

Macrocytosis, 117  
 MacroGLOSSIA, 140  
 MacroGLOSSUS, 68  
 Macroscopic specimens, examination of, 341  
 preservation of, xxii, p. 326  
 Macrostoma, 68  
 Magnifying-glass, 33  
 Magnus's test of circulation, 48  
 Malachite green, toxicology of, 422  
 Malacia, 223  
 Malaria, 322, 373  
 æstivo-autumnal, 322  
 eruptions in, 62  
 pernicious, 322, 323  
 quartan, 322  
 simple, 322  
 tertian, 322  
 Malarial cachexia, 323  
 chronic, 322  
 cirrhosis of liver, 217  
 fever, 322  
 Mal de caderas, 324  
 Male fern, toxicology of, 422  
 genital tract, examination of, 185, 211  
 Malformations as cause of death, 408  
 due to disease, 462  
 in anæmia, 72  
 aneurism, 72  
 chronic lung condition, 72  
 nervous diseases, 72

Malformations of nose, 141  
 osteitis deformans, 72  
 osteomalacia, 72  
 rhachitis, 72  
 syphilis, 72  
 testicles, 69  
 tuberculosis, 72  
 Malignant angina, see Diphtheria.  
 disease, see Cancer of various organs.  
 pustule, 291, 449  
 Mallets, 31  
 Mallory's anilin-blue stain, 324  
 Malpractice, 397, 398  
 Malta fever, 300  
 Mammary glands, 81  
 colostrum in, 81  
 fibrous tissue in, 81  
 incisions in, 81  
 lactation of, 81  
 of new-born, secretions in, 279  
 tuberculosis of, 317  
 tumors of, 61  
 Manometer, 47  
 Marantic thrombi, 136  
 Marrow, atrophy of, 267  
 cells, hypertrophy of, 267  
 changes in, in osteomyelitis, 267  
 fatty degeneration of, 267  
 in children, appearance of, 267  
 middle life, appearance of, 267  
 lymphoid, 267  
 method of fixing, 338  
 Mask, death, 289  
 Massage, stiff joints prevented by, 44  
 Mast-cells, fixative for, 340  
 Mastoid disease, 246  
 Materials, collection of, for culture purposes, 346  
 used for inoculation, 353  
 Maturity and period of intra-uterine gestation determined at postmortem, 444  
 Maxilla, alveolar compartments in, 405  
 Maxillary glands, weight of, 366  
 Maximum weight of encephalon, 361  
 McKinley, postmortem on, 6  
 Measles, 300, 351, 448  
 complications of, 448  
 German, 300, 303  
 Measures and weights, xxiv, p. 357. See also various organs.  
 Mechanical congestion, 149  
 hydrocele, 417  
 Meckel's diverticulum, 165  
 typhoid ulcer in, 19, 20  
 Meconium, 214, 278  
 Media for isolation of pathogenic micro-organisms, 350, 351  
 Mediastinal adenitis, simple, 146  
 suppurative, 146  
 tuberculous, 146  
 hemorrhage, 95  
 Mediastinitis, 146  
 Mediastino-pericarditis, 95

- Mediastinum**, abscess of, 95  
     carcinoma of, 146  
     dermoid cysts of, 146  
     diseases of, 94, 95, 99, 146  
     emphysema of, 95  
     examination of, 95, 441  
     lympho-carcinoma of, 146  
     lymphoma of, 146  
     tuberculosis of, 95  
     tumors of, 66, 95, 100, 146  
**Medical examiners**, 1  
     men as experts, 400  
**Medicine**, practice of, compensation in, 398  
**Medicolegal and pathologic postmortems**,  
     difference between, 402, 437  
     cases, medical men as experts in, 400  
         Prussian regulations for performance of autopsies in, xxvii, p. 436  
     examinations, 53  
     postmortem examinations, form of report in, 403  
         in cases of poisoning, 402, 443  
     references, xxix, p. 474  
     suggestions, xxvi, p. 397  
**Mediterranean fever**, 300  
**Medulla oblongata**, diseases of, 253, 297  
     examination of, 236, 439  
     removal of, 236  
     incising of, 236  
     weight of, 362  
     pons, and basal ganglia, fixatives for, 339  
**Medullary cancer of stomach**, 190  
     leukæmia, 124  
**Melanin**, effect on urine, 425  
**Melanosarcoma of lungs**, 156  
**Melanosis**, arsenical, toxicology in, 422  
**Melanotic sarcoma of pericardium**, 99  
     diagnosis of, 99  
     of heart, 134  
**Melnikow-Raswedenkow fluid**, 343  
**Membrane**, diphtheritic, 294  
     false, of diphtheria, 294  
     mucous, appearance of, in hydrophobia, 297  
     condition of, in cholera Asiatica, 293  
     preservation of, 327  
     serous, preservation of, 327  
     tuberculosis of, 317  
     spinal, hemorrhage of, 251  
**Membranous, chronic, nasal catarrh**, 141  
     colitis, 174  
**Mendel's law**, a law of heredity experimentally shown to exist by Mendel in 1865 and but recently rediscovered, in which one-fourth of the characteristics will be like each parent and the other half will possess recessive attributes, 2  
**Ménière's disease**, aural vertigo.  
**Meningeal hemorrhage**, 251  
**Meningeal plague**, 301  
**Meninges**, diseases of, 14, 16, 128, 132, 251, 252, 253, 299, 301, 313, 408, 411  
**Meningitis**, acute cerebrospinal, 252, 408, 448, 452  
**Meningocele**, 68, 75  
**Meningo-encephalitis**, 16, 253, 313  
**Menorrhagia**, 128  
**Mental alienation**, forms of, 453  
**Mentha pulegium**, toxicology of, 424  
**Menthol**, effect on urine, 425  
**Merbane**, 411  
**Mercaptan**, odor of, 21  
**Mercurial poisoning**, toxicology of, 418, 421, 422, 423, 424, 425, 426, 432  
**Mercury**, bichlorid of, for fixing tissues, 333  
**Mesenteric artery**, superior, thrombosis of, 137  
     glands, diseases of, 89, 90, 160, 293, 306, 318  
     examination of, 89, 160  
     hyperplasia of, in typhoid fever, 318  
     tumors of, 160  
**Mesentery**, disease of, 86, 159, 160, 317  
     examination of, 14, 15, 159, 160, 442  
     shortened, 160  
     tumors of, 86, 160  
**Metabolism**, carbohydrates in, 222  
**Metals**, acid salts of, toxicology of, 424  
**Metapagus**, 67  
**Metastasis due to cancer of stomach**, 190  
**Methæmoglobinæmia**, 419  
**Method**, angular, of removing calvarium, 226  
     author's, of examining male genital organs, 186  
         fixing skull-cap, 284  
         preparing frozen sections, 12  
         removing brain in child, 276  
         testing gums for lead, 423  
     circular, of removing calvarium, 226  
     Dejerine's, of examining brain, 238  
     external, of examining testicles, 186  
     French, of opening adult skull, 228  
         dura, 228  
     Giacomini's, of examining brain, 241  
     Hamilton's, of examining brain, 239  
     Harke's, of examining nasopharynx, 256  
     Letulle's, of removing organs, 15  
     Meynert's, of examining brain, 237  
     of closing abdominal incision, 285  
         committing suicide, 410  
         cutting ribs of animals, 377  
         scalp, 225  
     diagnosing rabies, 298  
     dividing the dura, 231  
     embalming cavities, 286  
     injecting embalming fluid, 286  
     killing domestic animals for post-mortem examination, 374

- Method of loosening adherent dura, 227  
 calvarium, 227  
 making a death-mask, 289  
 post-mortem examination of  
 animals, 355  
 smear preparations, 349  
 opening and cleansing intestines,  
 166  
 brain, 239  
 canal of Wirsung, 188  
 cephalic cavities of horse, 387  
 intestines, 166  
 kidney, 177, 178  
 œsophagus, 188  
 spinal canal, 242  
 stomach, 188  
 performing autopsy without vis-  
 ible mutilation, 280  
 criminal abortion, 406, 407  
 infanticide, 414  
 postmortems on animals,  
 xxiii, p. 346; xxv, p. 373  
 preserving natural color of speci-  
 mens, 344  
 removing adrenal, 176  
 brain, 232  
 calvarium, 226  
 child's brain, 276, 277  
 extremities of an animal, 375  
 hide of an animal, 375  
 intestines from the body, 165  
 kidney from the body, 176  
 liver from the body, 193  
 male organs of generation,  
 185  
 posterior extremity of an ani-  
 mal, 375  
 spinal cord, 242  
 from body of baby, 277  
 spleen, 162  
 replacing scalp, 285  
 restoring and preserving the body,  
 283  
 skull, 283, 284  
 sectioning brain, 235  
 Pitres's, of examining brain, 237  
 quick, for removing brain, 233  
 Virchow's, of examining brain, 235  
 Methyl alcohol, toxicology of, 21, 432  
 Methylene blue, effect on urine, 425  
 Metritis, 459  
 Metrorrhagia, 128  
 Meynert's method of examining brain,  
 235, 239  
 Micrencephaly, 249  
 Microbic poisoning, 16  
 Microcephalic skull, 361  
 Micrococcus gonorrhœæ, 350  
 melitensis, 300  
 Microcytosis, 117  
 Microglossus, 68  
 Micromegaly, 247  
 Micromyelia, 240  
 Micro-organisms, diseases due to, xxi, p.  
 290  
 in appendicitis, 171  
 peritonitis, 160  
 pathogenic, media for isolation of,  
 350, 351  
 present at postmortem, 43  
 Microscopic examinations in postmortem,  
 438  
 specimens, preservation of, xxii, p. 326  
 study, preservation of tissues for, 326  
 Microsomia, 70  
 Microstoma, 68  
 Middle ear, 258, 304  
 Miliary, acute, tuberculosis of brain and  
 cord, 313  
 aneurism, 138  
 carcinosis, 160  
 fever, 351  
 tubercles, lymph mistaken for, 89  
 of adrenals, 180  
 heart, 134  
 peritoneum, 89  
 ureter, 182  
 tuberculosis, 311, 312, 313, 315  
 of lungs, 311  
 peritoneum, 317  
 serous membranes, 317  
 Milk spots, 151  
 Mineral acids, toxicology of, 424, 426  
 Mitral, buttonhole of, 133  
 diseases of, 103, 129, 133, 303  
 examination of, 101, 103  
 orifice of heart, dimensions of, 365  
 stenosis, 133  
 Molar death, 326  
 Molecular death, 326  
 necrosis, 262  
 Moles of uterus, 209  
 Mollineux trial, 55  
 Mollities ossium, 270  
 Monoxid, carbon, toxicology of, 425  
 Monsters, 67  
 double, 67  
 parasitic, 67  
 single, 67  
 triple, 67  
 Morbidity statistics, xxviii, p. 448  
 Morbilli, see Measles.  
 Morgue, 9  
 Morphine, 65, 418  
 administered before electrocution, 413  
 toxicology of, 413, 418, 421, 422, 423,  
 424, 425  
 Morphism, toxicology in, 421  
 Mortopsy, see Postmortem.  
 Mortuary room, 10  
 Morvan's disease, 452  
 Mosquitoes, anopheles, 322  
 as disseminators of dengue, 373  
 malaria, 373  
 yellow fever, 319, 373  
 stegomyia, 319



- Mouth and its adnexa, diseases of, 60, 64, 67, 68, 294, 295, 296, 301, 309, 310, 312, 426, 428, 432, 456  
examination of, 53, 60, 281
- Movable kidney, 202
- Mucoid material, fixatives for, 340
- Mucous fever, see Typhoid fever.
- membrane, appearance of, in hydrophobia, 297  
condition of, in cholera Asiatica, 297  
of corpse, 47  
stomach, action of poisons on, 419  
patch, 69, 306, 307
- Müller's fluid for fixing tissues, 234, 257, 332
- Multiparæ, weight and dimensions of uterus in, 371
- Multiple abscess of liver, 219
- Mummification, 49
- Mumps, 301, 351
- Murexid test for tophi, 73
- Muscarine, toxicology of, 422, 423, 424
- Muscle-fibres, preservation of, 327
- Muscles, atrophy of, due to rheumatism, 303  
progressive spinal, 253  
color of, in anæmia, 82  
contracture of, disappearance of, after death, 71  
diseases of, 63, 72, 82, 83, 94, 253, 254, 292, 297, 303, 305, 306, 318, 321, 416, 428, 431, 432, 433, 444  
examination of, 63, 81, 444  
external examination of, 61  
hemorrhages in, 82, 318  
inflammations of, 83  
involuntary, post-mortem contraction of, 49  
irritability of, loss of, 47  
lesions in, 82  
papillary, 129  
parasites in, 83  
syphilis of, 306  
tumors in, 83, 84  
twitchings of, in amputated limbs, 47  
waxy degeneration of, in typhoid fever, 318  
weight of, 360
- Museum, 12
- Mustard, oil of, 422
- Mycobacterium lepræ, 299, 350  
tuberculosis, 350
- Mycotic aneurism, 138  
gastritis, 192
- Myelin sheaths, fixatives for, 339
- Myelitis, acute, 254
- Myelocytosis, diseases present in, 116
- Myelomeningocele, 249
- Myelotome, Pick's, 29
- Myiasis, 321
- Myocarditis, acute circumscribed interstitial, 130  
associated with pneumonia, 153  
chronic interstitial, 131  
pathologic changes in, 131  
diffuse interstitial, 131  
due to rheumatic fever, 303  
fibroid, 133  
fibrous, 131  
parenchymatous, 130  
segmentary, 130
- Myocardium, examination of, 105
- Myoma of bladder, 205  
intestines, 171  
kidney, 204  
tonsils, 143  
vulva, 69
- Myomalacia cordis, 131  
of heart, 130
- Myositis, acute, 83  
chronic, 83  
parenchymatous, 83  
progressive ossifying, 83
- Mytilotoxismus, 418
- Myxœdema, 147  
presence of thymus gland in, 95
- Myxoma, fixatives for, 340  
of bone, 274  
brain, 274  
heart, 134  
kidney, 204  
nasal passages, 141
- N**
- Nagana, 324
- Nails, 77  
anomalies of, 77  
diseases of, 77, 307  
examination of, 77, 444  
growth of, 77  
matrix, suppuration of, how treated, 45
- Name, legal, for record, 53
- Narcosis, chloroform, death from, 198
- Narcotics, toxicology of, 421, 424
- Nasal and cranial cavities of ruminant, lines used in sawing, 390  
catarrh, acute, 141  
atrophic, 141  
chronic, 141  
fibrinous, 141  
hypertrophic, 141  
membranous, 141  
cavities, exposure of, 256  
diphtheria, 294  
fossæ, angiosarcoma in, 141  
carcinoma in, 141  
chondroma in, 141  
diseases of, 455  
passages, cysts in, 141  
diseases of, 141  
epithelioma in, 141

- Nasal passages, fibroma in, 141  
 fibromyxoma in, 141  
 foreign bodies in, 142  
 osteoma in, 141  
 rhinoliths in, 141  
 sarcoma in, 141  
 tumors in, 141  
 tuberculosis, 141
- Nasopharynx, 274, 294, 299  
 examination of, 14, 233, xvi, p. 256:  
 282  
 in animals, 389
- Natator gyrinus, toxicology of, 426
- Native Africans, variations in color of  
 skin of, 60
- Nauwerck's method of embalming, 287  
 discovering seminal vesicles,  
 187
- Navel, see Umbilicus.
- Neck, diseases of, 71, 72  
 dislocation of, 66  
 examination of, 66, 71, 72, 112, 438,  
 442, 444  
 in postmortem, 440, 442  
 fracture of, 66  
 inspection of, 66  
 tracheal fistula of, 68  
 visceral clefts of, cysts of, 68
- Necropsy, see Postmortem.
- Necrosis, appearance of bone in, 268  
 centralis, 268  
 diffuse, 268  
 fat, 196, 221, 222  
 molecular, 262  
 phosphorous, 268  
 superficialis, 268
- Necrotic alteration, 296  
 changes in bone, 267
- Needles, 33  
 tarnishing of, in living muscle, 48
- Negri bodies, 419
- Nematodes, 321
- Neoplasms, malignant, of cervix uteri, 211
- Nephritis, acute, 458  
 diffuse, in scarlet fever, 304  
 in typhoid fever, 318  
 as cause of death, 408  
 chronic, 407  
 diffuse, 223  
 epithelial, 199  
 fibrous, 199  
 interstitial, 199, 201  
 parenchymatous, 199, 202
- Nephrolithiasis, see Calculi, renal.
- Nerium, toxicology of, 424
- Nerve, ischiatic, section of, 51
- Nerves, spinal, 243  
 sympathetic, hemorrhage into, 112
- Nervous form of influenza, 299  
 irritability, loss of, how determined,  
 47  
 system, central, action of poisons on,  
 419
- Nervous system, central and peripheral:  
 alcoholism, 427  
 fixatives for, 338  
 other diseases of, 453
- Nettles, toxicology of, 422
- Neuralgia, synonyms of, 453
- Neurofibroma of brain, 255
- Neurogenous arthritis, 264
- Neuroglia fibres, 255, 339
- Neuroretinitis, 223
- Neuroses, as cause of death, 408
- New-born, examination of, post-mortem,  
 275, 444  
 lymphotoxaemia in, 279  
 post-mortem examinations of, xvii  
 p. 275  
 secretion in mammary glands of, 279  
 urine of, study of, 278
- New growths, see also tumors of various  
 parts, 21
- Nicotine, toxicology of, 421, 422, 423, 424
- Nipple, accidental vaccination of, 61  
 infection of, 61  
 Paget's disease of, 61, 324  
 supernumerary, 61, 81
- Nissl's staining methods, 234, 241, 341
- Nitric acid as a fixative of tissue, 336  
 toxicology of, 422, 423, 426
- Nitrite, amyl, toxicology of, 425  
 sodium, toxicology of, 423, 425
- Nitrobenzol, toxicology of, 21, 411, 423,  
 432
- Nitrogen dioxid, toxicology of, 422
- Nitroglycerin, toxicology of, 423
- Nodular arteriosclerosis, 135
- Nodules of sarcoma in omentum, 159
- Noma, a form of diphtheritic gangrene.
- Nomenclature of diseases and causes of  
 death, international system of, xviii, p.  
 448
- Non-venereal diseases of genital organs  
 of male, 459
- Nose, abnormal shapes of, 68  
 absence of, 68  
 deviation of septum, 68, 141  
 diseases of, 68, 70, 128, 141, 142, 252,  
 296, 297, 299, 308, 405, 427, 444  
 malformations of, 141  
 saddle-back, 68  
 septum of, deviation of, 141  
 tumors of, 141
- Note-book, 23
- Notes to be made during postmortem, 23  
 dictation of, 22  
 mistakes in, liability of, 23  
 value of, 23
- Note-taking at autopsy, 22
- Nothnagel's method, 235
- N-rays in diseases of cord, 404
- Nuck, canal of, ovary in, 69
- Nutmeg poisoning, toxicology of, 433
- Nutrition and weight, 59
- Nux vomica, toxicology of, 426

## O

- Obducent in medicolegal postmortems, 401, 402  
 Obduction, 2  
 Obesitas cordis, 129  
 Objects of medicolegal postmortems, 401  
   used for comparison, 18  
 Obligations of a patient to physician, 398  
 Obliterative appendicitis, 172  
   arthritis, 250  
 Obstructive endarteritis, 136  
 Occlusion or absence of urethra, 69  
 Occupations, inflammations of skin in cer-  
   tain, 63  
 Ochronosis, 223, 264  
 Odor, characteristic, of certain diseases, 21  
   from postmortem, to remove, 10, 40  
   of acetone in diabetes, 21  
     alcohol from brain, 21  
     ammonia, 21  
     amyl nitrite, 21  
     aromatic oil, 418  
     bacterial growths, 21  
     Bacterium coli commune, 21  
     benzene, 418  
     bromin, 21  
     bromoform, 21  
     camphor, 418  
     cancerous ulcerations, 21  
     carbolic acid, 21, 418  
     certain poisons, 21  
     chloral, 418  
     chlorin, 21  
     chloroform, 21  
     chlorphenol, 21  
     creosote, 21  
     drugs, 21, 418  
     ether, 21  
     ethereal oils, 21  
     formalin, 21  
     hydrochloric acid, 418  
     hydrocyanic acid, 21  
     iodin, 21  
     iodoform, 21  
     measles, 21  
     mercaptan, 21  
     methyl alcohol, 21  
     nicotine, 418  
     nitrobenzol, 21  
     opium, 21  
     organs, 21  
     paraldehyde, 21  
     phosphorus, 21  
     poison, detection of, on exposing  
       brain, 419  
     pus in leucocythæmia, 21  
     smallpox, 21, 306  
     sulphuretted hydrogen, 418  
     tellurium, 21  
     turpentine, 418  
     uræmia, 21  
   post-mortem, 10, 286
- Edema, 20  
   complicating croupous bronchitis, 145  
   of glottis, 423, 426, 448  
     larynx, 142  
     lungs, 408, 423  
     purulent, 145  
     vulva, 69  
   pulmonary, 149  
 Edematous conditions, fixatives for, 340  
   infiltration, post-mortem, 50  
   laryngitis, 142  
 Esophageal epithelium, 189  
 Esophagus, abscesses of, 190  
   dimensions and weight of, 366  
   diseases of, 99, 112, 190, 295, 297, 310,  
     312, 426, 457  
   diverticula of, 190  
   examination of, 14, 15, 96, 188, 189,  
     442, 444  
   in animals, dissection of, 384  
   measurements of, 189, 366  
   obstructions of, 112  
   peptic ulcers of, 190  
   removal of, 15, 96, 110, 189, 442  
     in animals, 381  
   stricture of, 190, 227  
   syphilitic ulcers of, 190  
   tongue, trachea, and adjacent struct-  
     ures, removal in one piece and sub-  
     sequent examination of, 110  
   tuberculosis of, 312  
   tuberculous ulcers of, 190  
   tumors of, 112, 190  
   typhoid ulcers of, 190  
   ulcers of, 190  
   weight of, 366  
 Office of coroner, 1  
 Ohlmacher's solution for fixation of tis-  
   sues, 334  
 Oidium albicans, 310  
 Oil, anilin, toxicology of, 421  
   croton, toxicology of, 422, 423, 424  
   curcas, toxicology of, 422  
   ethereal, of mustard, toxicology of,  
     422  
   of turpentine, effect on urine, 424  
   poley, toxicology of, 423  
 Olygæmia, 116, 163  
 Omentum and peritoneum, fixation of, 337  
   appearance of, in peritonitis, 88  
   diseases of, 89, 160, 220, 317  
   examination of, 14, 15, 88, 159, 442  
   fat in, 89  
   fish-bone in, 20  
   hernia of, 89, 159  
   miliary tubercles in, 159  
   nodules of sarcoma in, 159  
   openings in, 89  
   removal of, 159  
   situation of, 88  
   tumors of, 89, 160  
   volvulus of, 159  
 Omphalomesenteric duct, 88



- Paralysis complicating scarlatina, 448  
 diver's, see Caisson disease, 249  
 general, synonyms of, 453
- Paralytic dementia, 248
- Paramidophenol, effect on urine, 425
- Paraphenyldiamin, toxicology of, 412
- Parasitic hæmoptysis, 321
- Parasites, diseases due to, xxi. p. 319  
 double, 67  
 in bladder, 207  
   blood, 121  
   bones, 274  
   kidneys, 202  
   liver, 220  
   lungs, 151  
   muscles, 83  
 intestinal, synonyms of, 457  
 in ureters, 205  
 of brain, 255  
   liver, 220
- Parathyroids, discovery of, 96  
 internal secretion of, 97  
 number of, 96  
 situation of, 96  
 structure of, 96  
 tumors of, 97
- Paratyphoid fever, 169  
 infections, 319
- Parchment-like spots, 65  
 due to drying, acids or alkalies, 65
- Parenchymatous myocarditis, 83, 130  
 myositis, 83  
 nephritis, 199, 202  
   chronic, 203  
   subacute, 202
- Parotid gland, 260, 269, 301, 302, 306, 366, 439
- Parotitis, 306
- Patella, removal of, 444
- Pathogenic micro-organisms, media for  
 isolation of, 350, 351  
 organisms, danger from, 6
- Patient and physician, contract between, 397  
 obligations of, to a physician, 398  
   physician to, 397
- Pectus carinatum, 71
- Pediculus capitis, 319  
 corporis, 319  
 getting rid of, at postmortem, 59  
 pubis, 319
- Pelades, 461
- Pellagra, 351, 418, 449  
 toxicology of, 421, 425, 433
- Pelvic organs, 183  
 in postmortem, 442  
 removal of, in animals, 385
- Pelvimeter, 35
- Pelvis, exenteration of, in animals, 385  
 female, characteristics of, 59
- Penal offence to conceal death of illegitimate child, 55
- Penicillium brevicaulis, 21
- Penis, absence of, 68  
 arteriosclerosis of, 68  
 atresia of, 68  
 atrophy of, 68  
 cancer of, 68  
 diseases of, 68, 70  
 elongation of, 68  
 erection of, 68  
 examination of, 68  
 fissure of, 68  
 gangrene of, 68  
 hypertrophy of, 68  
 imperfect formation of, 68  
 phimosis of, 68  
 removal of, 185, 186  
 scars on, syphilitic, 68  
 tumors of, 68, 134
- Pennsylvania Hospital, Ayer Clinical  
 Laboratory of, 110
- Pentastoma of bladder, 207  
 heart, 134
- Pentastomum denticulatum of spleen, 164
- Pepper dust, toxicology of, 422
- Peptic glands, examination of, 444  
 ulcer, 169  
 ulcers of œsophagus, 190
- Percentage of water in foetus and adult, 360
- Percussion, pathologic conditions revealed by, 66
- Perforating ulcer of foot, 248
- Perforation, by poison, 426, 427  
 due to ulcer of stomach, 87  
 in typhoid fever, 319  
 of intestines, 164
- Performance of autopsies in medicolegal cases, Prussian regulations for, xxvii. p. 436
- Periarteritis nodosa, 139  
 syphilitic, of pancreas, 222
- Periarthritis, 261
- Peribronchial gland, caseating, 134
- Peribronchitis, 146
- Pericarditis, 99, 144  
 complicating pneumonia, 99, 153  
 rheumatic fever, 303  
 scarlatina, 448  
 gonorrhœal, 297  
 productive, 99  
 suppurative, 99  
 synonyms of, 454  
 tuberculous, 99
- Pericardium and heart, injuries to, 99  
 changes in, at autopsy, 19  
 diseases of, 95, 98, 99, 144, 146, 151, 153, 174, 291, 295, 297, 299, 302, 303, 304, 311, 323, 445  
 erosion of, in suppuration of, 99  
 examination of, 13, 14, 15, 98, 441  
   in animals, 384  
 heart, blood, blood-vessels, and lymph-vessels, viii. p. 113  
 in animals, examination of, 384

- Pericardium, method of opening, 98  
 normal amount of fluid in, 98  
 appearance of, 99  
 tumors of, 99
- Perichondritis, 143
- Perilymphangitis, 153
- Perinephritic abscess, 203
- Period of maturity in intra-uterine gestation, 405, 406, 444
- Periosteal sarcomata, 274
- Periosteum, conditions of, in necrosis, 268  
 in rickets, 272
- Periostitis, 268, 271, 281
- Peripheral nerve-fibres, fixative for, 339  
 diseases of, 248, 250, 254, 292, 297, 299
- Perithelioma of brain, 255
- Peritoneal fluid, specimen of, from inoculated animals, 163
- Peritoneum, actinomycosis in, 159  
 cancer of, 160  
 cysts in, 159  
 diseases of, 84, 85, 86, 89, 132, 144, 153, 160, 161, 165, 174, 189, 190, 217, 220, 291, 311, 317, 323, 411, 443  
 endothelioma in, 159  
 examination of, 14, 15, 84, 159, 160  
 fibromata in, 159  
 filaria in, 159  
 lipomata in, 159  
 mechanical irritation of, 3  
 method of examination of, 88  
 pulmonary tissue in, 159  
 tuberculosis of, 317  
 tumors of, 159, 160
- Peritonitis, 144, 153  
 acute general, 160  
 serous, 161  
 as a complication of typhoid fever, 448  
 cause of, 161  
 chronic, 161  
 diffuse adhesive, 161  
 diffusive adhesive, 161  
 fibrinous, 161  
 hemorrhagic, 161  
 in typhoid fever, 319  
 local adhesive, 161  
 micro-organisms in, 160  
 proliferative, 161  
 purulent, 161  
 putrid, 161  
 serofibrinous, 161  
 simple, synonyms of, 458  
 tuberculous, adhesions in, 85  
 ulcerative, 161
- Periurethral abscess from gonorrhœal infection, 297
- Permission to perform postmortem, 3  
 method of obtaining, 3  
 to remove portions of body for preservation, 5
- Pernicious anæmia, 121, 195
- Pernicious cachexia, see Malarial cachexia.
- fever, see Intermittent and Malarial fever.
- malaria, see Malarial cachexia.
- Peroxid of hydrogen, 404
- Pest, 449
- Petechial fever, see Exanthematous typhus.
- hemorrhages, 145, 192, 414, 428  
 spots, 171  
 typhus, see Exanthematous typhus.
- Petroleum spreeds, poisoning therefrom, 412
- Peyer's patches, 123, 125, 293, 295, 301, 305, 443
- Pfeiffer's bacillus, 298
- Phalloides, amanita, 418, 424
- Pharyngeal cavities, exposure of, 256  
 diphtheria, 294
- Pharynx, affections of, 141, 246, 296, 297, 308, 456  
 examination of, 15, 442  
 removal of, 110, 442
- Phenomena bearing upon future of pathology, 404
- Phenylendiamin, toxicology of, 422
- Philadelphia Hospital, postmortems at, 5  
 blank post-mortem record of, 26
- Phimosis of glans penis, 68  
 partial, of urethra, 69
- Phlebectasia, 140
- Phlebitis of veins, 139
- Phleboscclerosis, 140
- Phlegmasia alba dolens, puerperal, complications of, 460
- Phlegmon of iliac fossa, 458, 461
- Phlegmonous gastritis, 191
- Phloridzin, toxicology of, 425
- Phosphates and lime salts in bones of pregnant women, 70
- Phosphomolybic acid hæmatoxylin, 341
- Phosphorescence, as a pathologic force, 404
- Phosphorescent bacteria, 404
- Phosphoric acid, useful in axillary cellulitis, 44
- Phosphorism, toxicology in, 423
- Phosphorus, necrosis from, 268  
 toxicology of, 21, 411, 421, 423, 424, 425, 433
- Phosphotungstic acid hæmatoxylin, 341
- Phrenic nerves, exposure of, 98
- Phthisiogenic organism, 350
- Phthisis, fibroid, 311, 315, 316  
 florida, 311, 315  
 ulcerative, 311, 315
- Physician and patient, contract between, 397  
 obligations of patient to, 398  
 to patients, 397  
 practising, when intoxicated, a criminal offence, 398

- Physician, resident, 3, 6  
 Physician's certificate for return of a death, 464  
   responsibility in judicial hanging, 46  
 Physostigmine, effect on pupil after sudden death, 46  
   toxicology of, 421, 422, 423, 424  
 Pia and arachnoid, weight of, 362  
   diseases of, 244, 251, 252, 313, 434  
   removal of, 231, 241  
     in animals, 388  
 Pianese's solution, as a fixative, 337  
 Pick's myelotome, 29, 232  
 Picrates, toxicology of, 422, 423, 424  
 Picric acid, as a fixative, 327, 337  
   toxicology of, 423, 424, 433  
 Picro-acetic acid, as a fixative, 337  
 Picro-hydrochloric acid, as a fixative, 337  
 Picro-nitric acid, as a fixative, 337  
 Picro-sulphuric acid, as a fixative, 337  
 Picrotoxin, toxicology of, 421, 434  
 Pig, as a disseminator of trichinosis, 373  
 Pigeon-breast (pectus carinatum) of rhabdomyositis, 72  
 Pigmentation in chlorosis, 122  
 Pigments in bile, 214  
 Pilocarpine, toxicology of, 422, 423, 424, 426  
 Pineal gland, dimensions and weight of, 363  
   examination of, 236  
 Pins, 33  
 Pitres's method of examining brain, 235, 237  
   studying centrum ovale, 235  
 Pituitary body, 56, 70, 246, 247, 282, 363  
   removal of, through orbit, 247  
 Place at which a postmortem is made, 8  
 Placenta, diseases of, as causes of spontaneous abortion, 407  
   weight of, 359  
 Placental polyp, 209  
 Plagiocephalic skull, 361  
 Plague, bubonic, 301, 449  
   carbuncular, 301  
 Plague-corpuscle, 301  
 Plague, intestinal, 301  
   meningeal, 301  
   pneumonic, 301  
   septic, 301  
   subcutaneous hemorrhages due to, 302  
 "Plaques choriales" of sheep, 70  
 Plasma cells, fixative for, 340  
 Plaster of Paris, 289  
   for filling cranial cavity, 34  
   in the preservation of the body, 283  
 Plastic bronchitis, 145  
 Plastron, fig. 69  
 Plates, earthenware, 33  
 Platinico-acetico-osmic-acid solution, 336  
 Platinum loop, or öse, 347, 348  
 Platycephalic skull, 361  
 Plethora serosa, 115  
   vera, 115  
 Pleura, abscess of, 144  
   carcinoma of, 156  
   chondroma of, 156  
   diseases of, 95, 97, 99, 127, 140, 146, 147, 149, 153, 155, 156, 157, 1574, 190, 219, 291, 293, 295, 297, 301, 316, 416  
   endothelioma of, 156  
   examination of, 97, 108, 441  
   fibroma of, 156  
   hyatid cysts of, 156  
   interlobular, 158  
   lipoma of, 156  
   osteoma of, 156  
   sarcoma of, 156  
   teratoma of, 156  
   thickening of, as lesion in tuberculosis, 316  
   tuberculosis of, 156, 315  
   tumors of, 156  
   visceral, pathologic conditions of, 1  
 Pleural cavity, amount of fluid in, 97  
   empyema of, 156  
   examination of, 97  
   suppuration in, 156  
 Pleurisy, 156, 157  
   acute, 157  
   chronic, 157  
   dry, 157  
   diaphragmatic, 158  
   due to rheumatic fever, 303  
   encysted, 158  
   following dysentery, 174  
   gonorrhœal, 297  
   hemorrhagic, 157  
   primitive dry, 158  
   purulent, 157  
   synonyms of, 456  
 Pleuropneumonia, 151  
 Plexus, choroid, 439  
 Pneumococcal arthritis, 264  
 Pneumococci, 146  
 Pneumococcus (*Streptococcus lanceolatus*), 351  
 Pneumogastric nerves, 15, 276  
 Pneumomalacia, 150  
 Pneumomycosis, 151  
 Pneumonia, 151  
   as a complication of typhoid fever, 318, 448  
   cause of death, 408  
   bacillus of, 151  
   catarrhal, 151  
   and croupous, difference between, 154  
   chronic interstitial, 151, 152, 223  
   croupous, 151, 152, 223  
   and catarrhal, table showing difference between, 154  
   due to rheumatic fever, 303  
   fibrinous, 151, 152



- Pneumonia, hypostatic, in typhoid fever, 318  
 lobar, 151, 152  
 lobular, 151, 154, 299  
 stages of, 152  
 synonyms of, 456
- Pneumonic areas in tuberculosis, 316  
 plague, 301
- Pneumonoconiosis, 155
- Pneumopericardium, causes of, 99
- Pneumothorax, 156  
 localized, 156  
 method of detecting, 80, 98
- Pockets in walls of bladder, 206
- Podagra, 263
- Podophyllotoxin, toxicology of, 424
- Poikilocytosis, 117
- Poisons, see Toxicology of the various  
 poisons themselves.  
 classification of, 420  
 definition of, 417
- Poley oil, toxicology of, 423
- Poliomyelitis, acute anterior, 254, 351  
 from compression, 254
- Polycythemia rubra, 116
- Polydactylism, 2
- Polygnathus, 68
- Polymelus, 68
- Polyps of intestines, 171  
 rectum, 69  
 ureters, 204  
 uterus, 204
- Pons Varolii, diseases of, 313  
 examination of, 236, 439  
 fixatives for, 339  
 incising of, 236  
 removal of, 236  
 weight of, 362
- Porencephalon, 250
- Porencephaly, external and internal, 249
- Portal pyæmia, 219  
 vein, 15, 61, 139, 162, 193, 194, 196, 365
- Post, see Postmortem.
- Posterior extremity of an animal, method  
 of removing, 375  
 sclerosis, see Locomotor ataxia.
- Posthumous circulation, 49
- Post-mortem books, 25  
 changes in conjunctiva, 48  
 cornea, 48  
 color scale, method of using, 113  
 considerations, general, i, p. 1; 436  
 cooling, hastened by use of water, 49  
 rectal temperature after forty  
 hours, 49  
 time required for completion, 49  
 decomposition, 49, 52  
 digestion of stomach, 87  
 drains, 11  
 emphysema, 95  
 examination of animals, 225  
 birds, 394  
 cats, 392
- Post-mortem examination of dogs, 392  
 law permitting, 5  
 lower animals, 374  
 new-born, 275, 403  
 restricted, 280  
 ruminants, 389  
 swine, 391  
 order of, and records, 13  
 rules for, 13
- expulsion of fœtus, 46  
 instruments, and how to use them,  
 iii, p. 28; 439  
 lividity or cadaveric lividity, 49  
 note-taking, ii, p. 13  
 odor, 286  
 records, ii, p. 13; 22, 23, 24, 26, 27, 394  
 of findings in lower animals, 394  
 references, xxix, p. 465  
 report, 447  
 rigidity or death-stiffening, 50  
 room, 10  
 drains of, 11  
 latticed flooring for, 12  
 ventilating shaft for, 11
- table, 9, 11  
 measuring rod for, 11  
 portable, 9  
 revolving, 11  
 weighing body on, 11
- transportation of corpses, 437
- wounds, care of, iv, p. 38  
 caustics used in treatment of, 43  
 how caused, 42, 43  
 virulence of, 41, 42
- Postmortem, abdominal cavity in, 442  
 after electrocutions, 7  
 Anatomical Board and, 5  
 anatomy, study of, at, 3  
 appearance of parts improved after, 9  
 ascertainment of special circum-  
 stances in, 437  
 assistants at, 6  
 at hospital, 9  
 morgue, 9  
 private house, 9  
 undertaker's, 9, 10  
 Boards of Health and, 4  
 care in handling specimens removed  
 at, 6  
 cases of poisoning in, 443  
 cause of death, determined by, 16  
 in child, whether or not  
 it has breathed, 444  
 cleanliness at, 9  
 closure of body in, 445  
 comparative, xxv, p. 373  
 cranial cavity in, 439  
 death-stiffening, 50  
 decomposed bodies in, examination  
 of, 436  
 defendant represented at, 420  
 definition of, 2  
 diagnosis at, 16

- Postmortem, disinfection after, 2
- dress at, 10
  - duties of obducent in, 437
  - early, in America, 1
  - establishment of judicial, 1
  - external examination in, 438
    - of decomposed bodies, 436
    - new-born, xviii, p. 275
    - vertebræ, as medicolegal precaution, 402
  - frozen bodies in, 437
  - general considerations, i, p. 1: 436
  - historical, 1
  - illumination at, 11
  - in charitable institutions, 5
    - France, 6, 10
    - Germany, 5, 7
    - insurance cases, 4, 5
    - medicolegal cases, Prussian regulations for performance of, 436
  - internal examination in, 439
  - interval to elapse before making, 7, 436
  - intra-uterine gestation in, 444
  - kidneys in, 442
  - laboratory, upon animals, 374
  - law permitting, 15
  - legal right to perform, 3
  - Letulle's method of making, 15
  - light at, 8
  - liver in, 443
  - medicolegal, 401, 437
    - and pathologic, difference between, 402
    - definition of, 2
    - in cases of poisoning, 402
    - treatment of contents of
      - stomach, 402
      - duodenum and ileum, 402
      - each organ, 402
    - obducent in, 401, 402
    - objects of, 401
  - order of performing, 12
  - micro-organisms present at, 43, xxiii, p. 346
  - microscopic examinations in, 438
  - multiple, 11
  - neck in, 442
  - nurse at, 9
  - object of, 2
  - odor at removal of, 10, 286
  - of Bishop,
    - Brooks, Phillips, 6
    - McKinley, 6
  - in anatomical cases, 5
    - mona's operative technic of, 374
    - babies suspected of hereditary syphilis, 277
    - birds, 324
    - bones, 301
    - cats, 302
    - child, preparation for, 9
    - dogs, 302
- Postmortem on ears, 256
- eyes, 256
  - horse, 374
  - infants, 403
  - joints, 261
  - lower animals, value of, 373
  - nasopharynx, 256
  - new-born, 444
  - ruminants, 389
  - swine, 391
  - operations at, 3
  - order of examination at, 13
  - pelvic organs in, 442
  - performed in coffin, 9
    - without legal permission, 3
  - performing of, dangerous, 45
  - permission to perform, 3
    - method of obtaining, 3
  - persons present at, 6
  - physicians and their duties in, 436
  - place at which made, 8
    - for and its lighting, 437
  - portable table for, 9
  - precautions to be taken in performing, 401
  - preparation for, 9
  - prevention of rheumatism while performing, 12
  - prohibition of smoking at, 6
  - Prussian regulations for the performance of medicolegal, xxvii, p. 436
  - purpose of, 2
  - record books for, form of, 24, 26, 304
  - records of, ii, p. 13
  - references to, xxix, p. 465
  - removal of entire organs at, 15
    - odor at, 10
    - specimens at, 5
  - repetition of, 430
  - report of, 447
  - resident physicians making, 3, 6
  - restricted, xix, p. 280
  - rigidity in, 50
  - room for, 11
    - fittings of, 11
  - rules for performing, 13
  - scales at, 10
  - scheme for record of, in lower animals, 304
  - skull at, 10
  - small and large intestines in, 443
  - societies for performing, 4
  - spinal column and cord in, 440
  - spinal cord in, 442
  - spleen in, 442
  - stomach and duodenum in, 443
  - stool for instruments at, 12
  - study of normal structures in, 2
    - anatomy at, 3
  - synonyms of, 12
  - substitute for daylight at, 8
  - substitutes in, 436
  - suggestions at, 9

- Postmortem, surgical procedures at, 3  
 synonyms of, 2  
 technic of, 437  
 thoracic cavity in, 440  
 those present at, 6  
 time after death before making, 7, 436  
     for performance of, 7  
     of completing a, 7  
 to be made before the body is dressed, 8  
 transportation of corpses for, 437  
 two main divisions of, 438  
 undertaker at, 6  
 value of, in accident cases, 2  
 veterinary, xxv, p. 373  
 where made, 9  
 wounds in, treatment of, iv, p. 38
- Post-partum hemorrhage, 128
- Potassium, bichromate of, as a fixing fluid, 331  
 toxicology of, 8  
 binoxalate, toxicology of, 424, 426  
 cantharidate, effect on urine, 426  
 cantharidinate, effect on urine, 424  
 chlorate, toxicology of, 60, 203, 423, 425, 434  
 cyanid, toxicology of, 411, 419, 421
- Pott's disease, 72, 198, 450  
 complications of, 450
- Powder markings, 409
- Powerful convulsive remedies, toxicology of, 421
- Pregnancy, accidents of, 460  
 ectopic, ovarian, tubal, 207
- Premature interments, 46  
 belief in, in Munich, 46  
 labor, case of, 279
- Preparation of inoculum, 353  
 material in a sterile bulb for inoculation, 353  
 smear, 347
- Preservation of body, xx, p. 286  
 disembowelled body of child, 276  
 muscle-fibres, 327  
 serous and mucous membranes, 327  
 tissues, by alcohol, 338  
     cedar oil, 338  
     nitrate solution, 338  
     sugar formation in liver, 338  
     for macroscopic purposes, 326  
     microscopic study, 326  
     general considerations in, 342
- Preservative injection for embalming bodies, 287  
 fluids, 342
- Primitive dry pleurisy, 158
- Principles, active, of digitalis, toxicology of, 421
- Probes, 32
- Processes, acute infectious, fixative for, 338
- Prognathous skull, 361
- Progressive locomotor ataxia, synonyms of, 452  
 muscular atrophy, 253
- Prolapse of rectum, 69  
 vagina, 211
- Proliferative endarteritis, 136  
 chronic peritonitis, 161
- Prosopothoracopagus, 67
- Prostate, abscesses of, 212  
 cystic formation in, 212  
 dimensions of, 372  
 diseases of, 205, 212, 297, 459  
 enlargement of, 212  
 examination of, 14, 15, 186, 212, 442  
 inflammation of, 212  
 measurements of, 372  
 removal of, 185, 186  
 tumors of, 212  
 weight of, 372
- Prostatitis, gonorrhœal, 297
- Prostration, 463
- Proteus, see *Bacterium vulgare*, 350
- Protocol of postmortem 22, 445, 446  
 in animals, 394
- Protozoa, diseases due to, 351
- Protozoön of smallpox, 305
- Provisional opinion of obducent, at close of postmortem, 446
- Prussian regulations for medicolegal autopsies, 7, xxvii, p. 436
- Psammoma of brain, 255
- Pseudencephalos, 67
- Pseudoarthrosis, 266
- Pseudoleukæmia, 124
- Pseudomembranous angina, see Diphtheria and Croup.
- Pseudomuscular hypertrophy, 253
- Psilosis, 306
- Psoas muscles, 14, 198
- Psorospermiasis, 323, 324
- Psorosperms in liver, 323
- Ptomain and toadstool poisoning, toxicology of, 434
- Ptoxis, acquired or congenital, 73
- Puerperal accidents, other, 460  
 diseases of the breast, 460  
 eclampsia, 270  
     and albuminuria, 460  
     latest theory of cause of, 278  
 hemorrhage, 460  
 scarlatina, see *Scarlatina*.  
 septicæmia, 460
- Pulex irritans, 319  
 penetrans, 319
- Pulmonary abscess, 213  
 apoplexy, 456  
 arteries, 15, 16, 102, 104, 105, 110, 148, 150, 223, 441  
 congestion, 456  
     of typhoid fever, 448  
 embolism, 148  
 œdema, 149



- Pulmonary orifices of heart, dimensions  
 of, 365  
 osteo-arthropathy, hypertrophic, 247  
 roots, 15, 110, 441  
 thrombosis, 148  
 tissue below diaphragm, 91  
   in peritoneum, 159  
 valves, 15, 133, 365  
 veins, thrombosis of, 148  
 vessels, examination of, 110  
 Pulse, absence of, on palpation, 47  
 Pupils, contraction of, after death, 48  
   dilatation of, after death, 48  
   in death by electricity, 413  
   measurement of, 74  
   Ripault's test on, 48  
 Purpura, 125  
   cachectic, 126  
   forms of, 126  
   hæmorrhagica, 126  
   Henoch's, 126  
   iodic, 126  
   mechanical, 126  
   morbid anatomy of, 126  
   neurotic, 126  
   rheumatica, 126  
   simplex, 126  
   toxic, 126  
 Purpuric blotches mistaken for bruises,  
 126  
 Purulent and septicæmic infection, 449  
   bronchitis in typhoid fever, 319  
   œdema of the lung, 145  
   osteitis, 270  
   periostitis, 268, 271  
   pleurisy, 157  
   vulvitis (gonorrhœal), 69  
 Pus in post-mortem wounds, treatment of,  
 43  
   or purulent conditions, fixatives for,  
   340  
 Putrefaction, by poison, 427  
   early manifestations of, 49  
   in sunstroke, 414  
   most positive sign of death, 49  
 Putrid bronchitis, 144, 145  
 Pyæmia, chronic specific, 297  
   portal, 219  
 Pyæmic hepatitis, 219  
 Pyelitis and pyelonephritis, 203  
 Pyelonephritis, 203  
 Pylephlebitis, 149  
 Pyloric stenosis, 190  
 Pylorus, 190  
 Pyopagus, 67  
 Pyopneumothorax, 150  
 Pyosepticæmia, 219  
 Pyothorax, 156  
 Pyramid, Lalouette's, 96  
 Pyramidon, toxicology of, 425  
 Pyrogallic acid and caustic soda for ab-  
 sorption of oxygen, 353  
 Pyrogallol, toxicology of, 423, 425
- Q**
- Quartan malaria, see Intermittent fever  
 and Malarial cachexia.  
 Questions, medicolegal, 400, 409  
 Quillaja bark, toxicology of, 422  
 Quinine, toxicology of, 422, 425  
   useful in axillary cellulitis, 44
- R**
- Rabic tubercle, 298  
 Rabies, 297, 351, 449  
   method of diagnosing, 298  
 Race, 55  
 Racial characteristics, 58  
 Rachipagus, 67  
 Radio-activity of dead body, 404  
 Ranunculus acris, toxicology of, 422  
   sceleratus, toxicology of, 422  
 Rape, evidence of, 54  
 Rarefying osteitis, 269  
 Rash of smallpox, 305  
   rose, 351  
 Raspatory of Chiara, 31  
 Rat, disseminator of bubonic plague, 373  
   used for inoculation, 353  
 Ratio of weight of organs to body, xxiv.  
   p. 357  
 Ray-fungus, 290  
 Raynaud's disease, 254  
 Razor, 29  
 Reaction, agglutinative, diseases diagnosed  
   by, 115  
   for diagnosis of human blood, 118  
   in blood and other substances, 118  
   semen, 417  
   amyloid, of kidney, 179  
   glycogenic, in blood, 120  
   of diabetic blood, 224  
   organs, 20  
 Receptaculum chyli, 197  
 Von Recklinghausen's disease (hæmo-  
 chromatosis), 254  
 Records, post-mortem, 13, 22, 23, 24, 26, 27  
   of findings in lower animals, 304  
   necropsy, scheme for, in  
   lower animals, 304  
 Rectal enema, reversed peristalsis after,  
 169  
 Rectum, congenital polypi of, 69  
   diseases of, 69, 305, 306, 308, 312  
   examination of, 15, 69, 166, 442  
   fistulæ of, 69  
   hemorrhoids of, internal and external,  
   69  
   imperforate, 279  
   removal of, 184, 442  
   in animals, 377  
   prolapse of, 69  
   tumors of, 69  
 Recurrent fever, see Relapsing fever.  
 typhus, see Relapsing fever.

- References, medicolegal, xxix, p. 474  
 post-mortem, xxix, p. 465
- Refrigerator box, 10, 11
- Regional landmarks, 17
- Regulations of court, 397  
 Prussian, for performance of medicolegal postmortems, xxvii, p. 436
- Regurgitation, tricuspid, 133
- Relapsing fever, 302
- Relation between cortex and medulla of kidney, 179
- Remedies, internal, toxicology of, 422
- Remittent fever, see Intermittent and Malarial fever.
- Removal of abdominal contents of animals, 378  
 dog, 393  
 brain, almost intact, 281  
   in animals, 388  
 cerebellum, medulla oblongata, and pons Varolii, 236  
 child's brain, 276  
 extremities of an animal, 375  
 female genitalia, 183  
 head of animal from trunk, 386  
 hide of an animal, 375  
 ileum of animals, 378  
 jejunum of animals, 378  
 kidneys of animals, 380  
 liver of animals, 380  
 male organs of generation, 185  
 spinal cord in animals, 389  
 sternum, 93  
 stomach of animals, 380  
 thoracic contents of animals, 384  
 vertebral column, 242
- Renal, see Kidney.
- Report, final, of postmortem, 445, 447  
 form of, in medicolegal post-mortem examinations, 403
- Reptiles in comparative pathology, 374
- Resident physicians, 3, 6
- Resolution of lung, 153
- Respiration, absence of, how determined, 47  
 time of cessation of, 47  
 Winslow's test for, 47  
 X-rays used to detect any movement of, 47
- Respiratory apparatus, diseases of, ix, p. 141; 455, 456
- Responsibility of physician, 46, xvii, p. 397
- Restoration and preservation of body, xx, p. 283
- Restricted post-mortem examinations, xix, p. 280
- Retina, examination of, 257  
 fixatives for, 339
- Retractors, hooked, 32
- Retroperitoneal glands, examination of, 14, 196  
 tumors of, 196
- Retropharyngeal abscess, 292
- Reversed peristalsis, 169
- Rhabdomyomata of heart, 134  
 kidney, 204
- Rhachioschisis, 249
- Rhachiotome, Luer's, 29
- Rhachipagus, 67
- Rhachitic cirrhosis of liver, 217  
 rhinitis, 272
- Rhachitis, 71, 72, 272
- Rheumatic arthritis, 263  
 fever, 303
- Rheumatism, 12, 303, 351, 451
- Rheumatoid arthritis, 262, 266
- Rhinitis, 141, 142, 272
- Rhinoliths in nasal passages, 141
- Rhinoscleroma, 141
- Rhubarb root, effect on urine, 425
- Rhus toxicodendron, toxicology of, 422
- Ribs, caries of, 81  
 cervical, 68  
 diseases of, 68, 72, 93, 94, 99, 246, 290  
 examination of, 68, 72, 81, 94  
 of animals, method of cutting, 377  
 tumors of, 273, 274
- Rice bodies, 261
- Rice-water, 293
- Richardson blue, 345
- Ricin, toxicology of, 434
- Rickets, 71, 72, 272
- "Riders" bones, 83
- Rigidity, post-mortem, or death-stiffening, 50, 51, 52, 281  
 absent in immature fœtus, 52  
 after high mental tension, 51  
 ante-natal, 404  
 changes in muscle sheaths found in, 51  
 delayed by rapid cooling of body, 52  
   section of one ischiatic nerve, 51  
 duration of, 51  
 in cachectic subjects, 50  
   cholera, 51  
   chronic alcoholism, 50  
   heat-stroke, 50  
   scarlet fever, 304  
   spinal poisons, 51  
   strychnine poisoning, 50, 51  
   suffocation, 50  
   suicide, 51  
   tetanus, 51  
   veratrum viride poisoning, 51  
 instantaneous, 51  
 location of beginning of, 51  
 of soldiers on battle-field, 51  
 overcome by force, 52  
 position of hands in, 51  
 post-mortem, 51, 52  
 reaction of muscles in, 52  
 time of beginning of, 51  
   completion of, 51
- Rigor mortis, 50, 51, 52, 281

- Ripault's test of pupils, 48  
 Robin's injecting fluid, 344  
 Rokitsky's 30,000 postmortems, 7  
 Rose rash, 331  
 Rötheln, 303  
 Rough on rats, 429  
 Roux's inoculation syringe, 354  
 Rubber gloves, 34, 343  
 Rubella, 303  
 Rubeola, 300  
 Rudimentary uterus, 209  
 Rue, toxicology of, 424  
 Rule for determining weight of cerebrum  
     from height of body, 363  
     steel, in measuring an organ, 360  
     to convert grammes into ounces and  
     grains, and *vice versa*, 357  
     two-foot, 32  
 Rules for determining lunar months of  
     embryo, 358  
 Ruminant, nasal and cranial cavities in,  
     lines used in sawing, 390  
 Ruminants, postmortems on, 389  
 Rupia, 307  
 Rupture of aneurism, 138  
     bladder, 206  
     heart, spontaneous, cause of, 129  
     tubal pregnancy, as cause of  
         death, 460  
     uterus, as cause of death, 460  
     varicose veins of vulva, 69  
     spontaneous, of heart, 129
- S**
- Sabina, toxicology of, 424  
 Saccular aneurism, 138  
 Sacculated aneurism treated by electroly-  
     sis, 139  
 Saddle-back nose, 68  
 Sago spleen, 164  
 Salicylic preparations, toxicology of, 422  
 Salivary calculi, 260  
     glands, diseases of, 66, 259, 294  
     examination of, 66, 260, 442  
     tumors of, 260  
     weight of, 366  
 Salivation, drugs producing, 423  
 Salpingitis, 208, 297, 314  
 Salt water, drowning in, 416  
 Sand, false, in intestines, 170  
 Sand-flea, 319  
 Santonica seeds, 425  
 Santonine, toxicology of, 421, 425  
 Saponiferous substances, toxicology of,  
     425  
 Saponine, toxicology of, 423  
 Sarcinae, 351  
 Sarcoma, as cause of death, 408  
     in larynx, 143  
     nasal passages, 141  
     metastatic, 75  
     nodules of, in omentum, 159  
 Sarcoma of bladder, 205  
     bone, 274  
     brain, 255  
     choroid, 75  
     ciliary body, 75  
     eye, 16  
     heart, 134  
     intestines, 171  
     kidney, 204  
     liver, 16, 219  
     mediastinum, 146  
     periosteum, 274  
     pituitary body, 247  
     pleura, 156  
     spleen, 164  
     tonsils, 143  
     urethra, 70  
     vulva, 69  
     spindle-celled, in mediastinum, 146  
         of lungs, 156  
 Sarcophages, 320  
 Satchel, contents of, for post-mortem  
     work, 35  
 Saturnism, toxicology in, 421, 422, 432  
 Sausage and fish poisoning, toxicology of,  
     422, 423  
 Sawdust in preservation of body, 283  
 Sawing skull, 226  
 Saws, 29  
     butcher's meat-saw, 29  
     Hey's, for opening skull, 29  
     Luer's double rhachiotome, 29  
     metacarpal, 29  
 Scalds and burns, extent of injuries in,  
     412, 463  
 Scale, moisture, 114  
     post-mortem color, method of using,  
         113  
     Tallqvist's blood-color scale, 113  
 Scales, 10, 32  
 Scalp, diseases of, 97, 295  
     examination of, 14, 225  
     method of replacing, 285  
 Scalpel for solids in smear preparations,  
     348  
 Scalpels, 29  
 Scarification on dead subject, 47  
 Scarlatina, synonyms of, 448  
 Scarlatinal cirrhosis of liver, 217  
 Scarlatinous angina, *see* Scarlatina.  
 Scarlet fever, 303, 351  
     complications of, 448  
     to be distinguished from Duke's  
         fourth disease, 300  
     virus of, toxicology of, 424  
 Scars, 64, 65  
     from hypodermics, 65  
     in death from electricity, 65  
     location of, 65  
     of kidneys, 179  
     significance of, 64  
     surgical, 64, 65  
     syphilitic, of penis, 68



- Scheme, author's, for classification of  
   poisons, 420  
 Schweinfurth-green, toxicology of, 423  
 Scilla, toxicology of, 424  
 Scirrhus cancer of lungs, 156  
   stomach, 190  
 Scissors, 31  
 Sclera, 75  
 Scleroderma, circumscribed, 305  
   diffuse, 305  
   organism of, 293, 350  
 Sclerosis, arterial, 223  
   disseminated, 254  
   insular, 254  
   posterior, see Locomotor ataxia.  
 Scoliosis, 198  
 Scopolamine, toxicology of, 422, 423, 424,  
   426  
 Scorbutus, 452  
 Scrofula, 450  
 Scrotum, absence of, 69  
   atrophy of, 69  
   cleft, 69  
   contraction of, 69  
   diseases of, 69, 140  
   examination of, 69, 73, 444  
   hematocele of, 69  
   hernia of, 69  
   hydrocele of, 69  
   hypertrophy of, 69  
   lymphangitis of, gangrenous, 69  
   tumors of, 69  
 Scurvy, 126  
 Seat-worms, odor in, 170  
 Sea-weed in body, 283  
 Sectio cadaveris, see Postmortem.  
   anatomica, see Postmortem.  
 Section, see Postmortem.  
   of oral cavity and cervical organs in  
   animals, 384  
 Sectioning brain, methods of, 234, 235, 237  
   cranium in animals, 386  
 Sections, frozen, preparation of, 12  
   of organs for preservation, 327  
   tumors for preservation, 327  
 Sedative alkaloidal poisons, 421  
   poisons, 421  
 Seeds, santonica, effect on urine, 425  
 Segmentation of heart, 129  
 Selection of culture-media, 349  
   site for inoculation, 355  
 Selenium, toxicology of, 426  
 Semen, 417  
   biologic blood-test for, 417  
   Florence test for, 417  
   precipitin method for, 417  
 Semilunar cartilages, dislocations of, 266  
   ganglia, 181  
 Seminal vesicles, dimensions of, 372  
   Nauwerck's method of finding,  
   187  
 Senile arteriosclerosis, 135  
   atrophy, 129  
 Senile debility, synonyms of, 462  
 Senna leaves, effect on urine, 425  
 Septicæmia, puerperal, synonyms of, 460  
 Septic plague, 301  
 Septum of nose, deviation of, 68, 141  
 Serous and mucous membranes, preser-  
   vation of, 327, 337  
   membrane, appearance of, when dis-  
   eased, 85, 97  
   when normal, 97  
   tuberculosis of, 317  
 Serum blood-test, 118  
 Setons, scars from, 64  
 Sex, 55  
   in hermaphrodites, 55  
 Shaft, ventilating, 11  
 Shape of organs, 16, 18  
   skull, 360  
 Shavings in preservation of body, 283  
 Sheaths, 272  
 Sheep, "plaques choriales" of, 70  
   post-mortem examination of, 391  
 Shiga, bacillus of, 172  
 Shock, death due to, 409  
 Shortened mesentery, 160  
 Shortening of limb, 56  
 Shoulders, circumference of, 56  
 Siderosis, 155  
 Sigmoid flexure, redundancy of, 20, 160  
 Signs of death, 46  
   degeneration, 70  
 Silicosis, 155  
 Silver nitrate solution for post-mortem  
   wounds, 43  
   Hume's intravenous injection of,  
   44  
   toxicology of, 411, 422, 423, 434  
 Single monsters, 67  
 Sinus of Valsalva, aneurisms of, 131, 138  
 Sinuses, accessory, exposure of, in ani-  
   mals, 389  
   longitudinal, 304, 439  
 Site of inoculation in animals, 354  
 Sitotaxismus, 418  
 Size, how estimated, 18  
 Skeleton, 58  
   peculiarities of female, 58, 59  
   male, 58, 59  
 Skin and its appendages, v. p. 59: 296, 461  
   acne of, 63  
   angioneurotic œdema of, 64  
   atrophy of, 63  
   bedsores of, 63  
   blots of, 63  
   bronzing of, 16, 62  
   color of, 59, 60, 62, 122  
   dirt on, 59, 61  
   diseases of, 44, 53, 60, 62, 63, 64, 65,  
   66, 116, 122, 123, 125, 126, 128, 132,  
   140, 252, 290, 291, 294, 296, 299, 300,  
   302, 305, 306, 307, 317, 318, 319, 320,  
   324, 411, 412, 414, 426, 427, 428, 431,  
   433, 438

- Skin, epithelioma of, 63  
 examination of, 53, 59, 60, 61, 62, 63,  
 64, 65, 66, 405  
 fixatives for, 340  
 in actinomycosis, 63  
   Addison's disease, 60  
   albinism, 60  
   anæmia, 62  
   anthrax, 63  
   arsenical poisoning, 63  
   blastomycosis, 63  
   Bright's disease, 64  
   chimney-sweeper's cancer, 63  
   chloasma, 62  
   chlorosis, 60, 122  
   cholera Asiatica, 203  
   dermatitis herpetiformis, 63  
   dermatomyositis, 63  
   diabetes, 64  
   drowning, 66  
   ecthyma, 63  
   eczema, 63  
   ergotism, 64  
   erythema multiforme, 62, 63  
   furunculosis, 63  
   gangrenous lymphangitis, 64  
   herpes, 62  
   ichthyosis, 63  
   impetigo, 63  
   keloid, 64  
   keratosis, 63  
   leprosy, 60, 63  
   lichen, 63  
   locomotor ataxia, 63, 64  
   lupus vulgaris, 63  
   measles, 62, 63  
   mercury poisoning, 62  
   miliaria, 63  
   moles, 62  
   myxœdema, 63  
   nævus pigmentosa, 62  
   pediculosis, 62  
   pelioid rheumatica, 62  
   pemphigus, 63  
   pernicious anæmia, 60, 63  
   pityriasis, 62, 63  
   planus, 63  
   prurigo, 63  
   psoriasis, 63  
   puerperal fever, 64  
   purpura, 62, 63  
   rheumatism, 62  
   ringworm, 63  
   scabies, 63  
   scarlet fever, 63  
   scleroderma, 63  
   scrofulosis, 63  
   scurvy, 64  
   seborrhœa, 63  
   smallpox, 62, 63  
   snake poisoning, 62  
   syphilis, 62, 63  
   tinea sycosis, 63
- Skin in tuberculosis, 318  
 typhoid fever, 318  
 typhus fever, 62  
 uræmia, 64  
 varicella, 63  
 vitiligo, 60  
 xanthoma, 64  
 parchment-like spots of, 65  
 tumors of, 62, 63, 64, 66  
 ulcers of, 63  
 vesicles of, 62, 63
- Skull, altitudinal index of, 360  
 base of, examination of, 233  
 brachycephalic type of, 361  
 cephalonic type of, 361  
 characteristic measurements of, 361  
 clamp, 32  
 clinoccephalic type of, 361  
 diseases of, 72, 226, 247, 251  
 dolichocephalic type of, 361  
 examination of, 14, 58, 72, xiii, p. 8  
 fracture of, 1, 408, 411  
 French method of opening, 228  
 hydrocephalic type of, 361  
 measurements of, 229, 361  
 methods of restoring and preserving, 283, 284  
 microcephalic type of, 361  
 opening of, with a hammer, 228  
 orthocephalic type of, 361  
 orthognathous, 361  
 oxycephalic type of, 361  
 pachycephalic type of, 361  
 pathologic types of, 361  
 plagiocephalic type of, 361  
 platycephalic type of, 361  
 prognathous type of, 361  
 rhachitic shape of, 72  
 sawing of, 226  
 shape of, 360  
 sphenocephalic type of, 361  
 sutures in restoring, 283  
 thickness of, 229  
 trigonocephalic type of, 361  
 trochocephalic type of, 361  
 tumors of, 273, 274  
 weights of contents of, 360
- Skullcap, see Calvarium, 230
- Sleeping sickness, 324
- Sloughing, as a complication of typhoid fever, 448
- Small intestine, see Intestine, small.
- Smallpox, 16, 48, 64, 305, 351, 448  
 bronchopneumonia, as a complication of, 306  
 cause of sudden death, 16  
 complications of, 448  
 hemorrhagic, 305  
 odor of, 305
- Smear culture, 352  
 preparations, 247  
 fixing of, 348
- Smell, sense of, development of, 21

- Smokeless powder, indistinct markings made by, 409
- Smoking, prohibition of, at postmortem, 6
- Snake venom, toxicology of, 434
- Soapstone, ground, for preserving rubber gloves, 34
- Society, American Anthropometric, 4
- Sodium chlorid and bichlorid of mercury solution, 334
- nitrite, toxicology of, 423, 425
- thiosulphate of, toxicology of, 424
- urate, 207
- Soft chancre, 450
- Solanine, effect on urine, 425
- Solanus pseudocapsicum, toxicology of, 419
- Soldiers, post-mortem rigidity in, 51
- Solid material, method of removing from interior of an organ, 347
- Solids in smear preparations, 348
- Soluble cells, preservation of, 328
- Solution, bichlorid-tablet, in preservation of tissues, 326
- Sore, syphilitic, 349
- Spanish fly, toxicology of, 422
- Spasms, as a complication of whooping-cough, 448
- Spatula, spear-headed, 347, 348
- Specific gravity of bile, 214
- blood, 114
- brain, 363
- kidney, 368
- liver, 367
- lung, 366
- pancreas, 369
- Specimens, general method of preserving, 343
- injected, 344
- Kaiserling's method of preserving, 343
- Littlejohn's method of preserving, 345
- macroscopical, examination of, 341
- microscopic examination of, xxii, p. 326
- natural color of, method of preserving, 344
- of peritoneal fluid from inoculated animals, 356
- urine from inoculated animals, 356
- various tissues from inoculated animals, 356
- preservation of, xxii, p. 326
- removed at autopsies, 6
- washing of, 342
- Spectroscopic examination of blood, 113, 418, 444
- Spermatic cord, diseases of, 212
- examination of, 14, 186, 212
- duct, 14
- Spermatozoa, 46, 417
- Sphenocephalic skull, 361
- Spina bifida, 68, 255
- Spinal arthritis, 264
- Spinal canal, method of examination of, xiv, p. 242
- column in postmortem, 440
- cord, appearance of, in anthrax, 440
- atrophy of, 253, 254
- diseases of, 68, 82, 122, 205, xv, p. 246; 291, 297, 307, 308, 310, 313, 404, 413, 430, 433
- examination of, 14, 243, 440
- hemorrhage of, 251
- in postmortem, 440
- measurements of, 364
- method of removing from body of baby, 277
- miliary tuberculosis of, 313
- opening of, 244
- preservation of, 244
- other diseases of, 452
- removal of, 7, 14, 242, 243, 244, 254, 274, 277, 440
- from back, 242
- front, 245
- in animals, 380, 389, 394
- study of, for removal, 243
- tuberculosis of, 252, 313
- tumors of, 255
- weight and dimensions of, 364
- curvatures, 269
- ganglia, 243
- membranes, hemorrhage of, 251
- nerves, 243
- Spindle-celled sarcoma in mediastinum, 146
- of lungs, 156
- Spiral douche, use of, to produce abortion, 406
- Spirals, Curschmann's, 144
- Spirochaetæ of Obermeier, 302, 351
- Spleen, 162
- absence of, 162
- amyloid degeneration of, 164, 307
- anæmic infarcts in, 163
- angiomata of, 164
- cancer of, 164
- changes in, in scarlet fever, 304
- coal-dust in, 163
- dimensions of, 369
- diseases of, 18, 66, 85, 117, 123, 124, 125, 127, 132, 144, 153, 161, 162, 163, 164, 181, 190, 217, 228, 291, 295, 299, 302, 305, 307, 308, 317, 323, 428, 431, 458
- echinococci cysts in, 164
- elastic tissue of, 163
- embolic infarct of, 163
- enlargements of, 62, 163, 305
- examination of, 14, 15, 162, 163, 442
- fixatives for, 340
- hyperæmia of, 163
- leukæmic, 164
- measurements of, 162, 369
- method of removing, 162
- normal color of, 163



- Spleen, of animals, removal of, 379, 380  
 inoculated animals, securing material from, 356  
 oligæmia of, 163  
 Pentastomum denticulatum of, 164  
 removal of, 15, 162, 280  
   in animals, 356, 379, 381, 382  
 rupture of, 303  
 sago, 164  
 sarcoma of, 164  
 tuberculosis of, 164, 311  
 tumors of, 116, 164  
 volume of, 369  
 weight of, 162, 360, 369
- Splenitis, acute, 163
- Spondylitis deformans, 263
- Sponges, 33
- Spontaneous rupture of heart, 129
- Spores, bacterial, longevity of, 356
- Spots, Tardieu's, 130
- Sprains, synonyms of, 463
- Sprue, 306
- Spurious aneurism, 137
- Stab cultures, 352
- Stain, trypan red, 325
- Stains, iodine, removal of, by ammonia, 286  
   on clothing, 54, 117  
   location of, to be marked, 54
- Staphylococci, 151  
   immunity to, 43
- Staphylococcus aureus, 160, 351
- Starvation, death by, indications of, 414
- Statistics, constant revision of, 3  
   morbidity and mortality, xxviii, p. 448
- Stature, abnormalities of, 56
- Status lymphaticus, 279
- Steel tape measure, 32
- Stegomyia, in yellow fever, 319
- Stenosis, 145  
   aortic, 133  
   laryngeal, 143  
   mitral, 133  
   of bronchi, 126  
     pulmonary valves, 133  
     trachea, 146  
     ureters, 204  
     uterus, 209  
   pyloric, 190
- Sterility in dwarfism, 56
- Sterilizing instruments, 28, 45, 352  
   syringes, 354
- Sternum, absence of, 68  
   and ribs, altered shape of, 94  
     examination of, 94  
   congenital luxation of, 266  
   diseases of, 68, 71, 72, 94, 246, 265, 266, 290  
   ensiform appendix of, 94  
   examination of, 68, 71, 94  
   erosion of, 94  
   fissure of, 68  
   fracture of, location of, 94  
   in carcinoma of breast, 94
- Sternum, marrow of, changes in, 94  
   pressure atrophy of, 94  
   regulations for removal of, 93  
   removal of, 93  
   restoration of, 285  
   tuberculous caries of, 94  
   tumors of, 274
- Stillbirths, 464
- Stillborn babes, hemorrhage in, 95  
   thymus gland of, 95
- Stomach and duodenum, of animals, removal of, 379  
   œsophagus, method of examining, 188  
   cancer of, 190  
   cancerous, removal of, during life, 19  
   capacity of, estimation of, 90  
   colloid cancer of, 190  
   contents of, examination of, 189  
     in peritoneal cavity, 87  
   dilatation of, 190  
   dimensions and weight of, 366  
   diseases of, 88, 90, 122, 151, 153, 191, 192, 217, 220, 224, 297, 299, 300, 318, 319, 419, 426, 427, 428, 430, 431, 432, 433, 444  
   examination of, 14, 15, 90, 188, 189, 190, 191, 192, 442, 443, 444  
   extenteration of, in animals, 381  
   gas in, post-mortem, 90  
   hemorrhage from, 192  
     petechial, from, 192  
   hour-glass contracture of, 90, 191  
   in animals, examination of, 383  
   in postmortem, 443  
   measurements of, 366  
   mucous membrane of, action of poisons on, 419  
   medullary cancer of, 190  
   of animals, removal of, 380  
   position of, in babe, 90  
   relative position to umbilicus, 17  
   removal of, 15, 188, 189, 443  
     in animals, 379, 381, 382, 393  
   scirrhus cancer of, 190  
   serous covering of, examination of, 90  
   size and situation of, 90  
   tuberculosis of, 312  
   tumors of, 90, 122, 190  
   ulcer of, synonyms of, 457  
     with perforation, 87  
   unusual articles found in, 189  
   weight of, 360
- Stomach-pump, 33
- Stomatitis aphthosa epizootica, 295
- Stone formations in kidney, 204  
   urine, 204
- Stools, appearance of, in cholera Asiatica, 293
- Strabismus, 73, 74
- Stramonium, toxicology of, 411, 414, 422

- Strangulation, 415  
 Streptococci, 151  
   immunity to, 43  
 Streptococcus, 295  
   intracellularis, 351  
   lanceolatus, 160, 351  
   pyogenes, 160, 351  
 Streptothrix actinomycosis, 290, 351  
 Stricture of œsophagus, 190  
   syphilitic, of bowel, 306  
   urethral, location of, in male, 70  
 Stroke culture, 352  
 Strongylus, 151  
   annatus, 138  
   gigas, 202  
 Strophanthus, toxicology of, 424  
 Strychnine, toxicology of, 411, 419, 421.  
   426, 435  
   effect on urine, 426  
   poisoning, post-mortem rigidity in, 51  
   test for, in urine, 435  
   toxicology of, 421, 426, 435  
 Struma aneurysmatica, 147  
   varicosa, 147  
 Subacute bronchitis, 144  
 Subcutaneous hemorrhages in plague, 302  
   inoculation of animals, 354  
 Subdiaphragmatic abscesses, 159  
 Sublamin, 335  
 Sublingual glands, 15, 366  
 Submaxillary glands, 80, 366  
 Submersion, accidental, 463  
 Subphrenic abscesses, 159  
 Sudden death, causes of, 407, 464  
   due to pressure of thymus, 96, 279  
 Suffocation, 415  
   death by, delays loss of vital warmth.  
   49  
   possible findings in, 415  
   post-mortem lividity in, 50  
 Suggestions, medicolegal. xxvi, p. 397  
 Suicide, 409  
   by asphyxia, 462  
   crushing, 463  
   cutting instruments, 462  
   drowning, 462  
   firearms, 462  
   hanging, 462  
   jumping from high places, 463  
   poison, 462  
   strangulation, 462  
   ingenious ways of, 410  
   methods of committing, 410  
   number of, increasing, 409  
   order of frequency of, 410  
 Sulphide of antimony, toxicology of, 418  
 Sulphonal, toxicology of, 421, 425  
 Sulphuric acid, toxicology of, 426  
 Sulphurous acid, toxicology of, 422  
 Sunstroke, 414  
   blood in, 414  
   lividity and putrefactive changes in.  
   414  
   Sunstroke, temperature in, 414  
   venous engorgement in, 414  
 Supernumerary fingers and toes, 68  
   see Accessory under organs.  
 Suppuration complicating smallpox, 448  
   due to tuberculous lesions, 311  
   of lung, 153  
   pleura, 156  
 Suppurative appendicitis, 171  
   bronchitis, 145  
   cholangitis, 219  
   exudates, odor, etc., of, 86  
   gastritis, 191  
   hepatitis, 219  
   periostitis, 271  
 Suprarenals, see Adrenals.  
 Surgery, practice of, at postmortems, 3  
 Surgical instruments in abdominal cavity,  
   88  
 Surra, 324  
 Surroundings, study of, 54  
 Suspended animation, cases of, 46  
 Sutures in restoring skull, 283  
 Suturing of heart muscle, 100  
 Sweating (miliary) fever, 449  
 Swine, postmortems on, 391  
 Symelus, 67  
 Sympathetics, 14, 16, 97, 276  
 Symptoms observed after the administra-  
   tion of the more common poisons, 421  
 Syncephalus, 67  
 Syncope, fatal, 21  
 Syncytioma malignum, 209  
 Syndesmosis, 266  
 Synonyms of causes of death, xxviii, p.  
   448  
 Synostosis, 266, 269  
   premature, 361  
 Synovial membrane, 261  
 Synovitis, catarrhal, 262  
 Synthetical organic poisons, 421  
 Syphilis, 351, 450  
   acquired, 306  
   as a cause of spontaneous abortion,  
   406  
   chancre of, 307  
   choroiditis due to, 307  
   congenital, 307  
   as cause of death, 408  
   cutaneous eruptions in, 307  
   finger-nails in, 307  
   germs of, 296  
   gumma of, 306  
   hepatic cirrhosis due to, 216  
   hereditary, 277, 306  
   initial lesion of, 306  
   iritis due to, 306  
   mucous patch of, 306  
   of arteries, 137  
   brain, 255, 308  
   circulatory system, 307, 308  
   cord, 308  
   gastro-intestinal tract, 308

- Syphilis of heart, 133  
 kidneys, 308  
 larynx, 309  
 liver, 309  
 lung, 309  
 lymph-vessels, 140  
 nervous system, 307  
 skin, 306, 307  
 testes, 310  
 vulva, 69  
 parotitis due to, 306  
 periostitis due to, 306  
 tertiary stage of, 307  
 ulcerating lesions of, 63  
 Syphilitic arthritis, 264  
 bone lesions, 72  
 cirrhosis of liver, 217  
 congenital, pancreatitis, 221  
 hydrocele, 417  
 inflammation of veins, 140  
 pancreatitis, 221  
 periarteritis of pancreas, 222  
 sore, 349  
 ulcers of œsophagus, 190  
 vagina, 211  
 Syringe, hypodermic, in inoculation of animals, 353  
 Koch's inoculation, 353  
 method of holding in intravenous inoculation of animals, 355  
 Syringes, sterilizing of, 354  
 used in inoculation of animals, 353, 354  
 careful treatment of, 354  
 Syringomyelia, 255, 264
- T**
- Tabs, dislocations common in, 262  
 dorsalis, 248, 306  
 mesenterica, 160  
 Table, dimensions of postmortem, 11  
 of approximate weight of internal organs, 360  
 changes in urine effected by poisons and medicines, 424  
 portable post-mortem, 9  
 showing difference between croupous and catarrhal pneumonia, 154  
 tuberculous and typhoid ulcers, 167  
 in grammes the mean weights of the brain, 362  
 Tables, construction of, 33  
 of weights and measures, 17  
 rotating, 33  
 weighing, 33  
 zinc-covered, wood, slate, iron, or glass, 33  
 Tænia, bothriocephalus latus, 121, 320  
 Tænia flavopunctata, 320  
 mediocanellatæ, 170  
 nana, 170  
 saginata, 320  
 solium, 320  
 Talipes calcaneus, 269  
 cavus, 269  
 equinus, 248, 269  
 planus, 269  
 valgus, 269  
 varus, 269  
 Tallqvist's blood-color scale, use of, 113  
 Tannin, toxicology of, 435  
 Tardieu's spots, 101, 130  
 Tar, toxicology of, 422  
 "Tea" cigarettes, poisoning therefrom, 412  
 Technic of exposing thoracic cavity, vii, p. 92  
 opening abdominal cavity, vi, p. 79  
 postmortem, 437  
 Teeth, 78  
 actinomycosis of, 290  
 anomalies of, 78  
 condition of, 78  
 deformities of, 257  
 diseases of, 78, 140, 251, 290, 308  
 examination of, 53, 78, 438  
 in pernicious anæmia, 121  
 special authority for removal of, in France, 6  
 syphilitic, 78, 308  
 tumors of, 78  
 Tellurium, toxicology of, 21, 424, 426  
 Temperature at which pathogenic bacteria grow best, 352  
 in cases of certain abdominal disorders, 49  
 cholera, 49  
 injury to nervous system, 49  
 sunstroke, 414  
 of animals, how taken, 353  
 rectal, after death, 49  
 rise of, after death, 49  
 Tenaculum, 32  
 Tendons, 14, 272  
 Tenosynovitis, acute, 272  
 chronic, 272  
 hæmatogenous, 272  
 simple, 272  
 tuberculous, 272  
 Terata anadidyma, 67  
 anakatadidyma, 67  
 katadidyma, 67  
 Teratoma of pleura, 156  
 Tertian malaria, 322  
 Tertiary stage of syphilis, 307  
 Test, antiserum, 118  
 Gmelin's, for bile, 120  
 Widal, for distinguishing human blood, 120  
 Testes, see Testicles.



- Testicles, abnormal position of, 69  
   abscess of, 211  
   absence of one or both, 68  
   atrophy of, 69, 211, 419  
   dimensions of, 370  
   diseases of, 68, 211, 300, 301, 307, 310, 313, 324, 419  
   duplication of, 69  
   ectopia of, 69  
   examination of, 14, 15, 68, 73, 186, 442, 444  
   external method of examining, 186  
   hypertrophy of, 69  
   hypoplasia of, 69  
   inflammation of, 211  
   injury of, 211  
   malformation of, 69  
   measurements of, 370  
   removal of, 185, 186  
   swelling of, 69  
   syphilis of, 211, 307, 310, 314  
   true abscesses in, 211  
   tuberculosis of, 211, 314  
   tumors of, 211  
   undescended, 69  
   weight of, 360, 370  
 Testimony, expert, 398, 399  
   in court, 17  
   medicolegal cases, 400  
   of a medical man in medicolegal cases, 400  
 Tetanus, 40, 310  
   antitoxin, its subdural use, 45  
   neonatorum, 310  
   post-mortem rigidity in, 51  
   precautions when feared, 45  
   synonyms of, 453  
   toxin of, toxicology of, 421  
 Tetranol, toxicology of, 425  
 Thawing of frozen bodies, 289  
 Thermocautery, 35  
 Thionin, 341  
 Thiosulphate of sodium, effect on urine, 424  
 Thoracic cavity, critical examination of, vii, p. 92; 440  
   contents in animals, removal of, 384, 393  
   of animals, exposure of, 376  
   duct, 14, 15, 37, 197, 198, 365  
   organs, ablation of, 382  
   in animals, dissection of, 384  
   technic of exposing, vii, p. 92  
 Thorax, appearance of, in anthrax, 291  
   emphysema of lungs, 149  
   method of injecting embalming fluid into, 288  
   opening, 92  
 Thread, to remove food from teeth, may poison, 412  
 Thrombi in chlorosis, 123  
   of heart, 137  
   tuberculous, 16  
 Thrombo-arteritis, 136  
 Thrombosis, 20, 136  
   and embolism, synonyms of, 455  
   as cause of death, 408  
   of brain, 250  
   dura, 233  
   portal vein, 195  
   pulmonary vein, 148  
   superior mesenteric artery, 137  
   veins, 139  
 Throttling and hanging, differentiation of, 415, 416  
 Thrush, 192, 209, 310  
 Thuja, toxicology of, 424  
 Thymol, effect on urine, 425  
 Thymus gland, abscess of, 95, 96  
   color of, 95  
   dimensions of, 370  
   diseases of, 70, 95, 123, 147, 181, 246  
   enlargement of, causing asphyxia, 279  
   examination of, 15, 95, 441  
   loses coloring matter when washed, 19  
   hemorrhage into, 95  
   in acromegaly, 95  
   adult, 95  
   certain pathologic conditions, 95  
   exophthalmic goitre, 95  
   myxoedema, 95  
   removal of, 15, 445  
   tumors of, 95, 96, 246  
   weight of, 95, 370  
 Thyroid gland, accessory, 96  
   anatomic changes in, 249  
   arsenic in, 249  
   atrophy of, in dwarfism, 36  
   degenerations of, 96  
   dimensions of, 370  
   diseases of, 56, 66, 70, 73, 96, 123, 146, 222, 246, 249, 455  
   enlarged, 66, 96  
   examination of, 15, 66, 96, 442  
   fixatives for, 341  
   hypertrophy of, 96  
   measurements of, 370  
   myxoedematous degeneration of, 96  
   pyramid of, 96  
   removal of, 15, 110  
   symptoms following removal of, 96  
   tumors of, 96, 146, 147, 246  
   weight of, 370  
 Thyroiditis in diphtheria, 96  
 Time after death before making an autopsy, 59  
   for body to skeletonize, 59  
   weighing organs, 357  
   of complete post-mortem cooling, 49  
 T-incision, 259

- Tinea favus, 461  
     tonsurans, trichophyton, 461  
 Tissues, for preservation, permission to  
     remove, 5  
     preservation of, xxii, p. 326  
 Toadstool, 434  
 Toluidin, toxicology of, 423  
 Toluyldiamin, effect on urine, 425  
 Tongue, atrophied (microglossus), 68  
     cleft, 68  
     deformities of, 68  
     diseases of, 67, 68, 111, 128, 140, 302,  
         310, 312  
     examination of, 15, 96, 111, 438, 442  
     frænum of, too short or too long, 68  
     hypertrophy of, 68  
     macroglossus of, 68  
     œsophagus, trachea, and adjacent  
         structures together, removal of, 110  
     removal of, 15, 37, 96, 110, 280, 442  
     tuberculosis of, 312  
     tumors of, 111, 140  
 Tonsils, angioma of, 143  
     diseases of, 124, 143, 294, 304, 307, 312  
     epithelioma of, 143  
     examination of, 15, 111, 442  
     fibroma of, 143  
     hypertrophied, 143  
     lymphoma of, 143  
     myoma of, 143  
     papilloma of, 143  
     removal of, 111  
     sarcoma of, 143  
     tumors of, 143  
 Tophi, 72, 128  
     murexoid test for, 73  
 Topographic examination of abdominal  
     cavity, vi, p. 79  
 Tow, for packing large cavities, 34  
 Toxic angina, see Diphtheria and Croup.  
     arthritis, 264  
     gastritis, 192  
     inanition, 418  
     purpura, 126  
 Toxicology, xxvi, p. 417 and the various  
     drugs themselves.  
     definition of, Robert's tables on, 421  
 Toxin of tetanus, 421  
     increases constitutional symptoms, 43  
 Trachea, calcareous and papillomatous ex-  
     crescences of, 146  
     diseases of, 97, 112, 144, 297  
     examination of, 14, 96, 112, 442, 445  
     fistula in, 144  
     in animals, dissection of, 385  
     injuries of, 112  
     removal of, 96, 110, 444  
     stenosis of, 146  
     tongue, œsophagus, and adjacent  
         structures together, removal of, 110  
 Trachoma, contagious granular lids, 454  
 Tract, genito-urinary, xi, p. 199  
     intestinal, gall-stones in, 170  
     Tract, intestinal, worms in, 170  
         respiratory, diseases of, 141  
 Traction aneurism, 138  
 Trade or occupation intoxications, 452  
 Trance, cases of, 46  
 Transudate, 20  
     how distinguished from exudate, 86  
 Transverse colon, malpositions of, 90  
 Traumatic aneurism, 138  
 Traumatism, 129, 155, 202, 218, 246, 463  
 Trays, enamelled, 33  
 Treatment of post-mortem wounds, iv, p.  
     38  
 Trephining engines, 29  
 Trichina spiralis, 83, 321, 373, 449  
     in tongue of cat, 344  
     usual location of, in muscles, 83  
 Trichosis vesicæ, 206  
 Tricuspid orifice of the heart, dimensions  
     of, 365  
     regurgitation, 133  
 Trigonocephalic skull, 361  
 Trional, toxicology of, 425  
 Triple monsters, 67  
     staining, 341  
 Trocar, 32, 65  
     curved, method of injecting thorax by  
         means of, 288  
     use of, in injecting brain cavity, 288  
 Tropacocaine, toxicology of, 426  
 Trophocephalic skull, 361  
 Tropical diseases, Manson's work on, 296  
     hepatitis, 219  
 True aneurism, 137  
 Trypanosoma, 121, 324, 373, 423  
 Trypan red, 325  
 Tsetse flies, 324  
 Tubal pregnancy, see Extra-uterine preg-  
     nancy.  
 Tubercle, bacterium, 310, 349  
     rubic, 298  
 Tuberculin, use of, in anatomic wart, 45  
 Tuberculosis, 310  
     abdominal, 450  
     aneurismal arteries in, 316  
     as cause of death, 408  
     bovine, 373  
     calcification in, 312  
     caseous, 311  
         nodules as lesions in, 316  
     cavities as lesions in, 316  
     cutaneous forms of, 318  
     diffuse, 311, 315  
     distribution of, in body, 311  
     enlarged bronchial glands as lesions  
         in, 316  
     general, caused by anatomical wart, 44  
     generalized, 450  
     human, 373  
         as differing from bovine, discus-  
             sion of, 373  
     hypertrophy of heart in, 316  
     lesions of, 311

- Tuberculosis, lymphatic glands in, 316  
 miliary, 16, 311, 313, 315, 317  
 distribution of, in body, 311  
 of adrenals, 180  
   brain and cord, 313  
   omentum, 159  
   pericardium, 69  
   peritoneum, 317  
   ureter, 182  
 modes of invasion of, 311  
 nasal, 141  
 of alimentary tract, 312  
   arteries, 137  
   bladder, 205, 314  
   bone, 273  
   brain and cord, 313  
   circulatory system, 314  
   genito-urinary system, 314  
   heart, 134  
   intestines, 312  
   kidneys, 314  
   larynx, 315, 449  
   liver, 313  
   lung, 315, 449  
   lymph-vessels, 140  
   mammary gland, 317  
   meninges, 450  
   mouth, 83, 312  
   oesophagus, 190, 312  
   peritoneum, 317  
   serous membranes, 317  
   skin, 318  
   stomach, 312  
   testes, 314  
   thoracic duct, 198  
   ureters, 314  
   urethra, 70  
   vein, 140  
 pneumonic areas as lesions in, 316  
 thickening of pleura, as lesion in, 316  
 ulcerative, 315  
   of peritoneum, 317  
 Tuberculous affections, 450  
   arthritis, 264, 265  
   bronchitis, 145  
   coxalgia, 72  
   endometritis, 210  
   hydrocele, 417  
   knee-joint, 72  
   lesions, fate of, 311  
   periostitis, 271  
   spinal curvature, 72  
   tumors of brain, 313  
   ulcerations, 311  
   ulcers, 167  
     distinguished from typhoid, 168  
 Tubes, infectious material used in, 352  
   inoculated, cultivation of, 352  
   inoculation of, 352  
 Tumors, as cause of death, 408  
   deformity, 72  
   cells of, in blood, 120  
   dental, 78  
 Tumors, erectile, of arteries, 137  
   for the different forms of tumors, see  
   the organs or parts themselves.  
 lymph, 140  
 of adrenals, 180  
   appendix, 172  
   arteries, 137  
   bladder, 205, 206  
   bone, 273  
   brain, 255, 313  
   breast, 81  
   bronchi, 146  
   cord, 255  
   dura, 230  
   eye, 75  
   Fallopian tubes, 208  
   gall-bladder, 195  
   heart, 134  
   intestines, 171  
   kidney, 179, 204  
   larynx, 143  
   liver, 214, 219  
   lungs, 155  
   muscles, 84  
   nasal passages, 141  
   ovary, 209  
   pancreas, 196  
   peritoneum, 159  
   salivary glands, 260  
   skin, 64  
   spleen, 164  
   stomach, 190  
   testicles, 211  
   uterus, 209  
   vagina, 211  
   veins, 140  
   retrobulbar, 258  
   sections of, for preservation, 327  
 Turmeric paper, 34  
 Turpentine as a cause of amyloid degeneration, 418  
   toxicology of, 418, 424  
 Twine, 33  
 Twin pregnancy, history of, 279  
 Types of skull due to premature synostosis, 361  
 Typhoid bacilli, 169, 205, 350  
   fever, 318  
     as cause of death, 408  
     complications of, 448  
     intestinal lesions of, 318  
     mesenteric lymphatic glands in, 89, 318  
     post-mortem rigidity in, 51  
     synonyms of, 448  
   ulcers, 165, 167  
     cause of artificial anus, 20  
     congestion in, 89  
     how distinguished from tuberculous ulcers, 168  
     in Meckel's diverticulum, 20  
     of oesophagus, 190  
 Typhus fever, 351



## U

- Ulceration of larynx, 143  
 Ulcerations, tuberculous, 311  
 Ulcerative colitis, 175  
   endocarditis, 132  
   phthisis, 311, 315  
   tuberculosis, 315  
     of peritoneum, 317  
 Ulcers, diphtheritic, of vulva, 69  
   duodenal, 164  
   gastric, situation of, 189  
   intestinal, 167  
   of œsophagus, 190  
     stomach, 169, 457  
   peptic, 169  
     of œsophagus, 190  
   perforating, of foot, 248  
   syphilitic, of œsophagus, 190  
     of vagina, 211  
   tuberculous, 167  
     of œsophagus, 190  
   typhoid, 167  
     and tuberculous, table showing  
       differences between, 167  
     causing artificial anus, 20  
     in Meckel's diverticulum, 20  
     of œsophagus, 190  
 Umbilical cord, 128, 359, 405, 406, 444  
   hemorrhage of, 128  
   length of, 359  
 Umbilicus, 66, 68, 219, 276, 310  
 Uncinariasis, 321  
 Undertakers, duties of, at postmortem, 9  
 Ungentum Credé as local antiseptic, 43  
 Unna's alkaline methylene blue, 341  
   orcein stain, 341  
 Urachus, allantoic cysts of, at navel, 66  
   persistence of, 68  
 Uræmia, 64, 117, 422  
   as cause of death, 407  
   blood changes in, 117  
 Uranium, salts of, toxicology of, 425  
 Urate, ammonium, 207  
   sodium, 207  
 Ureters, accidental tying of, 182  
   cysts of, 204  
   dimensions of, 368  
   diseases of, 127, 182, 200, 204, 224, 314  
   examination of, 14, 15, 175, 176, 181  
   miliary tubercles in, 182  
   parasites in, 205  
   polyps of, 204  
   removal of, 176, 184  
     in animals, 381  
   situation of, 175  
   stenosis of, 204  
   tumors of, 181, 209  
 Urethra, abnormal openings of, 69  
   absence of, 69  
   angioma of, 70  
   atresia or partial phimosis of, 69  
   caruncle of, 70  
   cleft, 69  
     condyloma of, 70  
     deformity of, 69  
     dimensions of, 368  
     discharge from, 69  
     diseases of, 69, 205, 297, 459  
     epithelioma of, 70  
     examination of, 14, 15, 39, 89, 181,  
       209, 442  
     fibroma of, 70  
     inflammation of, 69  
     measurements of, 368  
     occlusion of, 69  
     removal of, 185, 186  
     sarcoma of, 70  
     stricture of, location of, in male, 70  
     tuberculosis of, 70  
     tumors of, 69  
 Urethritis, 69  
 Uric acid, 207  
   in blood, in gout, 128  
   infarct of kidney, 201  
 Urinary salts, precipitates of, mistaken  
   for pus, 178  
   tract, calculi of, 459  
   lesions in, 182  
 Urine, black, causes of, 206  
   frog test for strychnine in, 183  
   of new-born, study of, 278  
   specimen of, from inoculated animals,  
     356  
   stone formation in, 207  
   typhoid bacilli in, 205  
 Urinometer, 33  
 Urobilin, 214  
 Urobilinogen, 214  
 Uropygium, 394  
 Urtica, toxicology of, 422  
 Usual causes of death, xxviii, p. 448  
 Utensils in postmortems on large domes-  
   tic animals, 374  
 Uterine hemorrhage, non-puerperal, syno-  
   nyms of, 459  
   tumors, 209  
 Uterus, atresia of, 209  
   atrophy of, 209  
   bilocularis, 209  
   bicornis, 209  
     duplex, 209  
   bipartitus, 209  
   congenital abnormalities of, 209  
   cordiformis, 209  
   dermoid cysts of, 209  
   didelphys, 209  
   dimensions of, 371  
   diseases of, 128, 201, 209, 210, 407  
   displacements of, 209  
   double, 209  
   examination of, 14, 15, 69, 186, 442  
   foreign bodies in, 210  
   hemorrhages of, 209  
   hypertrophy of, 209  
   hypoplasia of, 209

- Uterus, infantile, 209  
   infarcts of, 209  
   measurements of, 371  
   moles of, 209  
   other diseases of, 459  
   polyps of, 209  
   removal of, in animals, 386  
   rudimentary, 209  
   rupture of, 209  
   septus duplex, 209  
   situation of, 209  
   stenosis of, 209  
   subseptus, 209  
   thrush fungi in, 209  
   tuberculosis of, 209  
   tumors of, 209  
   unicollis, 209  
   unicornis, 209  
   weight of, 371  
     twenty-four hours after de-  
     livery, 210
- V**
- Vagina, abscess of, 211  
   diseases of, 210, 211, 297  
   erosions of, 211  
   examination of, 15, 39, 210, 442  
   hæmatocele of, 211  
   hernia of, 211  
   poisons in, 211  
   prolapse of, 211  
   removal of, 183  
     in animals, 386  
   syphilitic ulcers of, 211  
   tumors of, 211  
   vegetations of, 211  
 Vaginal hysterectomy, 281  
 Vaginitis, gonorrhœal, 297  
 Valsalva, sinus of, aneurism of, 138  
 Valves, diseases of, 130, 132, 133, 142, 153,  
   201, 210, 218, 247, 291, 297, 301, 303,  
   308, 408  
   examination of, 103  
     in animals, 385  
   measurements of, 103  
   mitral and pulmonary, situation of,  
     107  
     how remembered,  
     107
- Vapor fixation, 336  
 Vapors of nitric acid, etc., inhaled, toxicology of, 423  
 Varicella (chicken-pox).  
   Varicocele, 140  
   Varicose veins of vulva, rupture of, 69  
 Variola, see Smallpox.  
 Varioloid, see Smallpox.  
 Vasa deferentia, 224  
 Vascular glands, changes in, in acromegaly, 56  
   nephritis, 199  
   system in alcoholism, 427
- Vegetable acids, toxicology of, 426  
   irritant poisons, 421  
   toxic poisons, 421  
 Vegetations, vaginal hypertrophic, 211  
 Veins, azygos, 15  
   diseases of, 136, 137, viii, p. 139; 219,  
     224, 416, 455  
   examination of, 440  
   fibrous endophlebitis in, 140  
   hemorrhoids of, 140  
   phlebitis of, 139  
   phleboscrosis of, 140  
   primary tumors in, 140  
   syphilitic inflammation of, 140  
   thrombosis of, 139  
   tuberculosis of, 140  
   tumors of, 140  
   varicocele of, 140  
 Velum palati, 15, 96, 257, 442  
 Vena cava, 14, 15, 101, 156, 197, 365, 442  
   in animals, 381  
 Venereal warts of vulva, 69  
 Venous engorgement in sunstroke, 414  
   naevus, 137  
 Ventilating shaft, 11  
 Ventricles of brain, examination of, 439  
   heart, examination of, 102, 441  
   thickness of, 105, 106  
 Veratrine, toxicology of, 419, 421  
 Veratrum viride, post-mortem rigidity in  
   poisoning by, 51  
 Vermiform appendix, dimensions of, 366  
   in hernia, 20  
   situation of, 20  
   weight of, 366  
 Vertebral column, 67, 72, 198, 248, 253,  
   254, 255, 265, 290, 303, 311, 402, 440, 442  
 Vertical transverse incision of brain, 239  
 Vesical calculi, 206  
 Vesicles, seminal, 15, 372, 442  
 Vesicular emphysema of lungs, 150  
 Veterinary postmortems, 373  
 Viability of child, how determined, 403  
 Vibrio cholerae, 351  
 Vienna method of exenteration of ani-  
   mals, 382  
 Violence as a cause of death, 403, 407, 408  
   external, forms of, 408, 463  
 Violent death, 408  
 Virchow's method of determining bulk, 18  
   examining heart, 101  
   sectioning brain, 235  
 Virus of scarlet fever, effect on urine, 424  
 Vise, 32  
 Vital warmth, loss of, 48, 49  
 Vocal cords, 73, 111, 142, 256, 309  
 Volume of brain, 363  
   heart, 365  
   liver, 367  
   spleen, 369  
 Volvulus, 90, 139  
 Vomit, suspicious undissolved foreign  
   bodies in, 418

Vulva, absence of, 69  
 acne of, 69  
 aphthæ of, 69  
 atrophy of, 69  
 carcinoma of, 69  
 chancre of, 69  
 chancroid, 69  
 chondroma of, 69  
 diphtheritic ulcers of, 69  
 diseases of, 69, 70  
 eczema of, 69  
 elephantiasis of, 69  
 examination of, 15, 68, 70  
 fibroma of, 69  
 gumma of, 69  
 hæmatoma of, 69  
 herpes of, 69  
 hypertrophy of, 69  
 imperforate, 69  
 injuries of, 69  
 karorosis of, 69  
 lacerations of, 69  
 lipoma of, 69  
 lupus of, 69  
 mucous patch of, 69  
 myoma of, 69  
 œdema of, 69  
 sarcoma of, 69  
 syphilis of, 69  
 tumors of, 68  
 varicose veins of, rupture of, 69  
 venereal warts, 69  
 Vulvitis, gonorrhœal, 69  
 purulent, 69

**W**

Waiting-room, adjoining mortuary, 11  
 Walls, gastric, atony of, 191  
 Warm abscess, synonyms of, 461  
 Warmth, vital, rapidity of loss of, 48, 49  
     delayed, 48, 49  
     hastened, 48, 49  
 Wart, anatomic, 44, 318  
     cause of general tuberculosis, 44  
     treatment of, 44, 45  
     syphilitic, 307  
     venereal, of vulva, 69  
 Washing tissues for preservation, 342  
 Water, effect of, on lowering temperature  
     of corpse, 49  
     presence of, in body, 360  
     use of, in cleansing parts, 10  
 Waxy degeneration of voluntary muscles  
     in typhoid fever, 318  
 Webbed fingers and toes, 68  
 Weigert's stain, 341  
 Weighing, proper time for, 357  
 Weight, American insurance standard of,  
     358  
     and measurement of child at birth, 358  
         fœtus, 359  
     nutrition, 59

Weight, approximate, of internal organs,  
     360  
     average, of brain stem, 362  
         of cerebellum, 362  
         cerebrum, 362  
         new-born, 358  
 avoirdupois, 357  
 greatest, attained by man, 358  
 how expressed, 17  
 maximum, of encephalon, 361  
 of adrenal bodies, 369  
     arachnoid and pia, 462  
     bladder, 368  
     body, by percentage, 360  
     breasts at birth, 372  
         during lactation, 372  
         of adult, 372  
     heart, 364  
     kidney, 367  
     large intestine, 366  
     liver, 366  
     lungs, 365  
     maxillary glands, 366  
     new-born, 358  
     œsophagus, 366  
     ovaries, 370  
     pancreas, 369  
     pia and arachnoid, 462  
     pineal gland, 363  
     pituitary gland, 363  
     placenta, 359  
     prostate, 372  
     small intestine, 366  
     spinal cord, 364  
     spleen, 162, 369  
     stomach, 366  
     supernumerary livers, 367  
     suprarenals, 369  
     testes, 370  
     thymus gland, 370  
     thyroid gland, 370  
     uterus, 371  
     vermiform appendix, 366  
 Troy, 357  
 Weil's disease, 458  
 Wharton, gelatin of, 276  
 Whetstone, 33  
 White swelling, synonyms of, 450  
 Whitish discoloration of mucous mem-  
     branes by corrosive alkalies, 423  
 Whooping-cough, 351, 448  
 Widal test, 169  
     for distinguishing human blood,  
         120  
     how made at postmortem, 319  
 Williams case, 420  
 Winslow's test for respiratory movement,  
     47  
 "Wish-bone," 394  
 Wooden tongue, 290  
 Woodhead's method for examining ear,  
     258  
 Worms, intestinal, in peritoneal cavity, 88




Wounds, iv, p. 38  
 character of, 408  
 gunshot, in violent death, 408  
 most liable to become infected, 43  
 of hands protected before postmortem, 38  
 pericardium and heart, 99  
 post-mortem, character of, 41  
 how caused, 42  
 results of, 41  
 treatment of, iv, p. 38  
   by antiseptic dressing, 45  
     bleeding, 43  
     bromin, 43  
     carbolic acid, 43  
     caustics, 43  
     cautery, 43  
     Crédé's ointment, 44  
     flax and poultices, 44  
     glacial acetic acid, 43  
     hemorrhage, as a therapeutic measure, 43  
     Hume's intravenous injection, 44  
     intravenous injections, 44  
     iodin, 44  
     silver nitrate solution, 44  
     sucking of, 43  
 virulence of, explained, 42  
 virulent kinds of, 41  
 'right's electrical engine, 30  
 'risberg's ganglion, 276

Xanthoma, 64  
 Xerosis bacillus, 350  
 Xiphopagus, 67  
 X-rays, effect of, on anatomic wart, 45  
   guinea-pigs, 261  
   photographic plates, 261  
 in arsenical poisoning, 429  
   examination of bones and joints, 261  
 medicolegal uses of, 413  
 no death yet attributed to, 413  
 sterility caused by, 211, 414  
 use of, in detecting movement of heart and lungs, 47

Yaws, 296  
 Yellow atrophy, acute, of liver, 213  
   fever, 319, 351, 373, 449  
   lesions in, 319  
   mosquito in, 319  
   organisms of, 351

Zenker's fluids, 333, 334, 338  
 Zinc salts, toxicology of, 423  
   chlorid, for hardening brain, 241



—

...

...

...

...



LANE MEDICAL LIBRARY

To avoid fine, this book should be returned on  
or before the date last stamped below.

FEB 4 1913

JAN 5 1914

FEB 21 1914

MAY 2 1916

NOV 1 1916

1/13-17

APR 29 '24

MAR 10 1937

FEB 23 1946

# PLATE III



J57  
C368  
1905

Cattell, Henry W.  
Postmortem pathology  
37520

NAME	DATE DUE
Dickson Nov. 25, 12	FEB 4 1913
McPherson	FEB 4 1913
T. T. Shea	Feb 8, 1913
Mr. Conley	MAR 12 1913
W. E. Key	APR 21 1913
E. Owen	May 25, 1916
E. Owen	May 25, 1916
Kattalburg	APR 4 1916
Kattalburg	APR 4 1916
A. J. Barocchi	APR 15 1917
R. H. Gutter	APR 15 1917
S. S. Paxton	APR 25 1917
Sam Zand	SEP 1 1917
Aug G. Zand	SEP 1 1917
Jawed	SEP 1 1917

